

POTENTIAL AMELIORATIVE ROLE OF N-ACETYLCYSTEINE AGAINST TESTICULAR DYSFUNCTION INDUCED BY TITANIUM DIOXIDE IN MALE ALBINO RATS

^{1,2}Ahmed Farag El-Kirdasy, ³Mohamed Abdo Nassan, ^{4,5}Ahmed Abdel Aziz Baiomy,
^{6,7}Tamer Ahmed Ismail, ^{6,8}Mohamed Mohamed Soliman and ^{6,9}Hossam Fouad Attia

¹Department of Biochemistry, College of Veterinary Medicine, Sadat City University, Egypt

²Department of Biomedical Sciences, College of Pharmacy, Shaqraa University, Saudi Arabia

³Department of Pathology, Faculty of Veterinary Medicine, Zagazig University, Egypt

⁴Department of Biotechnology, Faculty of Applied Medical Sciences, Turabah, Taif University, Saudi Arabia

⁵Department of Zoology, Faculty of Science, Cairo University, Egypt

⁶Department of Medical Laboratory, Faculty of Applied Medical Sciences, Turabah, Taif University, Saudi Arabia

⁷Department of Physiology, Faculty of Veterinary Medicine, Zagazig University, Egypt

⁸Department of Biochemistry,

⁹Department of Histology and Cytology,

Faculty of Veterinary Medicine, Benha University, Egypt

Received 2013-12-10; Revised 2013-12-22; Accepted 2013-12-31

ABSTRACT

In this study, we examined the ameliorative action of N-Acetylcysteine (NAC) against Titanium Dioxide (TiO₂) induced testicular degeneration in albino rats. Adult male albino rats were given saline as a control group, TiO₂ (1200 mg kg⁻¹ BW), NAC (100 mg kg⁻¹ BW) and co-treatment of NAC and TiO₂ as a protective group for 3 months. Testicular tissues were extracted for changes in testicular gene expression and histopathology. Administration of TiO₂ significantly increased mRNA expression of IL-6 and TNF- α that are normalized by NAC administration. TiO₂ administration down regulated Glutathione-S-Transferase (GST) while increased B-cell Lymphoma2 (Bcl2) expressions. Co-administration of rats by NAC together with TiO₂ normalized changes in GST and Bcl2 expression. Expression of steroidogenesis related genes [Androgen Binding Protein (ABP), 17 β -Hydroxysteroid Dehydrogenase (17 β -HSD), cytochrome P450 17A (CYP17 α) and aromatase] showed down regulation in TiO₂ administered groups and normalized when NAC given together with TiO₂. Moreover, TiO₂ induced toxicity in testes that accompanied by degeneration in seminiferous tubules with congestion, oedema and cell disruption that are partially normalized by co-administration of NAC with TiO₂. In conclusion, the present findings confirmed the beneficial effect of NAC to prevent apoptosis of sertoli cells and testicular dysfunction induced by TiO₂ in male albino rats.

Keywords: Titanium Dioxide, N-Acetylcysteine, Steroidogenesis Related Enzymes, Gene Expression, Testicular Dysfunction and Apoptosis

1. INTRODUCTION

Titanium Dioxide (TiO₂) is widely used as an additive, in paints, in sunscreens and in cosmetic creams as well as in the environmental decontamination of air,

water and soil by the destruction of pesticides (Fisher and Egerton, 2001; Kaida *et al.*, 2004; Choi *et al.*, 2006; Medina *et al.*, 2007). The degree of cellular damage and oxidative stress of nanoparticles are related to the particle size and its chemical composition (Hoet *et al.*,

Corresponding Author: Mohamed Mohamed Soliman, Department of Biochemistry, Faculty of Veterinary Medicine, University, Egypt Tel: 00966501531640; 002012-3722404

2004). The smaller the particles of TiO₂, the more reactivity, effectivity and toxicity (Oberdorster, 2006). Oxidative stress is one of the most important toxicity mechanisms in body tissues as in lung, kidney, testis, brain and spleen (Li *et al.*, 2010; Sang *et al.*, 2012). Higher doses of TiO₂ nanoparticles (324-2592 mg kg⁻¹ bw) are accumulated in spleen, liver, kidney, lung and testis (Chen *et al.*, 2009). Upon exposure to TiO₂, pathological lesions occur in the liver, spleen, kidneys, testis and brain (Shi *et al.*, 2013; Jia *et al.*, 2013). Consequently disruption in organ function occur and alteration in inflammatory and defensive mechanisms of body. Testis is rich in polyunsaturated fatty acids which is a main target of Reactive Oxygen Species (ROS). Recent studies have shown that ROS-mediated oxidative stress can cause cellular testis damage (Turner and Lysiak, 2008). ROS are scavenged by antioxidant defense systems including reduced glutathione.

As known, N-Acetylcysteine (NAC) is an antioxidant and free radical scavenger, used extensively in conditional nutrient (Moschou *et al.*, 2008). NAC acts as a cysteine donor (precursor of de novo GSH synthesis) and maintains or even increases the intracellular levels of glutathione, a tripeptide which protects cells from toxins and free radicals (Stanislowski *et al.*, 2000; 2003; Van de Poll *et al.*, 2006; Sadowska *et al.*, 2007; Atkuri *et al.*, 2007). NAC free thiol group is capable of interacting with electrophilic groups of ROS (Sen and Packer, 2000). It has a protective effect for a range of cell types, including cell death in testis induced by oxidative stress (Cay *et al.*, 2006). Zafarullah *et al.* (2003) reported that cell growth and survival rate increased in response to ROS-induced injuries that lead to growth arrest and apoptosis. Thus, NAC can restore the imbalance between pro-oxidant and antioxidant systems during oxidative stress. NAC is safe and well tolerated when administered orally but has documented risks with intravenous administration (Dodd *et al.*, 2008). Rarely, it can cause rashes, fever, headache, drowsiness, low blood pressure and liver problems in higher doses (Dodd *et al.*, 2008).

Cytokines are proteins secreted by most cell types (Feghali and Wright, 1997). They are pharmacologically active, exhibiting both beneficial and pathologic effects on the target cells. They are either pro or inflammatory or anti-inflammatory proteins secreted to maintain body defensive mechanism. Imbalanced expression of cytokines has been implicated in the progression of many diseases (Arend and Gabay, 2004). Their expression reflect the immune and health state of the body. In this

study, we investigated the effects of NAC on anatase nano-TiO₂ induced alterations in gene expression of steroidogenesis, reduced Glutathione (GST), testicular inflammatory and apoptotic activity (IL-6 and TNF- α and Bcl2 respectively) and testis histopathology and possible protection by N-acetylcysteine in male albino rats. The enzymes of steroidogenesis examined are Androgen Binding Protein (ABP), 17 β -hydroxysteroid dehydrogenase (17 β -HSD), Cytochrome P450 17A (CYP17 α) and aromatase.

2. MATERIALS AND METHODS

2.1. Materials and Animals

Ethidium bromide, agarose and TiO₂ anatase form, particle size (25-70 nm) were purchased from Sigma Aldrich Co., MO, USA. The male albino rats were purchased from Egyptian Co for Experimental Animals Import, Helwan, Cairo, Egypt. Gum acacia and NAC were obtained from El-Nasr Co., Cairo, Egypt. Cytokines primers were from MACROGEN, Seoul, Korea. Twenty four male albino rats weighting 150-200 g were obtained from the Animal House in Faculty of Veterinary Medicine, Benha University. Rats were given free access to food and water with 12h/12h dark light cycle. All animals were left one week for adaptation. They were housed in separate well-ventilated cages, under standard conditions.

2.2. Experimental Design

Twenty four rats were divided into 4 groups (6 rats per group). Group I (control group), were kept under standard conditions, fed on balanced diet for 3 months. Group II (NAC treated group) received 100 mg kg⁻¹ body weight NAC once daily by gastric gavage for 3 months. Group III (TiO₂ treated group) received 1200 mg kg⁻¹ body weight TiO₂ by gavage (1/10 LD 50) in 1mL of 5% gum acacia solution as a solvent once daily for 3 months. The dose of TiO₂ was used based on studies of (Wang *et al.*, 2007). Group IV (NAC+TiO₂) received 100 mg kg⁻¹ body weight NAC one hour before TiO₂ by gavage once daily for 3 months. At the end of the experiments, the rats were sacrificed. Blood and testis were taken for histopathology and gene expression.

2.3. RNA Extraction and cDNA Synthesis

For preparation of total RNA, testis (approximately 100 mg per sample) were collected from rats, flash

frozen in liquid nitrogen and subsequently stored at -70°C in 1 mL Qiazol (QIAGEN Inc., Valencia, CA). Frozen samples were homogenized using a Polytron 300 D homogenizer (Brinkman Instruments, Westbury, NY). Then 0.3 mL chloroform were added to the homogenate. The mixtures were shaken for 30 sec followed by centrifugation at 4°C and 12,500 rpm for 20 min. The supernatant layer was transferred to a new set of tubes and an equal volume of isopropanol was added to the samples, shaken for 15 sec and centrifuged at 4°C and 12,500 rpm for 15 min. The RNA pellets were washed with 70% ethanol, briefly dried up then, dissolved in Diethylpyrocarbonate (DEPC) water. The prepared RNA integrity was checked by electrophoresis. RNA concentration and purity were determined spectrophotometrically at 260 nm. The ratio of the 260/280 optical density of all RNA samples was 1.7-1.9.

For synthesis of cDNA, mixture of 2 μg total RNA and 0.5 ng oligo dT primer in a total volume of 11 μL sterilized DEPC water was incubated in the PeX 0.5 thermal Cycler (Thermo Electronic Corporation, Milford, Ma) at 65°C for 10 min for denaturation. Then, 4 μL of 5X RT-buffer, 2 μL of 10 mM dNTPs and 100 U Moloney Murine Leukemia Virus (M-MuLV) Reverse Transcriptase (SibEnzyme Ltd. Ak, Novosibirsk, Russia) were added and the total volume was completed up to 20 μL by DEPC water. The mixture was then re-incubated in the thermal Cycler at 37°C for 1 h, then at 90°C for 10 min to inactivate the enzyme.

2.4. Semi-Quantitative PCR Analysis

Specific primers for tested genes (Table 1) were designed using Oligo-4 computer program and synthesized by Macrogen (Macrogen Company, GAsa-dong, Geumcheon-gu. Korea). PCR was conducted in a final volume of 25 μL consisting of 1 μL cDNA, 1 μL of 10 picomolar (pM) of each primer

(forward and reverse) and 12.5 μL PCR master mix (Promega Corporation, Madison, WI) the volume was brought up to 25 using sterilized, deionized water. PCR was carried out using a PeX 0.5 thermal Cycler with the cycle sequence at 94°C for 5 min one cycle, followed by different cycles each of which consisted of denaturation at 94°C for one minute, minute with additional final extension at 72°C for 7 min. As a reference, expression of Glyceraldehyde-3-Phosphate Dehydrogenase (G3PDH) mRNA was detected by using specific primers (Table 1). PCR products were electrophorized on 1% agarose gel (Bio Basic INC. Konrad Cres, Markham Ontario), stained with ethidium bromide in TBE (Tris-Borate-EDTA) buffer. PCR products were visualized under UV light and photographed using gel documentation system. The intensities of the bands were quantified densitometrically using ImageJ software (<http://imagej.en.softonic.com/>).

2.5. Histopathological Examination

The testes of male albino rats were collected at the end of experiment from tested groups. The samples were fixed in Bouin's solution, then dehydrated in ascending grades of alcohols, cleared in xylene and embedded in paraffin. The samples were casted, then sliced into 5 μm in thickness and placed onto glass slides. The slides were stained by general and specific stains (Wilson and Gamble, 2008). PAS and Masson's Trichrome reactions were carried out based on protocols of (Kiernan, 1999; Carson, 1990) respectively.

2.6. Statistical Analysis

Results are expressed as means \pm S.E of 5 different rats per each group. Statistical analysis was done using ANOVA and Fischer's post hoc test, with $p < 0.05$ being considered as statistically significant.

Table 1. PCR conditions for rat cytokines and spermatogenesis genes

Gene	Forward primer	Reverse primer	PCR cycles and Ann. Temp.
IL-6 (450bp)	5'-AGTTGCCTTCTTGGGACTGATGT-3'	5'-TGCTCTGAATGACTCTGGCTTTG-3'	30 cycles, 57°C 1 min
TNF- α (256 bp)	5'-CCACCACGCTCTTCTGTCTAC-3'	5'-ACCACCAGTTGGTTGTCTTTG-3'	30 cycles, 58°C 1 min
Bcl2 (350 bp)	5' TCC ATT ATA AGC TGT CAC AGA GG -3'	5'-- GAA GAG TTC CTC CAC CAC C -3'	30 cycles, 55°C 1 min
GST (575 bp)	5'-GCTGGAGTGGAGTTGAAGAA-3'	5'-GTCCTGACCACGTC AACATAG -3'	33 cycles, 55°C 1 min
ABP (195 bp)	5'-CCATTCCTCCTTTGAGTTTCGA-3'	5'-CAGCTCCACCCGGGTGC-3'	35 cycles, 65°C 45 sec
Aromatase (142 bp)	5'-GCCTGTCGTGGACTTGGT -3'	5'-GGTAAATTCATTGGGGTTGG -3'	33 cycles, 57°C 1 min
CYP17 α (138)bp	5'-GCTAACATTGACTCCAGCATTGG-3'	5'-GAAGCGCTCAGGCATAAACC-3'	33 cycles, 60°C 1 min
17 β -HSD (114 bp)	5'-CTCTGGGCACTGCATCAC-3'	5'-CAAGTAACTCTGCGTGGGT-3'	33 cycles, 53°C 1 min
GAPDH (309 bp)	5'-AGATCCACAACGGATACATT-3	5-TCCCTCAAGATTGTCAGCAA-3'	25 cycles, 52°C 1 min

3. RESULTS

3.1. IL-6 and TNF- α Expression in Testis after TiO₂ and NAC Administration in Male Albino Rats

Administration of TiO₂ to rats for 3 months increased IL-6 and TNF- α expression in testis and their expression was normalized in rats administered NAC together with TiO₂ (Fig. 1a and b). NAC administration to rats alone has no effect on IL-6 and TNF- α expression.

3.2. Bcl2 and GST Expression in Testis after TiO₂ and NAC Administration in Male Albino Rats

To test the antiapoptotic and antioxidant effect of NAC, we tested the expression of Bcl2 and GST after administration of TiO₂ or NAC either alone or together. Figure 1c and d, show that TiO₂ increased Bcl2 expression and decreased GST expression and administration of NAC together with TiO₂ normalized the increase in Bcl2 expression and the decrease in GST expression.




Fig. 1. RT-PCR analysis of IL-6, TNF- α , Bcl2 and GST expression after administration of either NAC or TiO₂ alone or together in Albino rats. NAC and TiO₂ were administered for 3 months as described in materials and methods. RNA was extracted and reverse transcribed (3 μ g) and RT-PCR analysis was carried out for of IL-6, TNF- α , Bcl2 and GST genes. Densitometric analysis was carried for 3 different rats. *p<0.05 Vs control while # p<0.05 Vs NAC group and \$ p<0.05 Vs TiO₂

3.3. Expression of Steroidogenesis Regulating Enzymes in Testis of Albino Rats after TiO₂ and NAC Administration

To test the effect of the pollutant on genes of testosterone biosynthesis, rats were treated with and/or without TiO₂ in presence or absence of NAC. The expression of ABP, 17β-HSD, CYP17α and aromatase were examined using RT-PCR analysis. As seen in **Fig. 2a**, TiO₂ induced significant decrease in ABP expression but in presence of NAC the suppression was normalized and increased. The down regulation in ABP

expression was around 55% of normal expression. Also, NAC inhibited TiO₂ induced 70% down regulation in aromatase expression that normalized in presence of NAC (**Fig. 2b**). Next, we examined the expression of CYP17α that plays a role as a rate-limiting enzyme in the biosynthesis of steroid hormones. TiO₂ induced 43% down regulation in CYP17α expression in testis. Presence of NAC with TiO₂ significantly normalized the expression of CYP17α (**Fig. 2c**). Next, we tested the expression of 17β-HSD, the enzyme responsible for conversion of androstenedione to testosterone.




Fig. 2. RT-PCR analysis of steroidogenesis related gene expression after administration of either NAC or TiO₂ alone or together in Albino rats. NAC and TiO₂ were administered for 3 months as described in materials and methods. RNA was extracted and reverse transcribed (3 μg) and RT-PCR analysis was carried out for of ABP, aromatase, CYP17α and 17β-HSD genes. Densitometric analysis was carried for 3 different rats. *p<0.05 Vs control while # p<0.05 Vs NAC group and \$ p<0.05 Vs TiO




Fig. 3. Photomicrograph of testis of control, TiO₂ and NAC administered rats. In (a), testis characterized by well organized seminiferous tubules (st), which consisted of spermatogonium (g), primary and secondary spermatocytes (s), spermatid(td) and spermatozoa (z); H&E .X40. In (b), fine collagen fibres with blood vesseles represent the interstitial CT between the seminiferous tubules (arrow). Masson trichrome X40. In (c), the rat testis showed positive PAS reaction in the basement membrane (thick arrow), sertoli cells (*), and spermatozoa (→), while faint PAS reaction was detected in the spermatogenic cells (arrow) PAS X 40. In (d). The testis of the TiO₂ administered group showed appotosis of sertoli cells (arrow) H&EX 40. In (e), numerous interstitial oedema spreaded between the seminifrous tubules (o) and appotosis of germinal layers were also detected (arrow), H&E X40. In (f), some seminiferous tubules showed decrease in the thickness of the germinal layer (arrows) with rare sperms (rz) and interstitial oedema (o) H&E X10. In (g), The testis in NAC and TiO₂ administered rats showed partial improvement in the testicular cells while the congestion of the testicular blood vessels (C) and mild oedema (o) still persist PAS X40. In (h), the interstitial oedema was diminished in most of the seminiferous tubules while mild oedema still located in some parts of the interstitial CT (o) and the general architecture of the testis was restored, H&EX40.

As seen in **Fig. 2d**, 17 β -HSD expression is down regulated in TiO₂ administered rats and NAC normalized its expression. The overall effect of NAC alone as seen in **Fig. 2** is stimulatory for spermatogenesis related gene expression.

3.4. Histopathological Findings in Testis of Albino Rats after TiO₂ and NAC Administration

The testis of the control group was consisted of seminiferous tubules surrounded by interstitial Connective Tissue (CT). The seminiferous tubules lined by spermatogenic cells, primary and secondary spermatocytes, spermatid and spermatozoa. Pyramidal cells are located between the spermatogenic cells (Sertoli cells). Numerous interstitial cells (Leydig cells) were located in the interstitial CT (**Fig. 3a**). The interstitial CT mainly consisted of fine collagen fibers. Arterioles and venuoles were located between the interstitial CT in the testis of the NAC group (**Fig. 3b**). Positive PAS reaction was noticed in the basement membrane, sertoli cells and spermatozoa (**Fig. 3c**), while faint PAS reaction was detected in the spermatogenic cells. The testis of the TiO₂ administered group showed apoptosis of sertoli cells (**Fig. 3d**), numerous interstitial oedema spreaded between the seminiferous tubules and apoptosis of germinal layers were also detected (**Fig. 3e**), some seminiferous tubules showed decrease in the thickness of the germinal layer with rare sperms (**Fig. 3f**). The testis in NAC and TiO₂ administered rats showed partial improvement in the testicular cells while the congestion of the testicular blood vessels and mild oedema still persist (**Fig. 3g**), the interstitial oedema was diminished in most of the seminiferous tubules while mild oedema still located in some parts of the interstitial CT and the general architecture of the testis was restored (**Fig. 3h**).

4. DISCUSSION

In this study, we have shown the importance of NAC on testis function and its antiapoptotic and anti-inflammatory effect and its role in protection of testis. As known, NAC and L-carnitine are highly concentrated in the epididymis and spermatozoa, where it may serve as an intra-mitochondrial vehicle for the acyl group in the form of acyl CoA that acts as a substrate for the oxidation process to produce energy for sperm respiration and motility (Kanter *et al.*, 2010). Accumulation of TiO₂ in testis down regulated genes responsible for cholesterol transport and steroidogenesis including ABP, P450_{scc}, 17 β -HSD, CYP-17 α and aromatase. Expression of ABP, CYP17 α and 17 β -HSD

were significantly decreased in rats following exposure to TiO₂ for 3 months. These genes play important roles in testosterone biosynthesis, for example, Progesterone is then converted to 17 α -hydroxyprogesterone and androstenedione by CYP17 α . Finally androstenedione is converted to testosterone by 17 β -HSD. The decrease in gene expression of CYP17 α and 17 β -HSD indicates a possible contribution of these genes to the decrease in serum testosterone of male rats.

Apoptosis is a programmed cell death, it affects cell function and viability. Bcl2 is known as apoptotic biomarker. Administration of TiO₂ for 3 months up-regulated Bcl2 expression and NAC down regulate it confirming antiapoptotic activity of NAC as reported by ours and findings of (Aggarwal *et al.*, 2010). Germ cell apoptosis has been reported by several investigators to play an important role in normal testicular physiology (Modi *et al.*, 2003). It is required for normal spermatogenesis and is believed to ensure cellular homeostasis and maintain the fine balance between germ cells and Sertoli cells (Said *et al.*, 2004). Also, apoptosis is a key regulator of spermatogenesis in normal and pathological states (Kerr, 1992). High rates of apoptosis have been reported in testicular biopsies from infertile men with various degree of testicular insufficiency (Juriscova *et al.*, 1999). Sensitivity to factors induced apoptosis was higher in the spermatogonia and spermatocytes, while the Sertoli and Leydig cells were comparatively resistant (Kanter *et al.*, 2010). Such changes was ameliorated and normalized by NAC administration.

In this study, TiO₂ induced both oxidative stress that indicated by the decrease in reduced glutathione and inflammatory cytokines (IL-6 and TNF- α). It has been shown that NAC ameliorated the decrease in GST and lipid peroxidation induced by diazotization (Oksay *et al.*, 2013). Moreover, the changes in cytokines expression induced by TiO₂ are ameliorated by NAC administration. NAC acts as antioxidant that stimulate GSH synthesis (Dodd *et al.*, 2008) and is confirmed in this study by up-regulation of testicular GST expression in TiO₂ administered rats. So NAC can restore the imbalance between pro-oxidant and antioxidant systems during oxidative stress and that are coincided with our findings. In testis, NAC serve as an important intracellular cofactor for the transport of activated long-chain fatty acids into the mitochondrial matrix, thereby facilitating oxidation and enhancing energy production. In the epididymis, free antioxidants are taken up from the

circulating plasma and actively transported through the epithelial cells into the epididymal plasma, using a specific carrier under the regulation of androgens (Enomoto *et al.*, 2002). Moreover, antioxidants especially L-carnitine and NAC act as a factor capable of protecting the heart against ischemia/perfusion injury leading to myocyte death (Shug *et al.*, 1991).

The protective effect of NAC on testicular damage in rat varicocele, testis torsion models and tetracycline-induced toxicity were shown in studied of (Cay *et al.*, 2006; Payabvash *et al.*, 2007; Farombi *et al.*, 2008; Aktas *et al.*, 2010; Duarte *et al.*, 2010). They showed that NAC had good effects on testicular dysfunction but they did not examine testicular gene alterations that examined in our study. Toxicity induction by tetracyclins increased testicular Malondialdehyde (MDA) levels and decreased superoxide dismutase, catalase, GSH and glutathione-S-Transferase (GST) levels. NAC significantly attenuated these effects as reported in our findings (Farombi *et al.*, 2008). Moreover, increased doses of TiO₂ exhibited sever pathological changes, including rare sperm, sperm breakages, rarefaction of Sertoli cell and Sertoli cell apoptosis, necrosis of the seminiferous tubules, decreased germinative layer thickness, vaculation and irregular arrangement of sertoli cells of the seminiferous tubules (Gao *et al.*, 2013).

Numerous studies have unequivocally shown that TiO₂ exposure can migrate through different routes and accumulate in various organs (Chen *et al.*, 2009; Shi *et al.*, 2013; Jia *et al.*, 2013), which, in turn cause oxidative stress, inflammation and apoptosis resulting in organ injury and failure. Dysfunction of the testis in turn affects reproduction and affects the function of the reproductive system (Jia *et al.*, 2013). The cytochrome P-450 aromatase enzyme, the rate-limiting enzyme in estradiol synthesis, catalyzes the conversion of testosterone to estradiol. The present study demonstrated that aromatase mRNA levels decreased in TiO₂ administered rats and NAC normalized aromatase expression in a way to control estradiol biosynthesis (Shi *et al.*, 2007). It has been shown that NAC attenuates diazinon induced oxidative stress in rat testis (Oksay *et al.*, 2013). Michael *et al.* (2010) reported that NAC supplementation improved semen quality and canine fertility.

ABP is a glycoprotein secreted by Sertoli cells in the seminiferous tubules. It binds specifically to testosterone, dihydrotestosterone and 17-beta-estradiol.

The higher levels of these hormones enable spermatogenesis in the seminiferous tubules and sperm maturation in the epididymis (Patrao *et al.*, 2009). So, NAC probably acts to enforce spermatogenesis through increase in ABP expression. Moreover, aromatase, or estrogen synthase, is an enzyme responsible for a key step in the biosynthesis of estrogens. It is a member of the cytochrome P450 superfamily that catalyze many reactions involved in steroidogenesis. In particular, aromatase is responsible for the aromatization of androgens into estrogens (Lu *et al.*, 2010). In our results TiO₂ inhibited ABP, 17 β -HSD, CYP17 α and aromatase expression while NAC has significant upregulation of steroidogenesis gene expression. Moreover, NAC induced partial improvement in the testicular dysfunction, germinal layer apoptosis and diminished the interstitial oedema in most of the seminiferous tubules.

5. CONCLUSION

The findings of this study confirmed that co-administration of NAC inhibited harmful effects of TiO₂ that disrupted testicular function of male albino rats. NAC has antioxidant and antiapoptotic functions. Moreover, it improved steroidogenesis related gene expression which is ameliorated by TiO₂ administration. NAC is a good health supplement for protection of male fertility from environmental toxic biohazards.

6. REFERENCES

- Aggarwal, A., M.M. Misro, A. Maheshwari, N. Sehgal and D. Nandan, 2010. N acetylcysteine counteracts oxidative stress and prevents hCG-induced apoptosis in rat Leydig cells through down regulation of caspase-8 and JNK. *Mol. Reprod. Dev.*, 77: 900-909. PMID: 20824644
- Aktas, B.K., S. Bulut, S. Bulut, M.M. Baykam and C. Ozden *et al.*, 2010. The effects of N-acetylcysteine on testicular damage in experimental testicular ischemia/reperfusion injury. *Pediatr. Surg. Int.*, 26: 293-298. PMID: 19911182
- Arend, W.P. and C. Gabay, 2004. Cytokines in the rheumatic diseases. *Rheum. Dis. Clin. North Am.*, 30: 41-67. PMID: 15061568
- Atkuri, K.R., J.J. Mantovani, L.A. Herzenberg and L.A. Atkuri, 2007. N-Acetylcysteine-asafe antidote for cysteine/glutathione deficiency. *Curr. Opin. Pharmacol.*, 7: 355-359. PMID: 17602868

- Carson, F., 1990. Histotechnology: A Self-instructional Text. 3rd Edn., ASCP Press, Chicago, ISBN-10: 0891895817, pp: 400.
- Cay, A., A. Alver, M. Kucuk, O. Issik and M.S. Eminagaoglu *et al.*, 2006. The effects of N-acetylcysteine on antioxidant enzyme activities in experimental testicular torsion. *J. Surg. Res.*, 131: 199-203. PMID: 16412470
- Chen, J., X. Dong, J. Zhao and G. Tang, 2009. *In-vivo* acute toxicity of titanium dioxide nanoparticles to mice after intraperitoneal injection. *J. Applied Toxicol.*, 29: 330-337. DOI: 10.1002/jat.1414
- Choi, H., E. Stathatos and D.D. Dionysiou, 2006. Sol-gel preparation of mesoporous photocatalytic TiO₂ films and TiO₂/Al₂O₃ composite membranes for environmental applications. *Applied Catal. B Environ.*, 63: 60-67. DOI: 10.1016/j.apcatb.2005.09.012
- Dodd, S., O. Dean, D.L. Copolov, G.S. Malhi and M. Berk, 2008. N-acetylcysteine for antioxidant therapy: Pharmacology and clinical utility. *Expert Opin. Biol. Ther.*, 8: 1955-1962. PMID: 18990082
- Duarte, F., R. Blaya, P.E. Teloken, D. Becker and M. Fernandes *et al.*, 2010. The effects of N-acetylcysteine on spermatogenesis and degree of testicular germ cell apoptosis in an experimental model of varicocele in rats. *Int. Urol. Nephrol.*, 42: 603-608. PMID: 20012359
- Enomoto, A., M.F. Wempe and H. Tsuchida, 2002. Molecular identification of a novel carnitine transporter specific to human testis: Insights into the mechanism of carnitine recognition. *J. Biol. Chem.*, 277: 36262-36271. PMID: 12089149
- Farombi, E.O., M.C. Ugwuezunmba, T.T. Ezenwadu, M.O. Oyeyemi and M. Ekor, 2008. Tetracycline-induced reproductive toxicity in male rats: Effects of vitamin C and N-acetylcysteine. *Exp. Toxicol. Pathol.*, 60: 77-85. PMID: 18406588
- Feghali, C.A. and T.M. Wright, 1997. Cytokines in acute and chronic inflammation. *Front. Biosci.*, 2: 12-26. PMID: 9159205
- Fisher, J. and T. Egerton, 2001. Titanium Compounds, Inorganic. In: Kirk-Othmer Encyclopedia of Chemical Technology, Kroschwitz, J.I. and A. Seidel (Eds.), Wiley-Interscience, Hoboken, NJ.
- Gao, G., Y. Ze, X. Zhao, X. Sang and L. Zheng *et al.*, 2013. Titanium dioxide nanoparticles induced testicular damage, spermatogenesis suppression and gene expression alterations in male mice. *J. Hazard Mater.*, 258-259: 133-143. PMID: 23721730
- Hoet, P.H., I. Brske-Hohlfeld and O.V. Salata, 2004. Nanoparticles-known and unknown health risks. *J. Nanobiotechnol.*, 2: 12-12. DOI: 10.1186/1477-3155-2-12
- Jia, F., Z. Sun, X. Yan, B. Zhou and J. Wang, 2013. Effect of pubertal nano-TiO₂ exposure on testosterone synthesis and spermatogenesis in mice. *Arch. Toxicol.* PMID: 24241477
- Juriscova, A., S. Lopes and J. Merrano, 1999. DNA damage in round spermatids of mice with targeted disruption of the Pp1c 75 gene and in testicular biopsies of patients with non-obstructive azoospermia. *Mol. Hum. Reprod.*, 5: 323-330. PMID: 10321803
- Kaida, T., K. Kobayashi, M. Adachi and F. Suzuki, 2004. Optical characteristics of titanium oxide interference film and the film laminated with oxides and their applications for cosmetics. *J. Cosmet. Sci.*, 55: 219-220. PMID: 15190897
- Kanter, M., Y. Topcu-Tarlacalisir and S. Parlar, 2010. Antiapoptotic effect of L-carnitine on testicular irradiation in rats. *J. Mol. Hist.*, 41: 121-128. PMID: 20446105
- Kerr, J.B., 1992. Spontaneous degeneration of germ cells in rat testis: Assessment of cell types and frequency during the spermatogenic cycle. *J. Reprod. Fertil.*, 95: 825-830. PMID: 1404097
- Kiernan, J.A., 1999. Histological and Histochemical Methods: Theory and Practice. 3rd Edn., Butterworth Heinemann, Boston, ISBN-10: 0750631066, pp: 502.
- Li, N., Y.M. Duan, M.M. Hong, L. Zheng and M. Fei *et al.*, 2010. Spleen injury and apoptotic pathway in mice caused by titanium dioxide nanoparticles. *Toxicol. Lett.*, 195: 161-168. PMID: 20381595
- Lu, X., T. Yu, M. Ma, C. Jin, Q. Liu and S. Wu *et al.*, 2010. Effect on the testis development and StAR, CYP19a1 and CYP11a1 expression of prepubertal male rats after sub-acute exposure to DEHP. *Wei. Sheng.* 39: 263-267. PMID: 20568448
- Medina, C., M.J. Santos-Martinez and A. Radomski, 2007. Nanoparticles: Pharmacological and toxicological significance. *Br. J. Pharmacol.*, 150: 552-558. PMID: 17245366
- Michael, A.J., C. Alexopoulos, E.A. Pontiki, D.J. Hadjipavlou-Litina and P. Saratsis *et al.*, 2010. Effect of N-acetyl-L-cysteine supplementation in semen extenders on semen quality and reactive oxygen species of chilled canine spermatozoa. *Reprod. Domest. Anim.*, 45: 201-217. PMID: 19508647

- Modi, D.N., S. Sane and D. Bhartiya, 2003. Accelerated germ cell apoptosis in sex chromosome aneuploid fetal human gonads. *Mol. Hum. Reprod.*, 9: 219-225. PMID: 12651904
- Moschou, M., E.K. Kosmidi, M. Kaloyianni, A. Geronikaki and N. Dabarakis *et al.*, 2008. *In vitro* assessment of the neurotoxic and neuroprotective effects of N-Acetyl-L-Cysteine (NAC) on the rat sciatic nerve fibers. *Toxicol. Vitro*, 22: 267-274. PMID: 17959349
- Oberdorster, G., 2006. Toxicology of air born environment and occupational particles.
- Oksay, T., M. Naziroglu, O. Erguun, S. Dogan and O. Ozatik *et al.*, 2013. N-acetyl cysteine attenuates diazinon exposure-induced oxidative stress in rat testis. *Andrologia*, 45: 171-177. PMID: 22742659
- Patrao, M.T., E.J. Silva and M.C. Avellar, 2009. Androgens and the male reproductive tract: An overview of classical roles and current perspectives. *Arq. Bras. Endocrinol. Metabol.*, 53: 934-945. PMID: 20126845
- Payabvash, S., A.H. Salmasi, S. Kiumehr, S.M. Tavangar and B. Nourbakhsh *et al.*, 2007. Salutary effects of N-acetylcysteine on apoptotic damage in a rat model of testicular torsion. *Urol. Int.*, 79: 248-254. DOI: 10.1159/000107958
- Sadowska, A.M., B. Manuel-y-Keenoy and W.A. De Backer, 2007. Antioxidant and anti-inflammatory efficacy of NAC in the treatment of COPD: Discordant *in vitro* and *in vivo* dose-effects: A review. *Pulmonary Pharmacol. Therapeutics*, 20: 9-22. PMID: 16458553
- Said, T.M., U. Paasch and H.J. Glander, 2004. Role of caspases in male infertility. *Hum. Reprod. Update*, 10: 39-51. PMID: 15005463
- Sang, X.Z., L. Zheng, Q.Q. Sun, T. Zhang and N. Li *et al.*, 2012. The chronic spleen injury of mice following exposure to titanium dioxide nanoparticles. *J. Biomed. Mater. Res., Part A*, 100: 894-902. PMID: 22275130
- Sen, C.K. and L. Packer, 2000. Thiol homeostasis and supplements in physical exercise. *Am. J. Clin. Nutr.*, 72: 653S-69S. PMID: 10919972
- Shi, H., R. Magaye, V. Castranova and J. Zhao, 2013. Titanium dioxide nanoparticles: A review of current toxicological data. *Part Fibre. Toxicol.*, 10: 15-15. DOI: 10.1186/1743-8977-10-15
- Shi, Z., H. Zhang, Y. Liu, M. Xu and J. Dai, 2007. Alterations in gene expression and testosterone synthesis in the testis of male rats exposed to perfluorododecanoic acid. *Toxicol. Sci.*, 98: 206-215. PMID: 17400581
- Shug, A., A. Paulson, R. Subramanian and V. Regitz, 1991. Protective effects of propionyl-L-carnitine during ischaemia and reperfusion. *Cardiovasc. Drugs Ther.*, 5: 77-83. PMID: 2031875
- Stanislawski, L., M. Lefeuvre, K. Bourd, M.E. Soheili and M. Goldberg *et al.*, 2003. TEGDMA-induced toxicity in human fibroblasts is associated with early and drastic glutathione depletion with subsequent production of oxygen reactive species. *J. Biomed. Mat. Res.*, 66A: 476-482. PMID: 12918029
- Stanislawski, L., M.E. Soheili, A. Perianin and M. Goldberg, 2000. Dental restorative biomaterials induce glutathione depletion in cultured human gingival fibroblast: Protective effect of N-acetyl cysteine. *J. Biomed. Mat. Res.*, 51: 469-474. PMID: 10880090
- Turner, T.T. and J.J. Lysiak, 2008. Oxidative stress: a common factor in testicular dysfunction. *J. Androl.*, 29: 488-498. PMID: 18567643
- Van de Poll, M.C.G., C.H.C. Dejong and P.B. Soeters, 2006. Adequate range for sulfur-containing amino acids and biomarkers for their excess: Lessons from enteral and parenteral nutrition. *J. Nut.*, 136: 1694S-1700S. PMID: 16702341
- Wang, J., G. Zhou, C. Chen, H. Yu and T. Wang *et al.*, 2007. Acute toxicity and biodistribution of different sized titanium dioxide particles in mice after oral administration. *Toxicol. Let.*, 168: 176-185. PMID: 17197136
- Wilson, I. and M. Gamble, 2008. The Hematoxylin and Eosins. In: *Theory and Practice of Histological Techniques*, Bancroft, J.D. and M. Gamble (Eds.), Elsevier Health Sciences, London, UK., ISBN-10: 0443102791, pp: 125-138.
- Zafarullah, M., W.Q. Li, J. Sylvester and M. Ahmad, 2003. Molecular mechanisms of N-acetylcysteine actions. *Cellular Mol. Life Sci.*, 60: 6-20. PMID: 12613655