

## Measure of Center

### ❖ Measure of Center

the value at the center or middle of a data set

1. Mean
2. Median
3. Mode
4. Midrange (rarely used)

1

## Mean

### ❖ Arithmetic Mean (Mean)

the measure of center obtained by adding the values and dividing the total by the number of values

What most people call an *average*.

2

1

## Notation

$\Sigma$  denotes the **sum** of a set of values.

$x$  is the **variable** used to represent the individual data values.

$n$  represents the **number of data values in a sample**.

$N$  represents the **number of data values in a population**.

3

$\bar{x}$  is pronounced 'x-bar' and denotes the mean of a set of **sample values**

$$\bar{x} = \frac{\Sigma x}{n}$$

This is the **sample mean**

$\mu$  is pronounced 'mu' and denotes the mean of all values in a **population**

$$\mu = \frac{\Sigma x}{N}$$

This is the **population mean**

4

## Mean

### ❖ Advantages

Is relatively reliable.  
Takes every data value into account

### ❖ Disadvantage

Is sensitive to every data value, one extreme value can affect it dramatically; is not a *resistant* measure of center (cannot resist the extremes)

5

## Median

### ❖ Median

the **middle value** when the original data values are arranged in order of increasing (or decreasing) magnitude

❖ often denoted by  $\tilde{x}$  (pronounced 'x-tilde')

❖ is not affected by an extreme value - is a resistant (resists the extremes) measure of the center

6

2

## Finding the Median

First *sort* the values (arrange them in order), then follow one of these rules:

1. If the number of data values is odd, the median is the value located in the exact middle of the list.
2. If the number of data values is even, the median is found by computing the mean of the two middle numbers.

7

5.40	1.10	0.42	0.73	0.48	1.10
0.42	0.48	0.73	1.10	1.10	5.40

**MEDIAN is 0.915**

$$\frac{0.73 + 1.10}{2} \quad (\text{even number of values - no exact middle shared by two numbers})$$

5.40	1.10	0.42	0.48	0.73	1.10	0.66
0.42	0.48	0.66	0.73	1.10	1.10	5.40

exact middle **MEDIAN is 0.73**

8

## Mode

- ❖ **Mode**  
the value that occurs with the **greatest frequency**
- ❖ **Data set can have one, more than one, or no mode**
- Bimodal** two data values occur with the same greatest frequency
- Multimodal** more than two data values occur with the same greatest frequency
- No Mode** no data value is repeated

9

## Mode - Examples

- |                                  |                   |
|----------------------------------|-------------------|
| a. 5.40 1.10 0.42 0.73 0.48 1.10 | ↔ Mode is 1.10    |
| b. 27 27 27 55 55 55 88 88 99    | ↔ Bimodal 27 & 55 |
| c. 1 2 3 6 7 8 9 10              | ↔ No Mode         |

10

3

## Definition

- ❖ **Midrange**  
the value midway between the maximum and minimum values in the original data set

$$\text{Midrange} = \frac{\text{maximum value} + \text{minimum value}}{2}$$

11

## Midrange

- ❖ **Sensitive to extremes**  
because it uses only the maximum and minimum values.
- ❖ **Midrange is rarely used in practice**

12

## Round-off Rule for Measures of Center

Carry one more decimal place than is present in the original set of values

The mean of 3,4,5 is 3.3333.. Aprox to 3.3

The mean of 80.4 and 80.7 is 80.55

13

## Skewed and Symmetric

### ❖ Symmetric

distribution of data is symmetric if the left half of its histogram is roughly a mirror image of its right half


### ❖ Skewed

distribution of data is skewed if it is not symmetric and extends more to one side than the other


14

4


## Symmetry and skewness



Mode = Mean = Median  
(b) Symmetric



(a) Skewed to the Left  
(Negatively)



(c) Skewed to the Right  
(Positively)

15

## Measures of Variation

spread, variability of data  
width of a distribution

1. Standard deviation
2. Variance
3. Range (rarely used)

## Standard deviation

The **standard deviation** of a set of sample values, denoted by  **$s$** , is a measure of variation of values about the mean.

17

## Sample Standard Deviation Formula

$$s = \sqrt{\frac{\sum (x - \bar{x})^2}{n - 1}}$$

18

5

## Sample Standard Deviation (Shortcut Formula)

$$s = \sqrt{\frac{n\sum(x^2) - (\sum x)^2}{n(n - 1)}}$$

19

## Population Standard Deviation

$$\sigma = \sqrt{\frac{\sum (x - \mu)^2}{N}}$$

$\sigma$  is pronounced 'sigma'

This formula only has a theoretical significance, it cannot be used in practice.

20

## Variance

- ❖ The **variance** is a measure of variation equal to the square of the standard deviation.
- ❖ Sample variance:  $s^2$  - Square of the sample standard deviation  $s$
- ❖ Population variance:  $\sigma^2$  - Square of the population standard deviation  $\sigma$

21

## Variance - Notation

$s$  = *sample* standard deviation

$s^2$  = *sample* variance

$\sigma$  = *population* standard deviation

$\sigma^2$  = *population* variance

22

6

Usual values in a data set are those that are **typical** and **not too extreme**.

Minimum usual value = (mean) - 2 × (standard deviation)

Maximum usual value = (mean) + 2 × (standard deviation)

23

## Rule of Thumb

is based on the principle that for many data sets, the vast majority (such as 95%) of sample values lie within two standard deviations of the mean.

A value is unusual if it differs from the mean by more than two standard deviations.

24


## Empirical (or 68-95-99.7) Rule

For data sets having a distribution that is approximately bell shaped, the following properties apply:

- ❖ About 68% of all values fall within 1 standard deviation of the mean.
- ❖ About 95% of all values fall within 2 standard deviations of the mean.
- ❖ About 99.7% of all values fall within 3 standard deviations of the mean.

25


## The Empirical Rule



26


7

## The Empirical Rule



27

## The Empirical Rule



28

## Chebyshev's Theorem

- The above empirical rule applies only for bell-shaped distributions
- The Chebyshev's Theorem implies
  - 75% of all values lie within 2 standard deviation of the mean
  - 89% of all values lie within 3 standard deviation of the mean

29

## Range (rarely used)

The **range** of a set of data values is the difference between the maximum data value and the minimum data value.

$$\text{Range} = (\text{maximum value}) - (\text{minimum value})$$

It is very sensitive to extreme values; therefore not as useful as other measures of variation.

30

8

## Measures of Relative Standing

## Z score

❖ **Z score** (or standardized value)

the number of standard deviations that a given value  $x$  is above or below the mean

32

## Measure of Position: zscore

**Sample**

**Population**


$$z = \frac{x - \bar{x}}{s}$$

$$z = \frac{x - \mu}{\sigma}$$

Round zscores to 2 decimal places

33

## Interpreting Zscores



Whenever a value is less than the mean, its corresponding zscore is negative

Ordinary values:  $-2 \leq zscore \leq 2$

Unusual values:  $zscore < -2$  or  $zscore > 2$

34

9

## Percentiles

are measures of location. There are 99 **percentiles** denoted  $P_1, P_2, \dots, P_{99}$ , which divide a set of data into 100 groups with about 1% of the values in each group.

35

## Finding the Percentile of a Data Value

$$\text{Percentile of value } x = \frac{\text{number of values less than } x}{\text{total number of values}} \cdot 100$$

Round it off to the nearest whole number

36

## Converting from the $k$ th Percentile to the Corresponding Data Value


### Notation

$$L = \frac{k}{100} \cdot n$$

- $n$  total number of values in the data set
- $k$  percentile being used
- $L$  locator that gives the **position** of a value
- $P_k$   $k$ th percentile

37

## Converting from the $k$ th Percentile to the Corresponding Data Value



38

10

## Quartiles

Are measures of location, denoted  $Q_1$ ,  $Q_2$ , and  $Q_3$ , which divide a set of data into four groups with about 25% of the values in each group.


- ❖  $Q_1$  (First Quartile) separates the bottom 25% of sorted values from the top 75%.
- ❖  $Q_2$  (Second Quartile) same as the median; separates the bottom 50% of sorted values from the top 50%.
- ❖  $Q_3$  (Third Quartile) separates the bottom 75% of sorted values from the top 25%.

39

## Quartiles

$Q_1$ ,  $Q_2$ ,  $Q_3$

divide **ranked** scores into four equal parts



40

## Some Other Statistics

- ❖ Interquartile Range (or IQR):  $Q_3 - Q_1$
- ❖ Semi-interquartile Range:  $\frac{Q_3 - Q_1}{2}$
- ❖ Midquartile:  $\frac{Q_3 + Q_1}{2}$
- ❖ 10 - 90 Percentile Range:  $P_{90} - P_{10}$

41

## 5-Number Summary

- ❖ For a set of data, the **5-number summary** consists of the minimum value; the first quartile  $Q_1$ ; the median (or second quartile  $Q_2$ ); the third quartile,  $Q_3$ ; and the maximum value.

42