

Recall

Binomial Probability Distribution


1. The procedure must have a **fixed number of trials, n** .
2. The trials must be **independent**.
3. Each trial must have all outcomes classified into **two categories** (commonly, **success** and **failure**).
4. The probability of success **p** remains the same in all trials (the probability of failure is **$q=1-p$**).

1

Approximation of a Binomial Distribution with a Normal Distribution

If $np \geq 5$ and $nq \geq 5$

Then $\mu = np$ and $\sigma = \sqrt{npq}$
and the random variable has

a  distribution.
(normal)

Procedure for Using a Normal Distribution to Approximate a Binomial Distribution

1. Verify that **both** $np \geq 5$ and $nq \geq 5$. If not, you cannot use normal approximation to binomial.
2. Find the values of the parameters μ and σ by calculating $\mu = np$ and $\sigma = \sqrt{npq}$.
3. Identify the discrete whole number x that is relevant to the binomial probability problem. Use the continuity correction (See **continuity corrections** discussion later) Draw a normal curve and enter the values of μ , σ , and either $x - 0.5$ or $x + 0.5$, as appropriate.


2

Procedure for Using a Normal Distribution to Approximate a Binomial Distribution

4. Change x by replacing it with $x - 0.5$ or $x + 0.5$, as appropriate.
5. Using $x - 0.5$ or $x + 0.5$ (as appropriate) in place of x , find the area corresponding to the desired probability by first finding the z score and finding the area to the left of the adjusted value of x .

Example – Number of Men Among Passengers

Finding the Probability of “At Least 122 Men” Among 213 Passengers.



$$\mu = np = 213 * 0.5 = 106.5$$

$$\text{and } \sigma = \text{sqrt}(npq) = \text{sqrt}(213 * 0.5 * 0.5) = 7.29$$

Consider $x-0.5$ and $x+0.5$ for continuity correction

$$122 - 0.5 = 121.5 \text{ and } 122 + 0.5 = 122.5$$

$$Z = (121.5 - 106.5) / 7.29 = 2.06$$

Copyright © 2010, 2007, 2004 Pearson Education, Inc. All Rights Reserved.

5

3

Definition

When we use the normal distribution (which is a **continuous** probability distribution) as an approximation to the binomial distribution (which is **discrete**), a **continuity correction** is made to a discrete whole number x in the binomial distribution by representing the single value x by the interval from

$$x - 0.5 \text{ to } x + 0.5$$

(that is, adding and subtracting 0.5).

Copyright © 2010, 2007, 2004 Pearson Education, Inc. All Rights Reserved.

6


Procedure for Continuity Corrections

1. When using the normal distribution as an approximation to the binomial distribution, **always** use the continuity correction.
2. In using the continuity correction, first identify the discrete whole number x that is relevant to the binomial probability problem.
3. Draw a normal distribution centered about μ , then draw a **vertical strip area** centered over x . Mark the left side of the strip with the number $x - 0.5$, and mark the right side with $x + 0.5$. For $x = 122$, draw a strip from 121.5 to 122.5. **Consider the area of the strip to represent the probability of discrete whole number x .**


4

Procedure for Continuity Corrections


4. Now determine whether the value of x itself should be included in the probability you want. Next, determine whether you want the probability of at least x , at most x , more than x , fewer than x , or exactly x . Shade the area to the right or left of the strip, as appropriate; also shade the interior of the strip itself **if and only if x itself** is to be included. The total shaded region corresponds to the probability being sought.




$X =$ at least 122
(includes 122 and above)




$X =$ more than 122
(doesn't include 122)



$X =$ at most 122
(includes 122 and below)



$X =$ fewer than 122
(doesn't include 122)



$X =$ exactly 122

5

Ex. Continuity correction


EXAMPLE 2

Internet Penetration Survey A recent Pew Research Center survey showed that among 2822 randomly selected adults, 2060 (or 73%) stated that they are Internet users. If the proportion of all adults using the Internet is actually 0.75, find the probability that a random sample of 2822 adults will result in *exactly* 2060 Internet users.

Step 1: $np = 2822 \cdot 0.75 = 2116.5$ (Therefore $np \geq 5$)
 $nq = 2822 \cdot 0.25 = 705.5$ (Therefore $nq \geq 5$)

Step 2: $\mu = np = 2822 \cdot 0.75 = 2116.5$
 $\sigma = \sqrt{npq} = \sqrt{2822 \cdot 0.75 \cdot 0.25} = 23.002717$

Step 3: probability of *exactly* 2060
 strip from 2059.5 to 2060.5,




6

Recall: Sampling Distribution of the mean

- ❖ The sampling distribution of the mean is the distribution of the sample means, with all samples having the same sample size n taken from the same population

Recall: Example

- ❖ Consider rolling a die 5 times and find the mean \bar{x} of the results.




Copyright © 2010, 2007, 2004 Pearson Education, Inc. All Rights Reserved.

13

7

Recall: Sampling distribution of the variance

- ❖ The sampling distribution of the variance is the distribution of the sample variances, with all samples having the same size n taken from the population



Copyright © 2010, 2007, 2004 Pearson Education, Inc. All Rights Reserved.

14

Estimators Estimates and Sample Size

Some statistics work much better than others as estimators of the population. The example that follows shows this.

8

Review

- ❖ Chapters 2 & 3 used “descriptive statistics” when we summarized data using tools such as graphs, and statistics such as the mean and standard deviation.
- ❖ Chapter 6 introduced critical values:
 z_{α} denotes the z score with an area of α to its right.
If $\alpha = 0.025$, the critical value is $z_{0.025} = 1.96$.
That is, the critical value $z_{0.025} = 1.96$ has an area of 0.025 to its right.

Preview

This chapter presents the beginning of **inferential** statistics.

- ❖ The two major activities of inferential statistics are (1) to use sample data to estimate values of a population parameters, and (2) to test hypotheses or claims made about population parameters.
- ❖ We introduce methods for estimating values of these important population parameters: proportions, means, and variances.
- ❖ We also present methods for determining sample sizes necessary to estimate those parameters.

9

Section 7-2 Estimating a Population Proportion



Key Concept

In this section we present methods for using a sample proportion to estimate the value of a population proportion.

- The sample proportion is the best point estimate of the population proportion.
- We can use a sample proportion to construct a confidence interval to estimate the true value of a population proportion, and we should know how to interpret such confidence intervals.
- We should know how to find the sample size necessary to estimate a population proportion.

10

Definition

A **point estimate** is a single value (or point) used to approximate a population parameter.

Definition

The sample proportion \hat{p} is the best point estimate of the population proportion p .

11

Example:

In a Research Center poll, 70% of 1501 randomly selected adults in the U. S. believe in global warming, so the sample proportion is $\hat{p} = 0.70$. Find the best point estimate of the proportion of all adults in the United States who believe in global warming.

Because the sample proportion is the best point estimate of the population proportion, we conclude that the best point estimate of p is 0.70. When using the sample results to estimate the percentage of all adults in the U.S. who believe in global warming, the best estimate is 70%.

But how reliable (accurate) is this estimate?

We will see that its **margin of error- coming after- is 2.3%. This means the true proportion of adults who believe in global warming is between 67.7% and 72.3%. This gives an interval (from 67.7% to 72.3%) containing the true (but unknown) value of the population proportion.**