

Phytochemical Screening, DNA Fingerprinting, and Nutritional Value of *Plectranthus amboinicus* (Lour.) Spreng

Seham S. El-hawary, Rabie H. El-sofany, Azza R. Abdel-Monem*, Rehab S. Ashour

Department of Pharmacognosy, Faculty of pharmacy, Cairo University, Cairo 11562, Egypt.

ABSTRACT

In this study, *Plectranthus amboinicus* (Lour.) Spreng was subjected to phytochemical and genetic investigation, so that it could be properly identified. The results of phytochemical screening of the different plant organs (leaves, stems and roots) revealed the presence of steam volatile substances, sterols and/or triterpenes, flavonoids, carbohydrates and/or glycosides, catechol tannins in the three organs under investigation. A DNA sample of the plant was analyzed by Random Amplified Polymorphic DNA (RAPD) technique using eleven oligonucleotide primers. The analysis of RAPD data achieves the use of B-19 and B-6 primers for selective discrimination of this plant. Also, the nutritional value, including the total carbohydrates, total soluble sugars, proteins, amino acids and vitamin content, of the leaves, stems and roots was determined. The roots recorded the highest values of total carbohydrates, total soluble sugars and proteins content (66.04, 23.33 and 17.58 g%, respectively), followed by the leaves (48.12, 4.74 and 16.45 g%, respectively) and the stems (44.62, 3.10 and 9.52 g%, respectively). The three organs under investigation contain essential amino acids in moderate amounts compared to the WHO daily recommended doses. They are rich in vitamins and can be used as a vitamins supplement.

Key words: *Plectranthus amboinicus*, DNA fingerprinting, phytochemical screening, nutritional value

INTRODUCTION

Genus *Plectranthus* belongs to family Lamiaceae and comprises about 350 species cultivated as ornamental plants or as sources of essential oils.^[1] *Plectranthus amboinicus* (Lour.) Spreng (synonyms include *Plectranthus aromaticus* Roxb., *Coleus aromaticus* Benth. and *Coleus amboinicus* Lour.), is a perennial herb, native to Indonesia and is distributed in Tropical Africa, Asia and Australia. It is used as food, additive and fodder, and as medicine in treating a wide range of diseases.^[1] The leaves extract is used to treat inflammatory disease,^[2,3] chronic cough and urinary disease^[2]. It is also used as an aromatic carminative and anthelmintic.^[2] It was reported to have antimicrobial,^[2,4] cytotoxic^[2,3] and antioxidant activities^[2,5]. Essential oils, diterpenes, flavonoids

and phenolic acids are the main constituents isolated from the different *Plectranthus* species.^[6,7] Because of taxonomic similarities of the different *Plectranthus* species, the same species of *Plectranthus* usually shows a number of synonyms.^[1] To help in solving this terminology problem, in this study, the different organs (leaves, stems and roots) of *Plectranthus amboinicus* (Lour.) Spreng growing in Egypt were subjected to a phytochemical screening aiming for chemical identification of this plant and the DNA profile of the plant was analyzed for genetic identification. The total carbohydrates, total soluble sugars, proteins, amino acids and vitamin content were determined to evaluate the nutritional value of this plant as a food.

MATERIALS AND METHODS

Plant material

Plectranthus amboinicus (Lour.) Spreng were collected over the years (2008-2010) from El-Orman garden. The plant was identified by Dr. Mohamed el Gebaly and Madam Treze (Taxonomist). A voucher specimen was kept in the herbarium of Department of Pharmacognosy, Faculty of Pharmacy, Cairo University.

*Address for correspondence:

Azza R. Abdel-Monem
Faculty of Pharmacy, Cairo University, Cairo, Egypt
E-mail: azzaramy@yahoo.com

DOI: 10.5530/pj.2012.30.2

Preliminary Phytochemical Screening

The powdered air-dried leaves, stems and roots of *Plectranthus amboinicus* (Lour.) Spreng were tested for the presence of steam volatile substances,^[8] sterols and/or triterpenes,^[9,10] flavonoids,^[11,12] crystalline sublimate,^[13] carbohydrates and/or glycosides,^[14,15] tannins,^[16] saponins,^[17] alkaloids and/or nitrogenous bases,^[18] anthraquinones,^[19] cardiac glycosides,^[8,20-23] and oxidase enzyme^[8]. The results are given in table 1.

DNA fingerprinting

DNA fingerprinting was performed in Agricultural Genetic Engineering Research Institute using the Random Amplified Polymorphic DNA (RAPD) technique.^[24] DNA was extracted using CTAB extraction buffer (1% N-cetyl-N,N,N trimethyl ammonium bromide). Eleven oligonucleotide primers (Operon Technologies Inc., Alameda, California, USA) were used for analysis. Amplification of DNA was carried out in thermal cycler, programmed as follows: first cycle at 94 °C for 5 min. (initial strand separation), followed by one cycle for 40 seconds at 94 °C (denaturation), 1 min. at 36 °C (annealing), forty cycles each for 1 min at 72 °C (elongation) and the last cycle for 7 min at 72 °C (final extension).

PCR reactions were performed in polypropylene tubes containing 2.5 µl reaction buffer, 2 µl MgCl₂, 2.5 µl of each dNTPs (Pharmacia, Sweden), 3 µl primers, 0.5 µl Taq DNA polymerase (Perkin-Elmer/Cetus, USA; advanced Biotechnologies, UK), 3 µl template DNA, and enough sterilized water to obtain 25 µl.

PCR material was separated by horizontal electrophoresis in a 1.5% agarose gel plate (Sigma Co.). 10 µl of each PCR product was mixed with 3 µl loading buffer

and loaded onto wells of the gels. The gels were run at 95 volts.

After electrophoresis the RAPD pattern was visualized by staining the gel with ethidium bromide solution (0.5 µg/ml), visualized under UV light, and photographed using a gel documentation system. RAPD molecular weight markers (Biolab Co.) were used. The banding profile produced by the eleven decamer primers is given in table 2.

Determination of total carbohydrates and total soluble sugars

The total carbohydrate and total soluble sugars were determined in the leaves, stems and roots by colorimetric method according to Dubois et al, (1956).^[25]

Determination of protein content

The total protein content was determined, in the three organs under investigation adopting Micro-Kjeldahl method according to A.O.A.C. (1995).^[26]

Determination of amino acids

The amino acids content in the three organs was determined by spectrophotometric method using amino acids analyzer (AAA 400, INGOS Ltd) after acid hydrolysis.^[27] The results obtained are shown in table 3.

HPLC analysis of vitamins

HPLC analysis of vitamins^[28-31] was carried out on Agilent 1100 apparatus equipped with Hypersil-BDS-C₁₈ column (4.6 × 250 mm). The injection volume was 5 µl, the mobile phase was methanol at a flow rate of 1 ml/min. Detection was carried out with UV detector. The results are shown in table 4.

RESULTS AND DISCUSSION

Preliminary Phytochemical Screening

The results of phytochemical screening showed the presence of steam volatile substances, sterols and/or triterpenes, flavonoids, carbohydrates and/or glycosides, catechol tannins in all organs of *Plectranthus amboinicus* (Lour.) Spreng. Crystalline sublimate, saponins, alkaloids and/or nitrogenous bases, cardiac glycosides and oxidase enzyme are absent in all organs.

Genetic profiling

The RAPD electrophoretic profile of the DNA sample amplified with the eleven decamer primers generated 58 fragment patterns, produced by B-19 (9 bands), B-6 (9 bands), D-20 (8 bands), E-4 (6 bands), A-6 and G-5 (5 bands each), A-17 and G-2 (4 bands each), A-18 and G-17 (3 bands each) and G-19 (2 bands). Thus, primers B-19 and B-6 could be selected for discrimination of *Plectranthus amboinicus* (Lour.) Spreng.

Table 1: Results of phytochemical screening of the different organs of *Plectranthus amboinicus* (Lour.) Spreng

Constituent	Leaves	Stems	Roots
Steam volatile substances	++	+	±
Sterols and/ or triterpenes	++	+	+
Free aglycones	++	+	+
Flavonoids	+	+	+
Crystalline sublimate	-	-	-
Carbohydrates and/or glycosides	+	+	+
Catechol tannins	+	+	++
Pyrogallol tannins	-	-	-
Saponins	-	-	-
Alkaloids and/or nitrogenous bases	-	-	-
Free anthraquinones	-	-	-
Combined anthraquinones	-	-	-
Cardiac glycosides	-	-	-
Oxidase enzyme	-	-	-

+: Present, ++: Strongly positive, -: Absent, ±: Traces

Table 2: Molecular size in base pairs of amplified DNA fragments produced by eleven decamer primers in *Plectranthus amboinicus* (Lour.) Spreng

Approx. band size (bp)	Decamer Primers										
	A6	A17	A18	B6	B19	E4	G2	G5	G17	G19	D20
1375					+						
1343				+							
1311	+	+									
1295								+			
1280					+	+					
1265				+							
1205				+							
1191					+						
1122	+										
1108							+				
1095											+
1069					+						
1032	+			+							
1019						+					
995								+			+
983						+	+				
960					+				+		
949							+				
842						+					
822											+
812				+							
793					+						
774		+				+					
756								+			
747							+				
720		+		+							+
695						+					
670		+							+		
631								+		+	
609											+
602			+								
595	+				+						
574			+								
553					+				+	+	+
534				+							
521								+			
503											+
479				+							
468											+
451			+								
441					+						
391				+							
343	+										
Sum	5	4	3	9	9	6	4	5	3	2	8

+: Present

Nutritional value

Total carbohydrates were found to be 48.12, 44.62 and 66.04 g% of dry weight in the leaves, stems and roots, respectively. The total soluble sugars recorded 4.74, 3.10 and 23.33 g% of dry weight in the leaves, stems and roots, respectively. The protein content of the leaves, stems and roots was 16.45, 9.52 and 17.58 g% of dry weight, respectively. Aspartic acid (25.02, 25.80 and 19.35 mg/100 g of dry weight), glutamic acid (12.71, 10.79,

and 13.23 mg/100 g of dry weight) and glycine (12.45, 12.77 and 14.02 mg/100 g of dry weight) were the main identified amino acids in the leaves, stems and roots, respectively. Meanwhile, the major detected amino acids were non-essential amino acids, the three organs under investigation contained essential amino acids (threonine, valine, isoleucine, leucine, phenyl alanine, histidine and lysine) in moderate amounts compared to the WHO daily recommended doses.^[32] The three organs are rich in vitamins

Table 3: Amino acids in *Plectranthus amboinicus* Lour. (Spreng) leaves, stems and roots

Amino acid	Percentage of amino acid (mg/100 g of dry weight)		
	Leaves	Stems	Roots
Aspartic acid	25.02	25.80	19.35
Threonine	4.16	5.33	4.49
Serine	5.66	6.76	6.77
Glutamic acid	12.71	10.79	13.23
Proline	0.09	0.10	0.04
Glycine	12.45	12.77	14.02
Alanine	8.48	10.73	9.92
Valine	5.82	0.55	7.30
Isoleucine	2.61	2.93	4.17
Leucine	7.15	8.66	7.86
Tyrosine	1.04	0.69	0.84
Phenyl alanine	3.81	3.83	3.58
Histidine	2.09	2.35	4.27
Lysine	4.45	5.81	3.43
Arginine	4.43	2.85	0.71

Table 4: Vitamins content in the leaves, stems and roots of *Plectranthus amboinicus* Lour. (Spreng)

Vitamins	Leaves	Stems	Roots
Vitamin A (ppm)	0.38	0.12	0.02
Vitamin E (ppm)	2.93	1.26	0.33
Vitamin D (ppm)	4.471	0.0313	0.209
Vitamin C (%)	0.11	0.11	0.12
Vitamin B complex (%)			
Thiamin B1	0.03	0.01	0.02
Pyridoxin B6	0.01	--	0.008
Riboflavin B2	2.47	0.91	0.7
Nicotinic B3 (Niacin)	0.52	0.08	0.1
Cyanocobalamine B12	0.37	0.14	--
Folic acid	0.002	--	0.03

according to the WHO daily recommended doses,^[33] and can be used as a vitamins supplement. From the results obtained, it could be concluded that *Plectranthus amboinicus* (Lour.) Spreng is a useful edible plant, can be used as a herbal nutrition supplement and can be considered as a multivitamins supplementary agent, for vitamins deficiency patients, or individuals on a restricted diet. This may explain its traditional use as a food or food additive.

REFERENCES

- Lukhoba CW, Simmonds MSJ, Paton AJ. *Plectranthus*: A review of ethnobotanical uses. *Journal of Ethnopharmacology*. 2006; 103:1-24.
- Khare RS, Banerjee S, Kundu K. *Coleus aromaticus* Benth-A nutritive medicinal plant of potential therapeutic value. *International Journal of Pharma and Bio Sciences*. 2011; 2(3):488-500.
- Gurgel AP, da Silva JG, Grangeiro AR, Oliveira DC, Lima CM, da Silva AC, Oliveira RA, Souza IA. *In vivo* study of the anti-inflammatory and antitumor activities of leaves from *Plectranthus amboinicus* (Lour.) Spreng Lamiaceae. *Journal of Ethnopharmacology*. 2009; 125:361-3.
- Vijaya KS, Syed MA, Badami S, Anil TM, David B. Antibacterial activity of aqueous extract of *Coleus amboinicus*. *Pharmacognosy Online*. 2008; 3:224-6.
- Patel RD, Mahobia NK, Singh MP, Singh A, Sheikh NW, Alam G, *et al.* Antioxidant potential of leaves of *Plectranthus amboinicus* (Lour.) Spreng. *Der Pharmacia Lettre*. 2010; 2:240-5.
- Adel-Mogib M, Albar HA, Batterjee SM. Chemistry of genus *Plectranthus*. *Molecules*. 2002; 7:271-301.
- Kumaran A, Karunakaran RJ. Activity-guided isolation and identification of free radical-scavenging components from an aqueous extract of *Coleus aromaticus*. *Food-Chemistry*. 2007; 100:356-361.
- Wagner H, Baldt S, Zgainski EM. *Droger Analyse*. Springer-Verlag, Berlin; 1983.
- Liebermann C, Burchard H. *Chem. Zentr*. 1890; 61:7.
- Wall M, Krieder M, Krewson C, Eddy C, William J, Corel D, *et al.* *Amer. Pharm. Assoc*. 1954; 43:1-7.
- Geissman TA. *The Chemistry of Flavonoid Compounds*. The Macmillan Company, New York; 1962.
- Willstätter R. *Ber. Deutsch. Pharm. Ges*. 1964; 47:2831. [Through J. Ph. Sci. 1966; 55, 276].
- Claus EP, Tyler V. *Pharmacognosy*. 5th ed., Lea and Febiger, Philadelphia; 1967.
- Molisch H. *Montash Chem*. 1886; 7:198. [Through Stank J, Cerny M, Kocourek J, Pacok J. *The Monosaccharides*. Publishing House of Czechoslovak, Academy of Science, Prague; 1963].
- Coutts RT, Snail GA. *Polysaccharides, Peptides and Proteins*. William Heinemann Medical Book Ltd., London; 1973.
- Evans WC. *Trease and Evans' Pharmacognosy*. 14th ed., WB Saunders Company Ltd., London, Philadelphia, Toronto, Sydney, Tokyo; 1996.
- Walform ML, Morgan PW, Benton FL. *J. Am. Chem. Soc*. 1940; 62:1484.
- Josephna P, Romo J. Isolation and structure of peruvirin. *Tetrahedron*. 1966; 22(6):1723.
- Borntrager HZ. *Anal. Chem*. 1880; 19:156. [Through Fairbarin JW. *J. Pharm. and Pharmacol*. 1949; 1:683].
- Fieser LF, Fieser M. *Steroids*. Reinhold Publishing Co., New York; 1959.
- Keller CC. *Ber. Deutsch. Pharm. Ges*. 1895. [Through Fieser LF, Fieser M. *Steroids*. Reinhold Publishing Co., New York; 1959].
- Killiani H. *Arch. Pharm*. 1896. [Through Fieser LF, Fieser M. *Steroids*. Reinhold Publishing Co., New York; 1959].
- Baljet H. *Scheveiz. Apoth. Ztg*. 1918. [Through chemical abstract. 12, 1336, 1918].
- Doyle JJ, Doyle JL. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull*. 1987; 19:11-5.
- Dubois M, Smith F, Gilles KA, Hamilton JK, Rebers PA. Colorimetric method for determination of sugar and related substances. *Anal. Chem*. 1956; 28(3):350-6.
- A.O.A.C. *Method of analysis*. Association of Official Agriculture Chemists. 16th ed., Arlington, VA, USA; 1995.
- Block RJ, Durrum EL, Zweig G. *Annual of paper chromatography and paper electrophoresis*. 2nd ed. Academic Press, New York; 1958.
- Pyka A, Sliwiok J. Chromatographic separation of tocopherols. *J Chromatography A*. 2001; 935:71-6.
- Romeu NM, Morera PS, Castellote AI, Lopez SMC. Rapid high-performance liquid chromatographic method for vitamin C determination in human milk versus an enzymatic method. *J chromatography B*. 2006; 830:41-6.
- Blanco D, Fernandez MP, Gutierrez MD. Simultaneous determination of fat-soluble and provitamins in dairy products by a liquid chromatography with narrow-bore column. *Analyst*. 2000; 125:427-31.
- Batifoulie F, Verry MA, Besson C, Demigné C, Rémésy C. Determination of thiamine and its phosphate esters in rat tissues analyzed as thiochromes on a RP-amide C16 column. *J Chromatography B*. 2005; 816:67-72.
- Protein and amino acid requirements in human nutrition. WHO Press, World Health Organization, Avenue Appia, Geneva, Switzerland; 2007.
- Vitamin and Mineral Requirements in Human Nutrition. 2nd ed., WHO Press, World Health Organization, Avenue Appia, Geneva, Switzerland; 2004.