



ISSN NO. 2320-5407

Journal homepage: <http://www.journalijar.com>INTERNATIONAL JOURNAL  
OF ADVANCED RESEARCH

## RESEARCH ARTICLE

**One pot synthesis of pyridine, thiazolidine, pyrazole and 2, 3-dihydro-1, 3, 4-thiadiazole derivatives under solvent-free condition**Anhar Abdel-Aziem<sup>1</sup> and Abdou Osman Abdelhamid\*<sup>2</sup><sup>1</sup>Department of Chemistry, Faculty of Science (Girls), Azhar University, Nasr City, Cairo, 11754, Egypt.<sup>2</sup>Department of Chemistry, Faculty of Science, Cairo University, Giza, 12613, Egypt.**Manuscript Info****Manuscript History:**

Received: 10 October 2013

Final Accepted: 25 October 2013

Published Online: November 2013

**Key words:**

2-Acetylbenzofuran, Hydrazonoyl halides, Pyridins, 1,3-Thiazoles, 1,3,4-Thiadiazoles, Pyrazoles; Solvent-free.

**Abstract**

Claisen condensation of 2-acetylbenzofuran and 1,3-diphenyl-1*H*-pyrazole-4-carbaldehyde **2** gave chalcone **3** which was exploited as a starting material for the syntheses of hitherto unknown different types of new heterocyclic compounds incorporating the benzofuran moiety via the reaction with various active methylene compounds. On the other hand, several new thiadiazolines, pyrazoles and pyrazolo[3,4-*d*]pyridazine were synthesized by the reaction of 3-(benzofuran-2-yl)-3-oxopropanenitrile **12** with  $\alpha$ -halo ketone,  $\alpha$ -halo ester and hydrazonoyl halides.

Copy Right, IJAR, 2013.. All rights reserved.


**Introduction**

Many organic solvents are hazardous and can be deleterious to human health. They are volatile and cause an environmental threat by polluting the atmosphere [1,2]. The replacement of volatile organic solvents in organic reaction processes is an important green chemistry goal. Furthermore, the solvent-free or solid-state reactions have numerous advantages: reduced pollution, low costs, high yields and purities of products, simplicity in process and handling [3,4]. On the other hand, various benzofuran derivatives possess different pharmacological and biological activities, of which the most potent is anti-inflammatory [5], pesticidal & insecticidal [6], antihistaminic [7], anticonvulsant [8], antiallergic [9] and In-vitro anti-HIV-1, anticancer, anti-microbial activities [10, 11]. The antimicrobial activity of benzofuran derivative appears to be dependent on substitution at the heterocyclic furan ring than the aromatic moiety [12]. As a continuation of our program, we report a facial, eco-friendly approach with excellent yield for the synthesis of pyridine, pyrazole, thiazoline and 2,3-dihydro-1,3,4-thiadiazoles bearing benzofuran moiety by grinding method.

**Result and Discussion**

The starting compound **3** was synthesized *via* grinding where 2-acetylbenzofuran (**1**), 1,3-diphenyl-1*H*-pyrazole-4-carbaldehyde (**2**) and moist potassium hydroxide were ground with a pestle in an open mortar (Scheme 1). Structure of the product **3** was confirmed on the basis of its correct elemental analysis, spectral data and chemical transportation. The IR spectrum of compound **3** revealed bands at 3064  $\nu$  (CH-arom), 1660  $\nu$  (CO), 1610  $\nu$  (C=N), 1570  $\nu$  (C=C). Its <sup>1</sup>H NMR spectrum revealed signals at  $\delta$  = 7.10-8.17 (m, ArH's and -CH=CH-). Thus, treatment of 1-(benzofuran-2-yl)-3-(1,3-diphenyl-1*H*-pyrazol-4-yl)prop-2-en-1-one (**3**) with 2-cyanoethanethioamide and ammonium acetate in acetic acid under reflux afforded the corresponding 2-mercapto-6-(benzofuran-2-yl)-4-(1,3-diphenyl-1*H*-pyrazol-5-yl)pyridine-2(1*H*)-thione (**4**). Similarly, compound **3** reacted with each of malononitrile, 2-cyanoacetamide or 2-cyanoacetohydrazide yielded the corresponding pyridine derivatives **5-7** (Scheme 1). The structures of **4-7** were confirmed on the basis of their elemental analysis, spectral data and alternative synthetic route. The IR spectrum of **4** revealed bands at 3340  $\nu$  (NH), 3058  $\nu$  (CH-aroma.), 2218  $\nu$  (CN), 1604  $\nu$  (C=N), 1234  $\nu$  (C=S) and the IR spectrum of **5** showed the presence of 3266, 3200  $\nu$  (NH<sub>2</sub>), 3062  $\nu$  (CH-aroma.), 2206  $\nu$  (CN), 1589  $\nu$  (C=C). The structure of **5** was supported by its <sup>1</sup>H NMR which showed signals at  $\delta$  = 4.24 (s, 2H, NH<sub>2</sub>), 7.20-7.85 (m, 17H, ArH's). Thus, one pot reaction of 2-acetylbenzofuran (**1**), 1,3-diphenyl-1*H*-pyrazole-4-carbaldehyde

(2) and the appropriate of 2-cyanoethanethioamide, malononitrile, 2-cyanoacetamide or 2-cyanoacetohydrazide and excess ammonium acetate in n-butanol gave products identical in all aspects (mp., mixed mp. and spectra) with the corresponding compounds **4-7** (Scheme 1). Structure **4** was also confirmed by chemical transformation, where the appropriate of iodomethane, ethyl chloroacetate, chloroacetone,  $\omega$ -bromoacetophenone and chloroacetonitrile were reacted with **4** by two different methods either with ethanolic triethylamine or (ethanolic potassium hydroxide) at room temperature *or via* grinding in which equimolar amounts of pyridine-2-thion **4**, potassium hydroxide and each of methyl iodide, ethyl chloroacetate, chloroacetone,  $\omega$ -bromoacetophenone and chloroacetonitrile were mixed and ground in a mortar to give the corresponding product **8a-c** and **9a** and **9b**, respectively (Scheme 1, Table 1).




Scheme 1

Structures of **8a-c**, **9a** and **9b** were elucidated on basis of analytical and spectral data. Thus, IR spectrum of **8b** showed the absence of SH band and the presence of bands at 3058  $\nu$ (CH-aromatic), 2981, 2931  $\nu$ (CH-aliphatic),

2214  $\nu(\text{CN})$ , 1732  $\nu(\text{C}=\text{O})$ , 1596  $\nu(\text{C}=\text{C})$ . Its  $^1\text{H NMR}$  ( $\text{DMSO-}d_6$ ) revealed signals at  $\delta$  1.20 (t, 3H,  $J = 7$  Hz,  $\text{CH}_2\text{CH}_3$ ), 4.22 (q, 2H,  $J = 7$  Hz,  $\text{CH}_2\text{CH}_3$ ), 4.70 (s, 2H,  $\text{SCH}_2$ ), 7.22-810 (m, 17H, ArH's).


However, compound **3** was reacted with thiosemicarbazide afforded 5-(benzofuran-2-yl)-2',5'-diphenyl-3,4-dihydro-2*H*,2'*H*-[3,3'-bipyrazole]-2-carbothioamide (**10**). Structure **10** was elucidated *via* elemental analysis, spectral data and chemical transformation. Thus,  $^1\text{H NMR}$  spectrum of **10** showed signals at  $\delta$  =



$\delta = 3.66$  (dd, 1H,  $J = 18.1, 5.8\text{Hz}$ ,  $\text{CH}_2(\text{pyraz})$ ), 4.01 (dd, 1H,  $J = 18.1, 12\text{Hz}$ ,  $\text{CH}_2(\text{pyraz})$ ), 5.33 (dd,  $J = 12.2, 5.8$  Hz,  $\text{CH}(\text{pyraz})$ ), 5.98 (s, 1H, pyrazole H-5), 7.00–7.82 (m, 15H, ArH's and pyrazole H-5), 9.34 (s, br., 2H,  $\text{NH}_2$ ). Compound **10** was reacted with the appropriate hydrazonoyl halides **11b,c** gave 2-(5-(benzofuran-2-yl)-3',5'-diphenyl-3,4-dihydro-1*H*,2*H*-[3,4'-bipyrazol]-2-yl)-4-methyl-5-(phenyl-diazenyl)thiazole (**12a**), and 2-(5-(benzofuran-2-yl)-3',5'-diphenyl-3,4-dihydro-1*H*,2*H*-[3,4'-bipyrazol]-2-yl)-4-phenyl-5-(phenyldiazenyl)thiazole (**12b**) (Scheme 2). Structures **12** were confirmed by elemental analyses, spectral data, and alternative synthetic routes. Thus, benzenediazonium chloride reacted with 2-(5-(benzofuran-2-yl)-2',5'-diphenyl-3,4-dihydro-2*H*,2'*H*-[3,3'-bipyrazol]-2-yl)-4-phenylthiazole (**13**), which prepared via reaction of **3** with **14** or **10** with *o*-bromoacetophenone, in pyridine to give a product identical in all aspects (m.p., mixed m.p., and spectra) with **12b** (Scheme 2, Table 1).

On the other hand, the reactivity of compound **15** [13, 14] towards a variety of hydrazonoyl halides was also studied. Thus, treatment of compound **15** with the appropriate hydrazonoyl halides **11a,b** and **11d** in moist sodium carbonate furnished without solvents, in each case, one isolable product. The isolable products were assigned as 3-substituted


5-(benzofuran-2-yl)-4-cyano-1-phenyl-1*H*-pyrazole **17a-c** on the basis of their analytical and spectral data (Scheme 3, Table 1). The IR spectrum of **17a-c** revealed bands near 2229  $\nu(\text{CN})$ , 1724-1693  $\nu(\text{C=O})$ . The  $^1\text{H}$  NMR spectrum of **17a** in (DMSO- $d_6$ ) revealed signals at  $\delta$  1.33 (t, 3H,  $J = 7$  Hz,  $\text{CH}_2\text{CH}_3$ ), 4.41 (q, 2H,  $J = 7$  Hz,  $\text{CH}_2\text{CH}_3$ ), 6.93 (s, 1H, CH-furan) and 7.41-7.63 (m, 9H, ArH's). From the above data the structures **18** were ruled out (Scheme 3).



**Scheme 3**

Structure of **17a-c** can also be confirmed by alternative synthetic routes. Thus, compound **15** together with the appropriate **11a-c** in ethanol in presence of sodium ethoxide gave product identical in all aspects (mp. mixed mp. and spectra) with **17**. The corresponding pyrazolopyridazine derivatives were obtained by the reaction of pyrazoles **17a** with hydrazine hydrate under reflux in ethanol for 5 hrs. The lack of nitrile and carbonyl groups in the IR spectrum of the isolated product supported the formation of pyridazine **19** (Scheme 3).

Finally, 2,3-dihydro 1,3,4-thiadiazoles **22a-f** were obtained by grinding (solvent-free conditions) of compound **15**, potassium hydroxide, phenyl isothiocyanate and hydrazonoyl halides (Scheme 4). Thus,  $^1\text{H}$  NMR spectrum of **22f** showed signal at  $\delta$  2.520 (s, 3H,  $\text{CH}_3$ ), 6.90 (s, 1H, CH-furan) and 7.08-8.30 (m, 13H, ArH's).



Scheme 4

Table1: Comparison between grinding and traditional method for synthesized compounds

Comp. No.	Grinding method		Traditional method	
	Time	Yield %	Time	Yield %
3	10 min.	95%	2 hrs	88%
4	10 min.	92%	4 hrs	85%
5	10 min.	92%	4 hrs	75%
6	10 min.	92%	4 hrs	80%
7	10 min.	92%	4 hrs	79%
8a	10 min.	90%	2hrs	75%
8b	10 min.	90%	2hrs	78%
8c	10 min.	90%	2hrs	75%
9a	10 min.	92%	2hrs	78%
9b	10 min.	94%	2hrs	82%
10	10 min.	90%	6hrs	74%
12a	10 min.	91%	4hrs	76%
12b	10 min.	90%	4hrs	75%
13	10min.	89%	6hrs	74%
17a	10 min.	92%	4hrs	75%
17b	10 min.	94%	4hrs	75%
17c	10 min.	90%	4hrs	75%
19	10 min.	90%	5hrs	75%
22a	10 min.	92%	4hrs	70%
22b	10 min.	93%	4hrs	70%
22c	10 min.	92%	4hrs	80%
22d	10 min.	90%	4hrs	70%
22e	10 min.	90%	4hrs	70%
22f	10 min.	93%	4hrs	71%

## Conclusion

Comparison between the above two methods showed that the solvent-free method (grinding method) was found to be better than the traditional method in terms of reaction time, yield, simplicity of the reaction procedure, ease of handling, low costs for performing the reaction and no solvents.

## Experiment

### Instrumentation

All melting points were determined on an electrothermal apparatus and are uncorrected. IR spectra were recorded (KBr discs) on a Shimadzu FT-IR 8201 PC spectrophotometer.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded in  $\text{CDCl}_3$  and  $(\text{CD}_3)_2\text{SO}$  solutions on a Varian Gemini 300 MHz and JNM-LA 400 FT-NMR system spectrometer and chemical shifts are expressed in ppm units using TMS as an internal reference. Mass spectra were recorded on a GC-MS QP1000 EX Shimadzu. Elemental analyses were carried out at the Microanalytical Center of Cairo University. Hydrazonoyl halides **11a-f** were prepared as previously reported [15-18].

### 1-(Benzofuran-2-yl)-3-(1,3-diphenyl-1H-pyrazol-4-yl)prop-2-en-1-one (3).

*Grinding method A:* 2-acetylbenzofuran (**1**) (1.6 g, 10 mmol), 1,3-diphenyl-1H-pyrazole-4-carbaldehyde (**2**) (1.24 g, 10 mmol), potassium hydroxide (0.56 g, 10 mmol) and few drops of water was ground with a pestle in an open mortar at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min while the reaction was monitored by TLC. The solid was washed with water and recrystallized from ethanol to give the corresponding **3**.

*Traditional method B:* Potassium hydroxide (10 mL, 10%) was added dropwise to a mixture of 2-acetylbenzofuran (**1**) and 1,3-diphenyl-1H-pyrazole-4-carbaldehyde (**2**) (10 mmol each) in ethanol (20 mL) while stirring at 0-5°C. The stirring was continued for 2 hrs. Then the resulting solid was collected and recrystallized from ethanol to give **3**. White crystals from ethanol. Yield: 88%, mp.: 168-170°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3064  $\nu(\text{CH-roma.})$ , 1660  $\nu(\text{CO})$ , 1610  $\nu(\text{C=N})$ , 1570  $\nu(\text{C=C})$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ,  $\delta$ , ppm): 7.10-8.17 (m, ArH's and  $-\text{CH}=\text{CH}-$ ). Anal. Calcd. for  $\text{C}_{26}\text{H}_{18}\text{N}_2\text{O}_2$  (390.43): C, 79.98; H, 4.65; N, 7.17. Found: C, 79.94; H, 4.61; N, 7.13%.

### Synthesis of pyridine derivatives 4-9

*Grinding Method A:* Equimolar amounts of **1** (0.8 g, 5 mmol), **2** (5 mmol) and the appropriate cyanothioacetamide, malononitrile, cyanoacetamide, or cyanoacid hydrazide (5 mmol) and ammonium acetate (0.77 g, 10 mmol) was ground with a pestle in an open mortar followed by the appropriate ethyl chloroacetate or iodomethane (10 mmol) at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min. while the reaction was monitored by TLC. The solid was washed with water and recrystallized from *N,N*-dimethylformamide to give the corresponding **4-7** respectively.

*Traditional Method B:* A mixture of 1-(benzofuran-2-yl)-3-(1,3-diphenyl-1H-pyrazol-4-yl)prop-2-en-1-one (**3**) (5 mmol), the appropriate 2-cyanoethanethioamide, malononitrile, cyanoacetamide,  $\omega$ -bromoacetophenone or cyanoacetohydrazide and ammonium acetate (5 mmol) was heated in acetic acid (10 mL) under reflux for 3 hrs. The reaction mixture was cooled; the separated solid was collected, filtered and crystallized from a proper solvent to give **4-7**.

*Traditional Method C:* A mixture of 2-acetylbenzofuran (**1**) (5 mmol), 1,3-diphenyl-1H-pyrazole-4-carbaldehyde (**2**) (5 mmol) and the appropriate 2-cyanoethanethioamide, malononitrile, cyanoacetamide,  $\omega$ -bromoacetophenone or cyanoacidhydrazide (5 mmol) and ammonium acetate (1.54 g, 20 mmol) was heated in *n*-butanol (10 mL) under reflux for 4 hrs. The excess solvent was evaporated and the residue was triturated with ethanol (10 mL). The solid formed was filtered off and crystallized from a proper solvent to give corresponding products which obtained in method A.

**6-(Benzofuran-2-yl)-2-mercapto-4-(1,3-diphenyl-1H-pyrazol-4-yl)pyridine-3-carbonitrile (4).** Orange crystals from ethanol. Yield: 85%, mp.: 288-90°C. FT-IR (KBr,  $\text{cm}^{-1}$ ): 3058  $\nu(\text{CH-roma.})$ , 2218  $\nu(\text{CN})$ , 1604  $\nu(\text{C=N})$ , 1234  $\nu(\text{SH})$ . Anal. Calcd. for  $\text{C}_{29}\text{H}_{18}\text{N}_4\text{OS}$  (470.54) C, 74.02; H, 3.86; N, 11.91; S, 6.81. Found: C, 74.10; H, 3.78; N, 11.96; S, 6.77. %.

**2-Amino-6-(benzofuran-2-yl)-4-(1,3-diphenyl-1H-pyrazol-4-yl)pyridin-3-carbonitrile (5).** Red crystals from ethanol. Yield: 75%, mp.: 89-90°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3288, 3200  $\nu(\text{NH}_2)$ , 3062  $\nu(\text{CH- arom.})$ , 2206  $\nu(\text{CN})$ , 1589  $\nu(\text{C=C})$ .  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 4.24 (s, 2H,  $\text{NH}_2$ ), 7.20-7.85 (m, 17H, ArH's). Anal. Calcd. For  $\text{C}_{29}\text{H}_{19}\text{N}_5\text{O}$  (453.49): C, 76.81; H, 4.22; N, 15.44. Found: 76.76; H, 4.33; N, 15.37%.

**1,2-Dihydro-6-(benzofuran-2-yl)-2-oxo-4(1,3-diphenyl-1H-pyrazol-4-yl)-pyridin-3-carbonitrile (6).** Yellow crystals from ethanol. Yield: 80%, mp.: 128-130°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3340  $\nu(\text{NH})$ , 3066  $\nu(\text{CH- arom.})$ , 2241  $\nu(\text{CN})$ , 1674  $\nu(\text{CO})$ , 1604  $\nu(\text{C=N})$ .  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 7.22-7.81 (m, 17H, ArH's), 11.21 (s, H, NH). Anal. Calcd. For  $\text{C}_{29}\text{H}_{18}\text{N}_4\text{O}_2$  (454.48): C, 76.64; H, 3.99; N, 12.33. Found: 76.75; H, 4.10; N, 12.24%.

**6-(benzofuran-2-yl)-2-hydrazinyl-4-(1,3-diphenyl-1H-pyrazol-4-yl)pyridine-3-carbonitrile (7).** Yellow crystals from ethanol. Yield: 79, m.p.: 220°C. FT-IR (KBr,  $\text{cm}^{-1}$ ): 3290, 3200  $\nu(\text{NH}_2)$ , 3062  $\nu(\text{CH- arom.})$ , 2210  $\nu(\text{CN})$ , 1689  $\nu(\text{C=O})$ . Anal. Calcd. for  $\text{C}_{29}\text{H}_{19}\text{N}_5\text{O}_2$  (469.49): C, 74.19; H, 4.08; N, 14.92. Found: 74.15; H, 4.04; N, 14.97%.

**2-(Substituted) 6-(benzofuran-2-yl)-4-(1,3-diphenyl-1H-pyrazol-4-yl)-pyridine-3-carbonitrile 8a-c.**

**Grinding Method:** Equimolar amounts of **4** (2.35 g, 5 mmol) and potassium hydroxide (0.28 g, 5 mmol) was ground with a pestle in an open mortar followed by the appropriate iodomethane, ethyl chloroacetate or chloroacetone (5 mmol) at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min. while the reaction was monitored by TLC. The solid was washed with water and recrystallized from *N,N*-dimethylformamide to give the corresponding **8a-c**, respectively.

**Traditional Method:** A mixture of **4** (2.35 g, 5 mmol) and potassium hydroxide (0.28 g, 5 mmol) in *N,N*-dimethylformamide (10 mL) was stirred for 2 hrs at room temperature. The appropriate of iodomethane, ethyl chloroacetate or chloroacetone (5 mmol) was added while stirring. Stirring was continued for 2 hrs. The resulting solid was collected and crystallized from *N,N*-dimethylformamide to afford **8a-c** respectively.

**6-(Benzofuran-2-yl)-2-(methylthio)-4-(1,3-diphenyl-1H-pyrazol-4-yl)-pyridin--3-carbonitrile (8a).** Yellow crystals from *N,N*-dimethylformamide. Yield: 75%, mp.: 234-36°C. FT-IR (KBr,  $\text{cm}^{-1}$ ): 3055  $\nu(\text{CH- aroma.})$ , 2923, 2854  $\nu(\text{CH- aliph.})$ , 2210  $\nu(\text{CN})$ , 1596  $\nu(\text{C=C})$ .  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 2.77 (s, 3H,  $\text{SCH}_3$ ), 7.27-8.49 (m, 17H, ArH's). Anal. Calcd. for  $\text{C}_{30}\text{H}_{20}\text{N}_4\text{OS}$  (484.57): C, 74.36; H, 4.16; N, 11.56; S, 6.62. Found: 74.42; H, 4.07; N, 11.44; S, 6.70 %.

**Ethyl 6-(benzofuran-2-yl)-4-(1,3-diphenyl-1H-pyrazol-4-yl)-pyridin-2-ylthioacetate-3-carbonitrile (8b).** Yellow crystals from *N,N*-dimethylformamide. Yield: 78%, mp.: 200-02°C. FT-IR (KBr,  $\text{cm}^{-1}$ ): 3058  $\nu(\text{CH- aroma.})$ , 2981, 2931  $\nu(\text{CH- aliph.})$ , 2214  $\nu(\text{CN})$ , 1732  $\nu(\text{C=O})$ , 1596  $\nu(\text{C=C})$ .  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 1.20 (t, 3H,  $J = 7$  Hz,  $\text{CH}_2\text{CH}_3$ ), 4.22 (q, 2H,  $J = 7$  Hz,  $\text{CH}_2\text{CH}_3$ ), 4.70 (s, 2H,  $\text{SCH}_2$ ), 7.22-8.10 (m, 17H, ArH's); MS (EI,  $m/z$  (%)): 557 (M+1, 25), 556 (M+, 33.3), 512 (33.3), 85 (29.2), 73 (100). Anal. Calcd. for  $\text{C}_{33}\text{H}_{24}\text{N}_4\text{O}_3\text{S}$  (556.63): C, 71.21; H, 4.35; N, 10.07; S, 5.76 Found: 71.29; H, 4.44; N, 10.13; S, 5.80 %.

**S-6-(benzofuran-2-yl)-3-cyano-4-(1,3-diphenyl-1H-pyrazol-4-yl)pyridin-2-yl ethanethioate (8c).** Yellow crystals from *N,N*-dimethylformamide. Yield: 75%, mp.: 100-02°C. FT-IR (KBr,  $\text{cm}^{-1}$ ): 3062  $\nu(\text{CH- aroma.})$ , 2931, 29854  $\nu(\text{CH- aliph.})$ , 2194  $\nu(\text{CN})$ , 1666  $\nu(\text{C=O})$ , 1596  $\nu(\text{C=C})$ .  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): (s, 3H,  $\text{COCH}_3$ ), 4.70 (s, 2H,  $\text{SCH}_2$ ), 7.20-8.15 (m, 17H, ArH's). Anal. Calcd. for  $\text{C}_{32}\text{H}_{22}\text{N}_4\text{O}_2\text{S}$  (526.61): C, 72.98; H, 4.21; N, 10.64; S, 6.09. Found: C, 72.98; H, 4.21; N, 10.64; S, 6.09 %.

**2-(Substituted-3-amino-6-(benzofuran-2-yl)-4-(1,3-diphenyl-1H-pyrazol-4-yl)thieno[2,3-b]pyridine (9a,b)**

**Grinding Method:** Equimolar amounts of **4** (2.35 g, 5 mmol) and potassium hydroxide (0.28 g, 5 mmol) was ground with a pestle in an open mortar followed by the appropriate phenacyl bromide or chloroacetonitrile (10 mmol) at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min. while the reaction was monitored by TLC. The solid was washed with water and recrystallized from *N,N*-dimethylformamide to give the corresponding **9a,b**, respectively.

*Traditional Method:* A mixture of **4** (2.35 g, 5 mmol) and potassium hydroxide (0.28 g, 5 mmol) in *N,N*-dimethylformamide (10 mL) was stirred for 2 hrs at room temperature. The appropriate of phenacyl bromide or chloroacetonitrile (10 mmol) was added while stirring. Stirring was continued for 2 hrs. The resulting solid was collected and crystallized from *N,N*-dimethylformamide to afford **9a,b** respectively.

**(3-amino-6-(benzofuran-2-yl)-4-(1,3-diphenyl-1H-pyrazol-4-yl)thieno[2,3-b]pyridin-2-yl)(phenyl)methanone(9a)**

Yellow crystals from ethanol. Yield: 76%. mp.: 199-200°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3370, 3286  $\nu(\text{NH}_2)$ , 3058  $\nu(\text{CH-roma.})$ , 1666  $\nu(\text{C=O})$ .  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 6.24 (s, 2H,  $\text{NH}_2$ ), 7.20-7.85 (m, 22H, ArH's). Anal. Calcd. for  $\text{C}_{37}\text{H}_{24}\text{N}_4\text{O}_2\text{S}$  (588.68): C, 75.49; H, 4.11; N, 9.52; S, 5.45. Found: C, 75.45; H, 4.15; N, 9.57; S, 5.41%.

**3-amino-6-(benzofuran-2-yl)-4-(1,3-diphenyl-1H-pyrazol-4-yl)thieno[2,3-b]pyridine-2-carbonitrile (9b)**

Yellow crystals from DMF. Yield: 78%. mp.: >300°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3463, 3332  $\nu(\text{NH}_2)$ , 3058  $\nu(\text{CH-roma.})$ , 2202  $\nu(\text{CN})$ , 1627  $\nu(\text{C=N})$ .  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 5.60 (s, 2H,  $\text{NH}_2$ ), 7.20-7.85 (m, 17H, ArH's). Anal. Calcd. for  $\text{C}_{31}\text{H}_{19}\text{N}_5\text{OS}$  (509.58): C, 73.07; H, 3.76; N, 13.74; S, 6.29. Found: C, 73.10; H, 3.72; N, 13.70; S, 6.25%.

**3-(benzofuran-2-yl)-4,5-dihydro-5-(1,3-diphenyl-1H-pyrazol-4-yl)pyrazole-1-carboxamide(10)**

*Grinding method:* Chalcone **3** (1.95 g, 5 mmol), thiosemicarbazide (0.485 g, 5 mmol), sodium hydroxide (0.4 g, 10 mmol) and few drops of water was ground with a pestle in an open mortar at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min while the reaction was monitored by TLC. The solid was washed with water and recrystallized from ethanol to give the corresponding **10**.

*Traditional method:* A mixture of chalcone **3** (1.95 g, 5 mmol) and thiosemicarbazide (0.485 g, 5 mmol) in acetic acid (20 mL) was heated under reflux for 6 hrs. The resulting solid that obtained after cooling was collected and recrystallized from ethanol to give **10** Page crystals from ethanol. Yield: 76%. mp.: 205-07°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3417, 3367  $\nu(\text{NH}_2)$ , 3058  $\nu(\text{CH-roma.})$ , 1600  $\nu(\text{C=N})$ . Anal. Calcd. for  $\text{C}_{27}\text{H}_{21}\text{N}_5\text{OS}$  (463.55): C, 69.96; H, 4.57; N, 15.11; S, 6.92. Found: C, 69.91; H, 4.53; N, 15.15; S, 6.96%.

**1-(2-(3-(benzofuran-2-yl)-4,5-dihydro-5-(1,3-diphenyl-1H-pyrazol-4-yl)pyrazol-1-yl)-4-substituted 1,3-thiazol-5-yl)-2-phenyldiazene (12a,b)**

*Grinding method:* carboxamide **10** (2.32 g, 5 mmol), the appropriate hydrazonoyl halides **11c,d** (5 mmol), sodium carbonate (1 g, 10 mmol) and few drops of water was ground with a pestle in an open mortar at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min while the reaction was monitored by TLC. The solid was washed with water and recrystallized from ethanol to give the corresponding **12a** and **12b**, respectively.

*Traditional method:* A mixture of **10** (2.32 g, 5 mmol) and hydrazonoyl halides **11c,d** (5 mmol) in ethanol containing triethylamine was heated under reflux for 4 hrs. The resulting solid that obtained after cooling was collected and recrystallized from ethanol to give **12a,b**, respectively.

**2-(5-(benzofuran-2-yl)-3',5'-diphenyl-3,4-dihydro-1'H,2H-[3,4'-bipyrazol]-2-yl)-4-methyl-5-(phenyldiazanyl)thiazole (12a)**

Red crystals from ethanol. Yield: 76%. mp.: 220°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3062, 2927  $\nu(\text{CH})$ , 1608  $\nu(\text{C=N})$ .  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ ):  $\delta$  = 2.53 (s, 3H,  $\text{CH}_3$ ), 3.20 (dd, 1H,  $J = 18.1, 5.8\text{Hz}$ ,  $\text{CH}_2(\text{pyraz})$ ), 3.82 (dd, 1H,  $J = 18.1, 12.0\text{Hz}$ ,  $\text{CH}_2(\text{pyraz})$ ), 5.23 (dd, 1H,  $J = 12.2, 5.8\text{Hz}$ ,  $\text{CH}(\text{pyraz})$ ), 7.20-7.75 (m, 21H, ArH's). Anal. Calcd. For  $\text{C}_{36}\text{H}_{27}\text{N}_7\text{OS}$  (605.71): C, 71.38; H, 4.49; N, 16.19; S, 5.29. Found: C, 71.34; H, 4.43; N, 16.15; S, 5.25%.

**2-(5-(benzofuran-2-yl)-3',5'-diphenyl-3,4-dihydro-1'H,2H-[3,4'-bipyrazol]-2-yl)-4-phenyl-5-(phenyldiazanyl)thiazole (12b)**

Red crystals from ethanol. Yield: 75%, mp.: 230°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3058, 2927  $\nu(\text{CH})$ , 1600  $\nu(\text{C=N})$ . Anal. Calcd. for  $\text{C}_{41}\text{H}_{29}\text{N}_7\text{OS}$  (667.78): C, 73.74; H, 4.38; N, 14.68; S, 4.80. Found: C, 73.70; H, 4.34; N, 14.72; S, 4.84%.



**4-(3-(benzofuran-2-yl)-4,5-dihydro-1-(4-phenylthiazol-2-yl)-1H-pyrazol-5-yl)-1,3-diphenyl-1H-pyrazole (13)**

*Grinding method:* Chalcone **3** (5 mmol), 1-(4-phenylthiazol-2-yl)hydrazine (**14**) (1.45 g, 5 mmol), sodium hydroxide (10 mmol) and few drops of water was ground with a pestle in an open mortar at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min while the reaction was monitored by TLC. The solid was washed with water and recrystallized from ethanol to give the corresponding **13**.

*Traditional method A:* A mixture of chalcone **3** (1.95 g 5 mmol) and 1-(4-phenylthiazol-2-yl) hydrazine (**14**) (1.45 g, 5 mmol) in acetic acid (20 mL) was heated under reflux for 6 hrs. The resulting solid that obtained after cooling was collected and recrystallized from ethanol to give **13**.

*Alternative method:* A mixture of **10** ((2.32 g, 5 mmol) and  $\omega$ -bromoacetophenone (1 g, 5 mmol) in ethanol containing triethylamine was heated under reflux for 4 hrs. The resulting solid that obtained after cooling was collected and recrystallized from ethanol to give **13**.

Brown crystals from ethanol. Yield: 74%, mp.: 160°C; FT-IR (KBr,  $\text{cm}^{-1}$ ): 3061, 2929  $\nu$  (CH), 1600  $\nu$  (C=N).  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-}d_6$ ):  $\delta$  = 3.23 (dd, 1H,  $J$  = 18.1, 5.8 Hz,  $\text{CH}_2$ (pyraz)), 3.95 (dd, 1H,  $J$  = 18.1, 12.0 Hz,  $\text{CH}_2$ (pyraz)), 5.33 (dd, 1H,  $J$  = 12.2, 5.8 Hz, CH(pyraz)), 6.58 (s, 1H, thiazole H-5), 7.25-7.72 (m, 21H, ArH's). Anal. Calcd. for  $\text{C}_{35}\text{H}_{25}\text{N}_5\text{OS}$ (563.67): C, 74.58; H, 4.47; N, 12.42; S, 5.69. Found: C, 74.54; H, 4.43; N, 12.46; S, 5.65%.

**Synthesis of 3-substituted 5-(benzofuran-2-yl)4-cyano-1-phenyl-1H-pyrazole (17a-c).**

*Grinding Method:* A mixture of **15** (0.925 g, 5 mmol), the appropriate hydrazonoyl halide **11a-c** (5 mmol), sodium carbonate (1 g, 10 mmol) and few drops of water was ground with a pestle in an open mortar at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min while the reaction was monitored by TLC. The solid was washed with water and recrystallized from the appropriate solvent to give the corresponding **17a-c**, respectively.

*Traditional Method:* Compound **15** (0.925 g, 5 mmols) was added to a stirred ethanolic sodium ethoxide solution (0.12 g-atom sodium metal in absolute ethanol 20 mL). After 20 min., the appropriate hydrazonoyl halide **11a-c** (5 mmols) was added and the reaction mixture was stirred for 4 h. The resulting solid was collected by filtration, dried and recrystallized from a proper solvent to give **17a-c**, respectively.

**Ethyl 5-(benzofuran-2-yl)-4-cyano-1-phenyl-1H-pyrazole-3-carboxylate (17a).** Yellow crystals from ethanol. Yield: 80%, mp.: 174°C. FT-IR (KBr,  $\text{cm}^{-1}$ ): 3066  $\nu$ (CH-aroma.), 2985, 2912  $\nu$ (CH-aliph.), 2229  $\nu$ (CN), 1724  $\nu$ (CO), 1596  $\nu$ (C=C).  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 1.33 (t, 3H,  $J$  = 7 Hz,  $\text{CH}_2\text{CH}_3$ ), 4.41 (q, 2H,  $J$  = 7 Hz,  $\text{CH}_2\text{CH}_3$ ), 6.93 (s, 1H, CH-furan), 7.41-7.63 (m, 9H, ArH's). Anal. Calcd. for  $\text{C}_{21}\text{H}_{15}\text{N}_3\text{O}_3$  (357.36): C, 70.58; H, 4.23; N, 11.76. Found: 70.50; H, 4.30; N, 11.85%.

**3-Acetyl-5-(benzofuran-2-yl)-1-phenyl-1H-pyrazole-4-carbonitrile (17b)** Yellow crystals from acetic acid. Yield: 82 %, mp.: 220°C. FT-IR (KBr,  $\text{cm}^{-1}$ ): 3062  $\nu$ (CH-arom.), 2237  $\nu$ (CN), 1693  $\nu$ (CO), 1604  $\nu$ (C=N).  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 2.68 (s, 3H,  $\text{CH}_3$ ), 7.05 (s, 1H, CH-furan), 7.25-7.61 (m, 9H, ArH's). Anal. Calcd. for  $\text{C}_{20}\text{H}_{13}\text{N}_3\text{O}$  (327.34): C, 73.38; H, 4.00; N, 12.84. Found: C, 73.45; H, 4.10; N, 12.73%.

**3-(Benzofuran-2-yl-carbonyl)-5-(benzofuran-2-yl)-1-phenyl-1H-pyrazole-4-carbonitrile (17c).** Red crystals from ethanol. Yield: 75%, mp.: 210-12°C.

$^1\text{H}$  NMR (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 7.33-7.61 (m, 15H, ArH's). Anal. Calcd. for  $\text{C}_{27}\text{H}_{15}\text{N}_3\text{O}_3$  (429.43): C, 75.52; H, 3.52; N, 9.79. Found: C, 75.44; H, 3.63; N, 9.69%.

**4-amino-3-(benzofuran-2-yl)-2-phenyl-2H-pyrazolo[3,4-d]pyridazin-7(6H)-one (19)**

*Grinding method:* A mixture of **17a** (1.8 g, 5 mmol) and hydrazine hydrate (0.5 g, 10 mmol) was ground with a pestle in an open mortar at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy reaction mixture solidified within 3-5 min. Grinding was continued for 5-10 min while the reaction was monitored by TLC. The solid was washed with water and recrystallized from ethanol to give the corresponding **19**.

*Traditional method:* A mixture of **17a** (1.8 g, 5 mmol) and hydrazine hydrate (0.5 g, 10 mmol) in ethanol (20 mL) was heated under reflux for 5 hrs. The resulting solid that obtained after cooling was collected and recrystallized from ethanol to give **19**.

White crystals from ethanol. Yield: 75%, mp.: 273-74°C.  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-}d_6$ ,  $\delta$ , ppm): 6.32 (br, 2H,  $\text{NH}_2$ ), 7.33-7.61 (m, 10H, ArH's), 9.50(s, 1H, NH). Anal. Calcd. for  $\text{C}_{19}\text{H}_{13}\text{N}_5\text{O}_2$ (343.34): C, 66.47; H, 3.82; N, 20.40. Found: C, 66.43; H, 3.86; N, 20.45%.

### Synthesis of 2,3-dihydro 1,3,4-thiadiazoles 22a-f

**Grinding Method :** 3-(benzofuran-2-yl)-3-oxopropanenitrile **15** (1.84 g, 10 mmol), phenyl isothiocyanate (1.37 g, 10 mmol) and potassium hydroxide (0.56 g, 10 mmol) was ground with a pestle in an open mortar followed by the appropriate hydrazonoyl halides **11a-c** (10 mmol) at room temperature for 2-3 min. until the mixture turned into a melt. The initial syrupy solidified within 3-5 min. Grinding was continued for 5-10 min. while the reaction was monitored by TLC. The solid was washed with ethanol-water and recrystallized from a proper solvent to afford **22a-f**, respectively.

**Traditional Method:** A mixture of Compound **15** (1.84 g, 10 mmol), phenyl isothiocyanate (1.37 g, 10 mmol) and potassium hydroxide (0.56 g, 10 mmol) in *N, N*-dimethylformamide (10 mL) was stirred for 2 hrs at room temperature. The appropriate hydrazonoyl halides **11a-c** (10mmol) was added while stirring. Stirring was continued for 2 hrs. The resulting solid was collected and crystallized from a proper solvent to afford **22a-c**, respectively.

#### Ethyl 5-(2-(benzofuran-2-yl)-1-cyano-2-oxoethylidene)-4-phenyl-4,5-dihydro-1,3,4-thiadiazole-2-carboxylate (**22a**)

Yellow crystals from ethanol. Yield: 70 %, mp.: 210-12 °C (lit mp. 210 [19]). FT-IR (KBr,  $\text{cm}^{-1}$ ): 3062  $\nu$ (CH-arom.), 2190  $\nu$ (CN), 1720, 1680  $\nu$ (CO).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ,  $\delta$ , ppm): 1.36 (t, 3H,  $J = 7.5$  Hz,  $\text{CH}_2\text{CH}_3$ ), 4.2 (q, 2H,  $J = 7.5$  Hz,  $\text{CH}_2\text{CH}_3$ ), 7.10 – 7.92 (m, 10H, ArH's). Anal. Calcd. for  $\text{C}_{22}\text{H}_{15}\text{N}_3\text{O}_4\text{S}$  (417.44): C, 63.30; H, 3.62; N, 10.07; S, 7.68. Found: C, 63.15; H, 3.52; N, 10.21, S, 7.86 %

#### 2-(5-Acetyl-3-phenyl-1,3,4-thiadiazol-2(3H)-ylidene)-3-(benzofuran-2-yl)-3-oxopropanenitrile (**22b**)

Red crystals from ethanol. Yield: 70 %, mp.: 200-202 °C (lit mp. 200 [19]). FT-IR (KBr,  $\text{cm}^{-1}$ ): 3062  $\nu$ (CH-arom.), 2190  $\nu$ (CN), 1665, 1655  $\nu$ (CO).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ,  $\delta$ , ppm): 2.11 (s, 3H,  $\text{CH}_3$ ), 7.10- 7.92 (m, 10H, ArH's). Anal. Calcd. for  $\text{C}_{21}\text{H}_{13}\text{N}_3\text{O}_3\text{S}$  (387.41): C, 65.11; H, 3.38; N, 10.85; S, 8.28. Found: C, 65.00; H, 3.41; N, 10.99, S, 8.41 %

#### 3-(Benzofuran-2-yl)-2-(5-benzoyl-3-phenyl-1,3,4-thiadiazol-2(3H)-ylidene)-3-oxopropanenitrile (**22c**)

Red crystals from ethanol. Yield: 80 %, mp.: 181-82 °C (lit mp. 181 [19]). FT-IR (KBr,  $\text{cm}^{-1}$ ): 3062  $\nu$ (CH-arom.), 2190  $\nu$ (CN), 1680, 1660  $\nu$ (CO).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ,  $\delta$ , ppm): 7.10 – 7.92 (m, 15H, ArH's). Anal. Calcd. for  $\text{C}_{26}\text{H}_{15}\text{N}_3\text{O}_3\text{S}$  (449.48): C, 69.48; H, 3.36; N, 9.35; S, 7.13. Found: C, 69.51; H, 3.27; N, 9.53, S, 7.25 %

#### 2-(5-(Benzofuran-2-carbonyl)-3-phenyl-1,3,4-thiadiazol-2(3H)-ylidene)-3-(benzofuran-2-yl)-3-oxopropanenitrile (**22d**)

Red crystals from ethanol. Yield: 70 %, mp.: 220-22 °C (lit mp. 220 [19]). FT-IR (KBr,  $\text{cm}^{-1}$ ): 3062  $\nu$ (CH-arom.), 3220  $\nu$ (NH), 2190  $\nu$ (CN), 1680, 1665  $\nu$ (CO).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ,  $\delta$ , ppm): 7.10 – 8.24 (m, 15H, ArH's). Anal. Calcd. for  $\text{C}_{28}\text{H}_{15}\text{N}_3\text{O}_4\text{S}$  (489.50): C, 68.70; H, 3.09; N, 8.58; S, 6.55. Found: C, 68.64; H, 2.95; N, 8.47, S, 6.72 %.

#### 5-(2-(Benzofuran-2-yl)-1-cyano-2-oxoethylidene)-N,4-diphenyl-4,5-dihydro-1,3,4-thiadiazole-2-carboxamide (**22e**)

Yellow crystals from ethanol. Yield: 70 %, mp.: 191-92 °C (lit mp. 192 [19]). FT-IR (KBr,  $\text{cm}^{-1}$ ): 3062  $\nu$ (CH-arom.), 3220  $\nu$ (NH), 2190  $\nu$ (CN), 1680, 1665  $\nu$ (CO).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ,  $\delta$ , ppm): 7.10 – 7.92 (m, 15H, ArH's), 9.31 (s, br., 1H, NH). Anal. Calcd. for  $\text{C}_{26}\text{H}_{16}\text{N}_4\text{O}_3\text{S}$  (464.50): C, 67.23; H, 3.47; N, 12.06; S, 6.90. Found: C, 67.42; H, 3.63; N, 11.89, S, 6.81 %.

#### 2-(5-(Benzofuran-2-yl-carbonyl)-3-*p*-tolyl-1,3,4-thiadiazol-2(3H)-ylidene)-3-(benzofuran-2-yl)-3-oxopropanenitrile (**22f**)

Brown crystals from acetic acid. Yield: 85% mp.: 220-22°C;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ - $d_6$ ,  $\delta$ , ppm): 2.520(s, 3H,  $\text{CH}_3$ ), 6.90 (s, 1H, CH-furan) and 7.08-8.30 (m, 13H, ArH's). Anal. Calcd. for  $\text{C}_{29}\text{H}_{17}\text{N}_3\text{O}_4\text{S}$  (503.53): C, 69.17; H, 3.40; N, 8.35; S, 6.37. Found: C, 69.10; H, 3.46; N, 8.41; S, 6.32%.

## Conclusion

A convenient, efficient and rapid method was developed for synthesis of pyridines, pyrazoles, thiadiazoline and 2,3-dihydro 1,3,4-thiadiazoles in a good to excellent yields by a grinding method under catalyst- and solvent-free conditions. This procedure is much simpler and faster than the traditional one. It is also consistent with a green chemistry approach since no solvent is needed.

## References

- [1] Mikami, K. *Green Reaction Media in Organic Synthesis*: (2005). Blackwell, Oxford.
- [2] Tanaka, K., Ed. (2004): *Solvent-Free Organic Synthesis*: Wiley-VCH, Weinheim.
- [3] Tanaka; Toda, F. K. (2000): *Solvent-Free Organic Synthesis*. *Chem. Rev.*, 100: 1025-1074.
- [4] Schmeyers, J.; Toda, F.; Boy, J.; Kaupp, G. (1998): Quantitative solid–solid synthesis of azomethines. *J. Chem. Soc., Perkin Trans.*, 2: 989-994.
- [5] Santana, L.; Teijeira, M.; Uriarte, E.; Teran, C.; Linares, B.; Villar, R.; Laguna, R.; Cano, E. (1998): AM1 theoretical study, synthesis and biological evaluation of some benzofuran analogues of anti-inflammatory arylalkanoic acids. *Eur. J. Pharm. Sci.*, 7: 161-166.
- [6] Findlay, J. A.; Buthelezi, S.; Li, G.; Seveck, M. (1997): Insect Toxin from an endophytic fungus from wintergreen. *J. Nat. Prod.*, 60(11): 1214-1215.
- [7] Leonardi, A.; Nava, G.; Nardi, D. (1983): Synthesis of benzofuran derivatives with H<sub>2</sub> antagonist activity. *Farmaco. Ed. Sci.*; 38(5): 290-308.
- [8] Dawood, K. M.; Abdel-Gawad, H.; Rageb, E. A.; Ellithey M.; Mohamad, H. A. (2006): Synthesis, anticonvulsant, and anti-inflammatory evaluation of some new benzotriazole and benzofuran based heterocycles. *Bioorg. Med. Chem.*, 14: 3672-3680.
- [9] Musser, J. H.; Brown, R. E.; Love, B.; Bailey, K.; Jones, H.; Kahen, R.; Huang, F. C.; Khandurala, A.; Leibowitz, M.; Sonniua-Goldman, P.; Donigi-Ruzza, D. (1984): Synthesis of 2-(2,3-dihydro-2-oxo-1,3,4-oxadiazol-5-yl) benzo heterocycles. A novel series of orally active antiallergic agents. *J. Med. Chem.*, 27: 121-125.
- [10] Rida, S. M.; El-Hawash, S. A. M.; Fahmy, H. T. Y.; Hazzaa, A. A.; ElMeligy, M. M. M. (2006): Synthesis of Novel Benzofuran and Related Benzimidazole Derivative for Evaluation of In Vitro Anti-HIV-1, Anticancer and Antimicrobial Activites. *Arch Pharm Res.*; 29(10): 826-833.
- [11] Raj, P. A.; Suddendra, G.; Shakeeland, A. S.; Girish, M. (2012): Synthesis of New Benzofuran Derivatives and their in Vitro evaluation for Antimicrobial Activity. *Inter J. OF Drug Formulation and Res.*, 3: 135-147
- [12] Yoo, S. E.; Lee, S. H.; Kim, S. K. (1997): The Conformation and activity Relationship of benzofuran derivative as Angiotension II Receptor antagonist. *Bioorg. Med. Chem.*, 5: 445-459.
- [13] Yilmaz, M.; Uzunalioglu, N.; Pekel, A. T. (2005): Manganese(III) acetate based oxidative cyclizations of 3-oxopropanenitriles with conjugated alkenes and synthesis of 4,5-dihydrofuran-3-carbonitriles containing heterocycles. *Tetrahedron*, 61: 8860.
- [14] Yılmaz, E. V. B.; Yılmaz, M.; Öktemer, A. (2011): Radical cyclizations of conjugated esters and amides with 3-oxopropanenitriles mediated by manganese (III) acetate. *011, Akkivoc*, (ii): 363-376.
- [15] Eweiss, N. E.; Osman, A. (1979): Synthesis of Heterocycles. One Step Synthesis of Acetylthiadiazolines. *Tetrahedron Lett.* 13: 1169-1170.

[16] Shawali, A. S.; Abdelhamid, A. O. (1976): Reaction of Dimethylphenacylsulfonium Bromide with *N*-Nitrosoacetaryl amides and Reactions of the Products with Nucleophiles. Bull. Soc. Chim. Jpn., 49, 321-324.

[17] Shawali, A. S.; Osman, A. (1971): Synthesis and Reactions of Phenylcarbamoyl-arylhydrazidic Chlorides. Tetrahedron, 27: 2517-2528.

[18] Abdelhamid, A. O.; Attaby, F. A.; Zaki, M. Y. (1990): Reactions with 2-(thiocyanatoacetyl) and 2-(selenacyanoacetyl)-2'-benzofuran. Synthesis of some new thiadiazoline, selenadiazoline, thiadiazolo[2,3-*b*]quinazoline and arylazothiazole derivatives. Phosphorous, sulfur, silicon, and Related element, 53: 403-410

[19] Abdelhamid, A. O.; Ismail, Z.H.; Abdel-Gawad, S. M.; Ghorab, M. M.; Abdel-Aziem, A. (2009): Synthesis of Some New Thieno[2,3-*b*]pyridines, Pyrimidino[4,5:4,5]thieno[2,3-*b*]pyridines, and 2,3-Dihydro-1,3,4-thiadiazoles. Phosphorus, Sulfur, Silicon, and Related element, 184: 58-75