

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2020 November 21; 26(43): 6706-6908



**OPINION REVIEW**

- 6706 Gastric acid level of humans must decrease in the future  
*Fujimori S*

**EXPERT CONSENSUS**

- 6710 First United Arab Emirates consensus on diagnosis and management of inflammatory bowel diseases: A 2020 Delphi consensus  
*Alkhatry M, Al-Rifai A, Annese V, Georgopoulos F, Jazzar AN, Khassouan AM, Koutoubi Z, Nathwani R, Taha MS, Limdi JK*

**MINIREVIEWS**

- 6770 Hepatocellular carcinoma after direct-acting antiviral hepatitis C virus therapy: A debate near the end  
*Muzica CM, Stanciu C, Huiban L, Singeap AM, Sfarti C, Zenovia S, Cojocariu C, Trifan A*

**ORIGINAL ARTICLE****Basic Study**

- 6782 *Lactobacillus bulgaricus* inhibits colitis-associated cancer via a negative regulation of intestinal inflammation in azoxymethane/dextran sodium sulfate model  
*Silveira DSC, Veronez LC, Lopes-Júnior LC, Anatriello E, Brunaldi MO, Pereira-da-Silva G*
- 6795 Antifungal activity and antidiarrheal activity via antimotility mechanisms of (-)-fenchone in experimental models  
*Pessoa MLS, Silva LMO, Araruna MEC, Serafim CAL, Alves Júnior EB, Silva AO, Pessoa MMB, Diniz Neto H, de Oliveira Lima E, Batista LM*
- 6810 Effects of Yue-Bi-Tang on water metabolism in severe acute pancreatitis rats with acute lung-kidney injury  
*Hu J, Zhang YM, Miao YF, Zhu L, Yi XL, Chen H, Yang XJ, Wan MH, Tang WF*
- 6822 Tissue microarray-chip featuring computerized immunophenotypical characterization more accurately subtypes ampullary adenocarcinoma than routine histology  
*Palmeri M, Funel N, Di Franco G, Furbetta N, Gianardi D, Guadagni S, Bianchini M, Pollina LE, Ricci C, Del Chiaro M, Di Candio G, Morelli L*

**Retrospective Cohort Study**

- 6837 Feasibility and nutritional impact of laparoscopic assisted tailored subtotal gastrectomy for middle-third gastric cancer  
*Liu H, Jin P, Ma FH, Ma S, Xie YB, Li Y, Li WK, Kang WZ, Tian YT*
- 6853 Role of doublecortin-like kinase 1 and leucine-rich repeat-containing G-protein-coupled receptor 5 in patients with stage II/III colorectal cancer: Cancer progression and prognosis  
*Kang XL, He LR, Chen YL, Wang SB*

**Clinical Trials Study**

- 6867 Comparison of two supplemental oxygen methods during gastroscopy with propofol mono-sedation in patients with a normal body mass index

*Shao LJZ, Zou Y, Liu FK, Wan L, Liu SH, Hong FX, Xue FS*

**Observational Study**

- 6880 Barriers for resuming endoscopy service in the context of COVID-19 pandemic: A multicenter survey from Egypt

*Elshaarawy O, Lashen SA, Makhlouf NA, Abdeltawab D, Zaghloul MS, Ahmed RM, Fathy H, Afifi S, Abdel-Gawad M, Abdelsameea E, Abd-Elsalam S, Mohamed SY, Tag-Adeen M, Tharwat M, Alzamzamy A, Bekhit AN, Eid AM, Awad A, Aamr M, Abd El Dayem WA, Wifi MN, Alborai M*

**SYSTEMATIC REVIEWS**

- 6891 Crohn's disease in low and lower-middle income countries: A scoping review

*Rajbhandari R, Blakemore S, Gupta N, Adler AJ, Noble CA, Mannan S, Nikolli K, Yih A, Joshi S, Bukhman G*

**ABOUT COVER**

Editorial Board Member of *World Journal of Gastroenterology*, Dr. Vasiliy I Reshetnyak is a Distinguished Professor at the AI Yevdokimov Moscow State University of Medicine and Dentistry in Moscow, Russia. In 1977, he graduated with honors from the Medical Faculty of IM Sechenov 1st Moscow Medical Institute. In 1982m Dr. Reshetnyak defended his PhD thesis. Working at the Central Research Institute of Gastroenterology (CRIG) led to obtainment of his doctoral thesis (DSc) in 1996 and his rise to Deputy Director on Scientific Work at the CRIG. Since 2003, Professor Reshetnyak has served as an Academic Secretary of the Research Institute of General Reanimatology. His scientific research activity has focused on the diagnosis, clinical and laboratory manifestations, and treatment of gastroenterological diseases with patients in critical states. Since 2019, Dr. Reshetnyak has held the rank of Professor of the Department of Propaedeutic of Internal Diseases and Gastroenterology. (L-Editor: Filipodia)

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastroenterology (WJG, World J Gastroenterol)* is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJG* mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

**INDEXING/ABSTRACTING**

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2020 edition of Journal Citation Report® cites the 2019 impact factor (IF) for *WJG* as 3.665; IF without journal self cites: 3.534; 5-year IF: 4.048; Ranking: 35 among 88 journals in gastroenterology and hepatology; and Quartile category: Q2.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Yu-Jie Ma*; Production Department Director: *Xiang Li*; Editorial Office Director: *Ze-Mao Gong*.

**NAME OF JOURNAL**

*World Journal of Gastroenterology*

**ISSN**

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

**LAUNCH DATE**

October 1, 1995

**FREQUENCY**

Weekly

**EDITORS-IN-CHIEF**

Andrzej S Tarnawski, Subrata Ghosh

**EDITORIAL BOARD MEMBERS**

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

**PUBLICATION DATE**

November 21, 2020

**COPYRIGHT**

© 2020 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Observational Study

**Barriers for resuming endoscopy service in the context of COVID-19 pandemic: A multicenter survey from Egypt**

Omar Elshaarawy, Sameh Aldesoky Lashen, Nahed A Makhlof, Doaa Abdeltawab, Mariam Salah Zaghloul, Rasha M Ahmed, Hayam Fathy, Shimaaf Afifi, Muhammad Abdel-Gawad, Eman Abdelsameea, Sherief Abd-Elsalam, Salem Youssef Mohamed, Mohammed Tag-Adeen, Mina Tharwat, Ahmed Alzamzamy, Ahmed Nasr Bekhit, Alshaimaa M Eid, Abeer Awad, Mohammad Aamr, Waleed A Abd El Dayem, Mohamed-Naguib Wifi, Mohamed Alborai

**ORCID number:** Omar Elshaarawy 0000-0002-6945-6204; Sameh Aldesoky Lashen 0000-0002-8599-1338; Nahed A Makhlof 0000-0003-2949-4369; Doaa Abdeltawab 0000-0002-1150-1113; Mariam Salah Zaghloul 0000-0002-4244-5396; Rasha M Ahmed 0000-0002-5912-920X; Hayam Fathy 0000-0002-4218-7550; Shimaaf Afifi 0000-0001-5937-4240; Muhammad Abdel-Gawad 0000-0002-0204-4715; Eman Abdelsameea 0000-0002-3225-7164; Sherief Abd-Elsalam 0000-0003-4366-2218; Salem Youssef Mohamed 0000-0003-2917-4293; Mohammed Tag-Adeen 0000-0001-9813-3191; Mina Tharwat 0000-0001-9834-1644; Ahmed Alzamzamy 0000-0002-3817-5370; Ahmed Nasr Bekhit 0000-0002-9689-3362; Alshaimaa M Eid 0000-0002-4753-9826; Abeer Awad 0000-0001-9945-9767; Mohammad Aamr 0000-0002-1871-4509; Waleed A Abd El Dayem 0000-0002-4897-2170; Mohamed-Naguib Wifi 0000-0002-3403-7106; Mohamed Alborai 0000-0002-8490-9822.

**Author contributions:** Elshaarawy O is guarantor of the article; Elshaarawy O and Alborai M contributed the study conception and design; Elshaarawy O, Lasheen SA and Alborai M provided methodology; Lasheen SA

**Omar Elshaarawy, Eman Abdelsameea,** Department of Hepatology, Gastroenterology and Liver Transplantation, National Liver Institute, Menoufia University, Shebine Elkom 32511, Egypt

**Omar Elshaarawy,** Department of Internal Medicine and Center of Alcohol Research and Liver Diseases, Salem Medical Center, Heidelberg University, Heidelberg 69121, Germany

**Sameh Aldesoky Lashen,** Department of Internal Medicine, Faculty of Medicine, University of Alexandria, Alexandria 21521, Egypt

**Nahed A Makhlof, Doaa Abdeltawab,** Department of Tropical Medicine and Gastroenterology, Faculty of Medicine, Assiut University, Assiut 71515, Egypt

**Mariam Salah Zaghloul,** Department of Hepatology, Gastroenterology and Infectious Diseases, Kafrelsheikh University, Faculty of Medicine, Kafrelsheikh 33565, Egypt

**Rasha M Ahmed, Hayam Fathy,** Department of Internal Medicine, Faculty of Medicine, Assiut University, Assiut 71515, Egypt

**Shimaaf Afifi,** National Hepatology and Tropical Medicine Research Institute, Cairo 11638, Egypt

**Muhammad Abdel-Gawad,** Department of Hepatology, Gastroenterology and Infectious Diseases, Faculty of Medicine, Al-Azhar University, Assiut 71524, Egypt

**Sherief Abd-Elsalam,** Department of Tropical Medicine and Infectious Diseases, Faculty of Medicine, Tanta University, Tanta 31527, Egypt

**Salem Youssef Mohamed,** Department of Internal Medicine, Gastroenterology and Hepatology Unit, Faculty of Medicine, Zagazig University, Zagazig 44519, Egypt

**Mohammed Tag-Adeen,** Department of Internal Medicine, Gastroenterology Unit, Qena Faculty of Medicine, South Valley University, Qena 83523, Egypt

**Mina Tharwat,** Department of Tropical Medicine and Gastroenterology, Faculty of Medicine, Aswan University, Aswan 81528, Egypt

contributed formal analysis and investigation; all authors wrote original draft preparation, reviewed and edited the manuscript; Elshaarawy O and Alborae M contributed supervision; all authors approved the final version of the manuscript.

#### Institutional review board

**statement:** The study was approved by the Institutional Review Board of the National Liver Institute, Menoufia University, Egypt (NLI IRB 00003413) in June 2020, protocol number 00203/2020.

**Informed consent statement:** It is a survey; no consent was needed from patients.

**Conflict-of-interest statement:** Nothing to disclose.

**Data sharing statement:** No additional data are available.

**STROBE statement:** The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Specialty type:** Gastroenterology and hepatology

**Country/Territory of origin:** Egypt

**Peer-review report's scientific quality classification**  
Grade A (Excellent): 0

**Ahmed Alzamzamy,** Department of Gastroenterology and Hepatology, Military Medical Academy, Cairo 11771, Egypt

**Ahmed Nasr Bekhit,** Department of Tropical Medicine, Zagazig General Hospital, Zagazig 42511, Egypt

**Alshaimaa M Eid,** Department of Hepatology and Gastroenterology and Infectious Disease, Faculty of Medicine, AlAzhar University, Cairo 11562, Egypt

**Abeer Awad, Mohamed-Naguib Wifi,** Department of Internal Medicine, Faculty of Medicine, Cairo University, Cairo 11759, Egypt

**Mohammad Aamr,** Department of Hepatology and Gastroenterology, Shebin Elkom Teaching Hospital, Shebine Elkom 3521, Egypt

**Waleed A Abd El Dayem,** Department of Tropical Medicine, Faculty of Medicine, Zagazig University, Zagazig 44519, Egypt

**Mohamed Alborae,** Department of Internal Medicine, Al-Azhar University, Cairo 11651, Egypt

**Corresponding author:** Mohamed Alborae, MD, MRCP, MSc, Lecturer, Department of Internal Medicine, Al-Azhar University, ElMokhayam El Daem St., Nasr City, Cairo 11651, Egypt. [alborae@azhar.edu.eg](mailto:alborae@azhar.edu.eg)

## Abstract

### BACKGROUND

The current coronavirus disease 2019 (COVID-19) pandemic has affected routine endoscopy service across the gastroenterology community. This led to the suspension of service provision for elective cases.

### AIM

To assess the potential barriers for resuming the endoscopy service in Egypt.

### METHODS

A national online survey, four domains, was disseminated over a period of 4 wk in August 2020. The primary outcome of the survey was to determine the impact of the COVID-19 pandemic on the endoscopy service and barriers to the full resumption of a disabled center(s).

### RESULTS

A hundred and thirteen Egyptian endoscopy centers participated in the survey. The waiting list was increased by  $\geq 50\%$  in 44.9% of areas with clusters of COVID-19 cases ( $n = 49$ ) and in 35.5% of areas with sporadic cases ( $n = 62$ ). Thirty nine (34.8%) centers suffered from staff shortage, which was considered a barrier against service resumption by 86.4% of centers in per-protocol analysis. In multivariate analysis, the burden of cases in the unit locality, staff shortage/recovery and the availability of separate designated rooms for COVID-19 cases could markedly affect the resumption of endoscopy practice ( $P = 0.029$ ,  $< 0.001$  and  $0.02$ , respectively) and Odds ratio (0.15, 1.8 and 0.16, respectively).

### CONCLUSION

The COVID-19 pandemic has led to restrictions in endoscopic volumes. The staff shortage/recovery and the availability of COVID-19 designed rooms are the most important barriers against recovery. Increasing working hours and dividing endoscopy staff into teams may help to overcome the current situation.

**Key Words:** COVID-19; Endoscopy; Practice; Pandemic; Egypt; Barriers

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Grade B (Very good): B, B  
 Grade C (Good): 0  
 Grade D (Fair): 0  
 Grade E (Poor): 0

**Received:** October 14, 2020

**Peer-review started:** October 14, 2020

**First decision:** November 3, 2020

**Revised:** November 3, 2020

**Accepted:** November 9, 2020

**Article in press:** November 9, 2020

**Published online:** November 21, 2020

**P-Reviewer:** Galloro G, Liu YC

**S-Editor:** Gao CC

**L-Editor:** A

**P-Editor:** Ma YJ



**Core Tip:** This survey study included 113 endoscopy centers from all over Egypt. Our findings highlighted the barriers for resuming endoscopy in different types of health care facilities in Egypt as well as reporting the current practice in Egyptian endoscopy units during the pandemic. Our study offers real life based snapshot of the current practice and recommendations for routine endoscopic practice during the second wave.

**Citation:** Elshaarawy O, Lashen SA, Makhlof NA, Abdeltawab D, Zaghoul MS, Ahmed RM, Fathy H, Afifi S, Abdel-Gawad M, Abdelsameea E, Abd-Elsalam S, Mohamed SY, Tag-Adeen M, Tharwat M, Alzamzamy A, Bekhit AN, Eid AM, Awad A, Aamr M, Abd El Dayem WA, Wifi MN, Alborai M. Barriers for resuming endoscopy service in the context of COVID-19 pandemic: A multicenter survey from Egypt. *World J Gastroenterol* 2020; 26(43): 6880-6890  
**URL:** <https://www.wjgnet.com/1007-9327/full/v26/i43/6880.htm>  
**DOI:** <https://dx.doi.org/10.3748/wjg.v26.i43.6880>

## INTRODUCTION

In December 2019, a novel an enveloped Ribonucleic acid (RNA) beta coronavirus caused an outbreak called severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) - resulted in coronavirus disease 2019 (COVID-19) began in Wuhan<sup>[1,2]</sup>. The virus rapidly spread throughout the country, then throughout the whole world, and was declared a global pandemic by the World Health Organization in March 2020<sup>[3]</sup>. As of September 23, 2020, it has infected over 31 million people worldwide and caused more than 965000 deaths<sup>[3,4]</sup>.

The clinical gastrointestinal manifestations are present and less severe when compared with SARS<sup>[1,5,6]</sup>. Early reports from Wuhan stated that around 10% of COVID-19 cases had diarrhea and nausea, 1 to 2 d before the beginning of respiratory symptoms<sup>[6]</sup>, and some cases may be missed if screening was applied only to those with respiratory symptoms. It is widely believed that SARS-CoV-2 spreads *via* droplets and contact (especially if within one meter of distance), but there is evidence that airborne spread is also possible during aerosol-generating procedures (AGPs)<sup>[2]</sup>.

In addition, new emerging evidence that SARS-CoV-2 can be found in the feces of patients, giving the possibility of fecal-oral route transmission<sup>[7,8]</sup>. This can be explained by the excessive expression of angiotensin-converting enzyme 2 protein, a receptor for SARS-CoV-2 required for cell entry in the epithelial layer of the gastrointestinal tract<sup>[9]</sup>.

Being highly infective, SARS-CoV-2 put the healthcare workers (HCWs) in clinical departments performing endoscopy in great challenges during this pandemic as being highly susceptible to getting the infection<sup>[10]</sup>. Endoscopy could be considered a high-risk procedure as pulmonary and gastric secretions, as well as fecal material, may contain high viral loads. So, infection to HCWs occurs either due to direct contact with the infected patients or to the lack or improper use of the personal protective equipment (PPE)<sup>[11]</sup>. In fact, several studies have discussed infection prevention and control measures that must be implemented to increase patient safety, prevent nosocomial outbreaks, protect HCWs, and ensure the rational use of limited PPE<sup>[11,12]</sup>.

Several endoscopy societies and expert groups have offered recommendations and position statements for endoscopy during the COVID-19 pandemic<sup>[11,13-16]</sup>. Several GI society guidelines recommended ceasing elective procedures during the pandemic to decrease the risk of infection<sup>[17-19]</sup>. Different recommendations/guidelines have been recently released to ensure the smooth reopening of endoscopy units and resuming elective procedures<sup>[17]</sup>. Whether or not endoscopy units in different parts of the world are ready to resume service based on these recommendations is not fully known.

This study is a multicenter study to discuss the barriers of resuming endoscopic maneuvers in different centers and governorates in Egypt as the coronavirus infection rate has remarkably decreased and there are international calls for resuming all the activities with caution and different precautions, and to assess whether the endoscopic service has returned to its near normal standard level.

## MATERIALS AND METHODS

### Methods

We designed an online survey that was based on four domains (Supplementary Figure 1). The first domain included the demographic data as regards the governorate, the type of health care facility, and the type(s) of endoscopy service(s) provided. The second domain was for the status of the COVID-19 pandemic and its impact on the endoscopy service(s). The third domain was directed to the readiness of the health care facility for dealing with COVID-19 cases (infrastructure, working staff, PPEs, case stratification, *etc.*), and the fourth domain was about the ability of the facility to resume the endoscopy service(s) and the different barriers which preclude service resumption. The whole survey included 20 main questions. All the questions were closed-ended questions that could be answered by selecting the appropriate answer(s).

The questionnaire was set-up by a nationally representative group of endoscopy physicians covering different Egyptian governorates. The survey was designed using the RedCap platform (v9.1.0., United States). The study was approved by the Institutional Review Board of the National Liver Institute, Menoufia University, Egypt (NLI IRB 00003413) in June 2020, protocol number 00203/2020. The questionnaire was distributed online for 4 wk (starting from July 30, 2020 till August 28, 2020). The primary outcome of the survey was to identify the current impact of the COVID-19 pandemic on the endoscopy service and the barriers against the full resumption of the participating centers.

### Statistical analysis

Data were collected and entered into personal computers and analyzed using Statistical Package for Social Sciences (SPSS version 26.0) software (IBM SPSS Inc. Chicago, United States). The incomplete responses were excluded from the analysis. For simplification, we divided the responding governorates according to the administrative division of the governorates of Egypt (see below in discussion). The data were expressed as numbers (proportions). Comparisons between groups were made by the  $\chi^2$  or Fisher's test (FET) for the qualitative variables. A uni- and multivariate regression analysis was done to identify the parameters which determine the decision making as regards the resumption of endoscopy service if not working. Correlation analysis also was done to explore the correlation between the current status of endoscopy service and the different domains.

## RESULTS

### Characteristics of the endoscopy Centers

The survey was sent to 115 endoscopy centers all over Egypt, of which 113 (98.3%) responded to the survey in complete response and two centers didn't give a full response, so they were excluded from the analysis. Most respondents ( $n = 67$ , 59.3%) were from high-volume endoscopy units either University or teaching facility general hospitals providing emergency endoscopy. Cairo and Giza regions have participated with 30 (26.5%) centers, Lower Egypt (Alexandria, Beheira, Sharqia, Qalyubia, Damietta, and Ismailia) and Upper Egypt (Assiut, Fayoum, Sohag, Qena, Aswan and Luxor) have participated through 34 (30.1%) centers, and the Delta region (Menoufia, Kafr El-sheik, Gharbia, and Dakahlia) have participated through 13 (11.5%) centers. COVID-19 designated facilities were present among 48 (42.5%) centers (Figure 1).


### Patient selection and classification

On the day of the planned endoscopic procedures patients were checked for being suspected cases of COVID-19 in 95 (84.1%) of centers. The process of selection and screening was based on symptoms among 90 centers (79.6%), temperature check among 82 centers (72.6%), non-contrast multi spiral computed tomography (MSCT) chest among 50 centers (44.2%), polymerase chain reaction (PCR) testing among 11 centers (9.7%), and antibody testing among 3 centers (2.7%).

### Impact of the current status of the pandemic on the waiting lists

Endoscopy practice was almost resuming or resumed in about 70% of the centers (Figure 2A). In general, there was an expansion of the waiting lists all over the country. The Quantification of the expansion is shown in Figure 2B. We found significant variation in the waiting lists according to the distribution of COVID-19 pandemic






**Figure 1** Distribution of different types of healthcare facilities and endoscopic services according to different regions. A: Distribution of health care facilities; B: Availability of different endoscopic services. ERCP: Endoscopic retrograde cholangiopancreatography; EUS: Endoscopic ultrasound; VCE: Video capsule endoscopy (Y-axis is standing for percentage).

status among different centers (Figure 2B and C). In areas with clusters of cases ( $n = 49$ ), the waiting lists increased by 0%-25% in 14.29%, 26%-50% in 32.65%, 51%-75% in 34.69%, 76%-100% in 10.2%, and by more than 100% in 8.17% of these centers. While in areas with sporadic cases ( $n = 62$ ), 45.16% of the centers had their waiting list increased by 0%-25%, however, in 19.35% and 20.9% of the centers the list increased by 26%-50%, and 51%-75%, respectively. In about 14.59% of the centers, the waiting list increased by more than 75%, meanwhile, areas with no new cases ( $n = 2$ ), 50% of centers had their list increased by only 0%-25% (FET = 19.1,  $P = 0.005$ ).

### Shortage in staff

During the COVID-19 outbreak, endoscopy staff has been deployed to COVID-19 designated wards and hospitals. In addition, a significant number of healthcare providers were infected. The shortage of endoscopy staff was present among 39 (34.8%) of centers (Figure 2D). Specialists shortage was present in 88 (77.9%) of centers, nursing staff shortage in 90 (79.6%) of centers, and trainees shortage in 110 (97.3%) of centers. This shortage was considered as a barrier against resuming endoscopy service among 38 (33.6% in intention to treat analysis, 86.4% in per-protocol analysis; 69 centers didn't answer). The shortage in specialists was more pronounced in Delta and lower Egypt than upper Egypt and Cairo and Giza (61.5% and 91.3% vs. 64.7% and 83.3% respectively,  $P = 0.019$ ). However, the shortage of nursing staff and trainees were not significantly different among different centers,  $P = 0.93$  and  $P = 0.24$ , respectively).



**Figure 2** Impact of the status of the pandemic in different regions on the increase of waiting lists as well as resumption of endoscopic services and the impact of staff shortage. A: Current status of endoscopy services; B: Percentages of increased waiting list; C: Status of coronavirus disease 2019 epidemic; D: Staff shortage. COVID-19: Coronavirus disease 2019 (X-axis is standing for percentage).

Only 41 units (35%) had to increase the working hours to compensate for this shortage. Seventy-six units (67.3%) recovered their staff either from sick leave or COVID-19 wards.

**Pre-procedural precautions applied by different Centers during the COVID-19 pandemic**

During the current COVID-19 epidemic in Egypt, 72 (63.7%) of endoscopy centers have provided suitable waiting areas for appropriate social distancing precautions, 65 (57.5%) of centers could provide an adequate number of recovery rooms to keep the social distancing strategy to the optimum, and 40 (35.4%) of centers have increased their working hours to accommodate the extra-burden of increased cases volume; at the same time, 77 (68.1%) of centers have adopted selection strategy to select the endoscopist and the assisting team based on the presence of a suspected/confirmed COVID-19 cases. Sixty-three (55.7%) units were planning to resume the pre-pandemic capacity with a designated track for confirmed COVID-19 cases. However, only 24 (21.2%) of centers could provide special endoscopy rooms for confirmed or suspected COVID-19 cases. Of note, training was already resumed in 16 units (14.2%).

The Intra-procedural precautions applied during the COVID-19 pandemic in Egypt varied according to the availability of PPE in each facility locality as shown in Table 1 (more details in Supplementary Table 1).

**Table 1** Intra-procedural precautions applied by different Centers during the coronavirus disease 2019 pandemic in Egypt, *n* (%)

Governorates	Usage of PPE				Action taken by centers	
	For all cases	For suspected cases	For confirmed cases only	Vary accordingly	Delay of turnover for cleaning	Increased time/case
Cairo and Giza	5 (16.7)	13 (43.3)	12 (40)	8 (26.7)	27 (90)	21 (70)
Upper Egypt	10 (29.4)	7 (20.6)	13 (38.2)	7 (20.6)	32 (94.1)	27 (79.4)
Lower Egypt	13 (36.1)	10 (27.8)	10 (27.8)	8 (22.2)	32 (88.9)	29 (80.6)
Delta	7 (53.8)	4 (30.8)	3 (23.1)	3 (23.1)	13 (100)	13 (100)

PPE: Personal protective equipment.

### Post-procedural precautions applied by different Centers during the COVID-19 pandemic

Post-procedural patient follow-up for the development of fever or suspected COVID-19 symptoms for 14 d after discharge was a strategy adopted by 18 (15.9%) of centers only. The follow up was performed by either a resident doctors (8%), endoscopy nurses (5.3%), administrative staff (6.2%), and/or specialist physicians (1.8%).

Multivariate regression analysis showed that burden of cases in the unit locality, staff shortage and their recovery and the availability of separate designated rooms for COVID-19 cases could markedly affect the resumption of endoscopy practice ( $P = 0.029$ ,  $< 0.001$  and  $0.02$ , respectively) and Odd's ratio (0.15, 1.8 and 0.16, respectively) (Table 2).

## DISCUSSION

The prolonged suspension of routine endoscopic services during the SARS-CoV-2 pandemic has significant implications on diagnostic endoscopic services such as delay in diagnosis and management of cancer patients as well as the expansion of waiting lists worldwide. Therefore, this encouraged the international gastrointestinal endoscopic societies to release position statements, recommendations, and guidance for the rapid and safe resumption of endoscopic services commensurate with facilities and pandemic situations of each country<sup>[14,17-19]</sup>. To our knowledge, whether or not endoscopy centers in Egypt are ready to resume elective services has not been studied. Accordingly, the results of the current survey represent the current status of resuming routine endoscopic services in Egypt and determine the barriers of resuming such services.

Our results showed that 70% of different centers all over Egypt have resumed routine endoscopic services as is illustrated in Figure 2A. This is in parallel to the current situation of early recovery phases of the pandemic and the decline in the number of new COVID-19 cases in Egypt and other countries worldwide<sup>[3]</sup>. Expansion of elective endoscopy waiting lists is one of the challenges to resume the full capacity of endoscopic services. According to the British Society of Gastroenterology guidance in the early recovery stages, triage mechanisms are needed to prioritize patients scheduling according to the indications<sup>[20]</sup>.

The highest percentages of waiting lists expansion are observed in areas with clusters of COVID-19 cases which indicate the awareness of senior decision-makers of endoscopy centers in Egypt with the importance of triaging and prioritization of patients scheduling in the light of clinical need with the available capacity. Also, this reflects their compliance with GI endoscopy societies' guidance. In addition, this observation complies with the current SARS-CoV-2 pandemic situation in Egypt, as there is a decrease in the number of areas with new cases allowing endoscopy centers to restore their full capacity as soon as possible.

Regarding the shortage in endoscopy staff, 34.5% of the centers reported a significant shortage of staff during the SARS-CoV-2 pandemic in nursing staff (79.6%) and specialists (77.9%). This is related to two factors; one was the reallocation of staff and medical equipment to the surge in demand to manage suspected and confirmed COVID-19 cases and the other was some of them get infected caused by frequent exposure to infected patients. The shortage in specialists was more pronounced in

Table 2 Regression analysis for factors that determine decision making as regard resuming endoscopy service

Factor	Univariate		Multivariate		
	P value	OR	P value	OR	95%CI
Status of pandemic	0.03	1.5	0.029	0.15	0.20-0.9
Shortage of PPE	0.04	4.2	0.42	2.24	0.31-16.1
Staff recovery	0.05	1.4	< 0.001	1.8	5.1-6.7
Having a COVID designed unit	0.04	4.2	0.01	0.5	0.11-0.82
Having separate endoscopy room	0.04	1.01	0.02	0.16	0.03-0.7
Patient classified before procedure	0.03	1.7	0.007	0.7	1.1-1.18

OR: Odd's ratio; COVID: Coronavirus disease; PPE: Personal protective equipment.

Delta and Lower Egypt geographically which may be related to the geographic distribution of a large number of confirmed COVID-19 cases and isolation hospitals that need more medical staff.

This shortage in endoscopy staff is considered as a barrier against resuming endoscopy service among 38 centers in the current study (33.6% in intention to treat analysis where  $n = 113$ , 86.4% in the per-protocol analysis where  $n = 44$ ). Regarding patient selection before endoscopy, a large number of participating centers (84.1%) screened their patients. The most common method used was symptoms-based screening for COVID-19 risk stratification (79.6%) of the centers and fewer centers used PCR testing (9.7%).

Similarly, Alborae *et al*<sup>[21]</sup> reported that most worldwide centers (93.9%) screened patients for possible COVID-19 disease prior to the procedure and 54 centers (33.13%) used PCR testing. However, asymptomatic infected patients are a known source for transmission<sup>[2,22]</sup>, accordingly, this type of screening is not sufficient to guide for COVID-19 risk stratification. A recent study from China, employed both symptoms-based screening and PCR in patient's screening, reported no cases of endoscopy-related nosocomial COVID-19 disease transmission in 1361 cases<sup>[23]</sup>.

Regarding pre-procedural precautions and endoscopy centers designation, our results show that most participating centers so far follow the different international GI endoscopy societies guidelines<sup>[14,17-19]</sup> including appropriate social distancing precautions or increase working hours to accommodate the extra-burden of increased cases volume. More than two-thirds of our centers have adopted a selection strategy to select the endoscopist and the assisting team based on the presence of a suspected/confirmed COVID-19 case. Also, Alborae *et al*<sup>[21]</sup> stated that the majority (78.5%) of the centers considered dedicated teams for the delivery of endoscopic services after the onset of the pandemic.

To reduce the risk of infection spread most of the international society's recommendations are in agreement with the existence of a restricted protocol for infection control and the awareness of medical staff for the donning and doffing of PPE. According to the results of our study, there is wide variability between different centers regarding intra-procedural precautions depending upon the availability of PPE and type of procedures. We can overcome those types of barriers by reuse some of PPE components such as respirator masks in case of shortage as it was reported by many studies as the methods of decontamination were explained<sup>[23-25]</sup>.

Most of the international recommendations on endoscopy encourage follow-up of patients after endoscopic procedures<sup>[11]</sup>. Unfortunately, in our study few participating centers (15.9%) are practicing the strategy of following their patients for the development of any suspected symptoms post-procedure. A similar study showed that 18.4% of centers only called patients back two weeks after procedures<sup>[21]</sup>. According to multivariate regression analysis that was done in our study, we found that endoscopy staff shortage, availability of separate designated rooms for COVID-19 cases, and the burden of cases in each locality are the most obvious barriers for the resumption of routine endoscopy practice in Egypt.

Although there are many barriers that may interfere with the complete restoration of endoscopy services in Egypt, according to the results of our study, most participating centers are preparing their facilities for the resumption of full endoscopy services.

We recommend increasing working hours and dividing endoscopy staff into teams

to overcome the shortage of endoscopy staff. Also, follow up of patients for two weeks after endoscopic procedures to detect any possible transmission of SARS-CoV-2 infection in endoscopy centers as this strategy may help in tracing the source of nosocomial transmission.

---

## CONCLUSION

---

In conclusion, the current study represents an important national multicenter survey addressing crucial information about resuming regular gastrointestinal endoscopy services and its barriers in Egypt.

## ARTICLE HIGHLIGHTS

### **Research background**

An outbreak of coronavirus disease 2019 (COVID-19) has hit the world and disturbed the whole healthcare system, including endoscopic practices which are a very risky procedures in terms of exposure to infection.

### **Research motivation**

Smooth resumption of routine endoscopic service has to be guaranteed as well as decreasing the burden of exposing patients and endoscopy staff to infection during endoscopic procedures. Many strategies have to be implemented in endoscopy units, however, these strategies face many barriers as shortage of personal protective equipment, working staff and post procedure tracing of infection and follow-up of patients.

### **Research objectives**

This survey study was designed to evaluate the feasibility and the difficulty of resumption of routine endoscopic service in the context of COVID-19 pandemic.

### **Research methods**

We conducted a survey study that included 20 questions to be answered by the head of endoscopy units in 113 units from all over Egypt.

### **Research results**

One hundred and thirteen centers participated in the study from all over Egypt. Due to halting the routine endoscopic services during the pandemic, the waiting lists were doubled in most of the centers. Third of the centers experienced significant shortage of endoscopy staff. The lack of dedicated endoscopy rooms for infected patients and the staff shortage were the main barriers to resume routine services smoothly.

### **Research conclusions**

We recommend increasing working hours and dividing endoscopy staff into teams to overcome the shortage of endoscopy staff. Also, follow up of patients for two weeks after endoscopic procedures to detect any possible transmission of infection in endoscopy centers as this strategy may help in tracing the source of nosocomial transmission.

### **Research perspectives**

For smooth resumption of endoscopic service between two waves of the current pandemic, several strategies have to be applied in a uniform manner in all endoscopy units. Our study showed how different centers acted differently during this pandemic in terms of their plan to safely getting routine endoscopy service on track. Future studies should propose modalities to guarantee uniform application of determined strategies that overcome the current barriers.

---

## REFERENCES

---

- 1 Xie J, Tong Z, Guan X, Du B, Qiu H. Clinical Characteristics of Patients Who Died of Coronavirus Disease 2019 in China. *JAMA Netw Open* 2020; 3: e205619 [PMID: 32275319 DOI: ]

- 10.1001/jamanetworkopen.2020.5619]
- 2 **Huang C**, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**: 497-506 [PMID: 31986264 DOI: 10.1016/S0140-6736(20)30183-5]
  - 3 **World Health Organization**. Coronavirus Disease (COVID-19) Situation Reports. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>
  - 4 **Joao BN**. Geographic Information Systems and COVID-19: The Johns Hopkins University Dashboard. 2020 Preprint. Available from: ResearchSquare: rs-15447 [DOI: 10.21203/rs.3.rs-15447/v1]
  - 5 **Chen N**, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J, Yu T, Zhang X, Zhang L. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020; **395**: 507-513 [PMID: 32007143 DOI: 10.1016/S0140-6736(20)30211-7]
  - 6 **Wang D**, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y, Zhao Y, Li Y, Wang X, Peng Z. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* 2020; **323**: 1061-1069 [PMID: 32031570 DOI: 10.1001/jama.2020.1585]
  - 7 **Gu J**, Han B, Wang J. COVID-19: Gastrointestinal Manifestations and Potential Fecal-Oral Transmission. *Gastroenterology* 2020; **158**: 1518-1519 [PMID: 32142785 DOI: 10.1053/j.gastro.2020.02.054]
  - 8 **Xiao F**, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for Gastrointestinal Infection of SARS-CoV-2. *Gastroenterology* 2020; **158**: 1831-1833. e3 [PMID: 32142773 DOI: 10.1053/j.gastro.2020.02.055]
  - 9 **Wang W**, Xu Y, Gao R, Lu R, Han K, Wu G, Tan W. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA* 2020; **323**: 1843-1844 [PMID: 32159775 DOI: 10.1001/jama.2020.3786]
  - 10 **Ni W**, Yang X, Yang D, Bao J, Li R, Xiao Y, Hou C, Wang H, Liu J, Yang D, Xu Y, Cao Z, Gao Z. Role of angiotensin-converting enzyme 2 (ACE2) in COVID-19. *Crit Care* 2020; **24**: 422 [PMID: 32660650 DOI: 10.1186/s13054-020-03120-0]
  - 11 **Soetikno R**, Teoh AYB, Kaltenbach T, Lau JYW, Asokkumar R, Cabral-Prodigalidad P, Shergill A. Considerations in performing endoscopy during the COVID-19 pandemic. *Gastrointest Endosc* 2020; **92**: 176-183 [PMID: 32229131 DOI: 10.1016/j.gie.2020.03.3758]
  - 12 **Mayol J**, Fernández Pérez C. Elective surgery after the pandemic: waves beyond the horizon. *Br J Surg* 2020; **107**: 1091-1093 [PMID: 32383479 DOI: 10.1002/bjs.11688]
  - 13 **Zhang Y**, Zhang X, Liu L, Wang H, Zhao Q. Suggestions for infection prevention and control in digestive endoscopy during current 2019-nCoV pneumonia outbreak in Wuhan, Hubei province, China. *Endoscopy* 2020; **52**: 312-314 [PMID: 32212122 DOI: 10.1055/a-1128-4313]
  - 14 **Catlow J**, Beaton D, Beintaris I, Shaw T, Broughton R, Healey C, Penman I, Coleman M, Rutter M. JAG/BSG national survey of UK endoscopy services: impact of the COVID-19 pandemic and early restoration of endoscopy services. *Frontline Gastroenterol* 2020; **0**: 1-7 [DOI: 10.1136/flgastro-2020-101582]
  - 15 **Repici A**, Maselli R, Colombo M, Gabbiadini R, Spadaccini M, Anderloni A, Carrara S, Fugazza A, Di Leo M, Galtieri PA, Pellegatta G, Ferrara EC, Azzolini E, Lagioia M. Coronavirus (COVID-19) outbreak: what the department of endoscopy should know. *Gastrointest Endosc* 2020; **92**: 192-197 [PMID: 32179106 DOI: 10.1016/j.gie.2020.03.019]
  - 16 **Francis N**, Dort J, Cho E, Feldman L, Keller D, Lim R, Mikami D, Phillips E, Spaniolas K, Tsuda S, Wasco K, Arulampalam T, Sheraz M, Morales S, Pietrabissa A, Asbun H, Pryor A. SAGES and EAES recommendations for minimally invasive surgery during COVID-19 pandemic. *Surg Endosc* 2020; **34**: 2327-2331 [PMID: 32323016 DOI: 10.1007/s00464-020-07565-w]
  - 17 . American Society for Gastrointestinal Endoscopy: guidance for trainees during the COVID-19 pandemic. *Gastrointest Endosc* 2020; **92**: 748-753 [PMID: 32485214 DOI: 10.1016/j.gie.2020.05.032]
  - 18 **Chiu PWY**, Ng SC, Inoue H, Reddy DN, Ling Hu E, Cho JY, Ho LK, Hewett DG, Chiu HM, Rerknimitr R, Wang HP, Ho SH, Seo DW, Goh KL, Tajiri H, Kitano S, Chan FKL. Practice of endoscopy during COVID-19 pandemic: position statements of the Asian Pacific Society for Digestive Endoscopy (APSDE-COVID statements). *Gut* 2020; **69**: 991-996 [PMID: 32241897 DOI: 10.1136/gutjnl-2020-321185]
  - 19 **Gralnek IM**, Hassan C, Beilenhoff U, Antonelli G, Ebigbo A, Pellissè M, Arvanitakis M, Bhandari P, Bisschops R, Van Hooft JE, Kaminski MF, Triantafyllou K, Webster G, Pohl H, Dunkley I, Fehrke B, Gazic M, Gjergjek T, Maasen S, Waagenes W, de Pater M, Ponchon T, Siersema PD, Messmann H, Dinis-Ribeiro M. ESGE and ESGENA Position Statement on gastrointestinal endoscopy and the COVID-19 pandemic. *Endoscopy* 2020; **52**: 483-490 [PMID: 32303090 DOI: 10.1055/a-1155-6229]
  - 20 **The British Society of Gastroenterology**. BSG Guidance on recommencing GI Endoscopy in the deceleration & early recovery phases of the COVID-19 pandemic. Available from: <https://www.bsg.org.uk/covid-19-advice/bsg-guidance-on-recommencing-gi-endoscopy-in-the-deceleration-early-recovery-phases-of-the-covid-19-pandemic/>
  - 21 **Alboraie M**, Piscoya A, Tran QT, Mendelsohn RB, Butt AS, Lenz L, Alavinejad P, Emara MH, Samlani Z, Altonbary A, Monged A, Lemmers A, Sudovykh I, Ho DQD, Ghazanfar S, Kamau E, Iqbal S, Tan DMY, Liao WC, Vignesh S; "The Global Endo-COVID working group". The global impact of COVID-19 on gastrointestinal endoscopy units: An international survey of endoscopists. *Arab J Gastroenterol* 2020; **21**: 156-161 [PMID: 32912748 DOI: 10.1016/j.ajg.2020.08.008]
  - 22 **Hoehl S**, Rabenau H, Berger A, Kortenbusch M, Cinatl J, Bojkova D, Behrens P, Böddinghaus B, Götsch U, Naujoks F, Neumann P, Schork J, Tiarks-Jungk P, Walczok A, Eickmann M, Vehreschild MJGT, Kann G, Wolf T, Gottschalk R, Ciesek S. Evidence of SARS-CoV-2 Infection in Returning Travelers from Wuhan, China. *N Engl J Med* 2020; **382**: 1278-1280 [PMID: 32069388 DOI: 10.1056/NEJMc2001899]
  - 23 **Lindsley WG**, Martin SB Jr, Thewlis RE, Sarkisian K, Nwoko JO, Mead KR, Noti JD. Effects of Ultraviolet Germicidal Irradiation (UVGI) on N95 Respirator Filtration Performance and Structural Integrity. *J Occup Environ Hyg* 2015; **12**: 509-517 [PMID: 25806411 DOI: 10.1080/15459624.2015.1018518]

- 24 **Mills D**, Harnish DA, Lawrence C, Sandoval-Powers M, Heimbuch BK. Ultraviolet germicidal irradiation of influenza-contaminated N95 filtering facepiece respirators. *Am J Infect Control* 2018; **46**: e49-e55 [PMID: 29678452 DOI: 10.1016/j.ajic.2018.02.018]
- 25 **Holmdahl T**, Walder M, Uzcátegui N, Odenholt I, Lanbeck P, Medstrand P, Widell A. Hydrogen Peroxide Vapor Decontamination in a Patient Room Using Feline Calicivirus and Murine Norovirus as Surrogate Markers for Human Norovirus. *Infect Control Hosp Epidemiol* 2016; **37**: 561-566 [PMID: 26861195 DOI: 10.1017/ice.2016.15]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

