



# Diagnosis and Molecular Identification of Virulent Infectious Bursal Disease in Naturally Infected Broiler Chickens.

Ghazi A. M. Zohair<sup>1\*</sup>, M. M. Amer<sup>2</sup>, A. EL-shemy<sup>3</sup>, M.A. Bosila<sup>4</sup>, Kh. M. Elbayoumi<sup>4</sup>

<sup>1</sup>Faculty of Agriculture, Department of animal production, Sana`a University, Republic of Yemen

<sup>2</sup>Dept of Poultry, Dis., Faculty Vet. Med., Cairo University, P.O. Code 12211 Giza, Egypt

<sup>3</sup>Department of Parasitology, Vet. Res. Division, NRC, P.O. Code 12311 Dokki, Giza, Egypt

<sup>4</sup>Dept of Poultry. Dis., Vet. Res. Division, NRC, P.O. Code 12311 Dokki, Giza, Egypt

## ABSTRACT

Infectious bursal disease (IBD) is a serious problem in young chickens and losses occurs in face of restricted biosecurity, frequently vaccination of breeder to maintain uniform high maternal antibody titers and trials to apply more effective vaccines in broiler. The disease was diagnosed based on clinical signs, pathology, and AGPT as well as virus isolation and detection of viral antigen in tissue by RT-PCR. Therefore, bursae were collected from 5 IBD suspected flocks aged 4 to 5 weeks having clinical signs, mortality (35-70%) and lesions suggestive to infected with virulent IBDV. AGP test results showed positive precipitation lines in 5/5 tested flocks (100%) between the prepared bursal homogenate or CAM of inoculated ECE with high embryo mortality of virulent IBDV. The 5 bursal homogenate extract were subjected to RT/PCR products of VP2 full gene and electrophoretic pattern of amplified samples revealed the presence of specific PCR products (642 bp). BY restriction end nuclease enzymes BstNI and SspI the samples are negative at 210 bp, 171 bp, 151 bp and 110 bp bands suggested either to be very virulent IBDV (vvIBDV), while Sspi restriction enzyme analyzed on 2 % agarose gel electrophoresis showed that Results clarify the isolates to be very virulent subtype. The recorded histopathological changes in the examined bursal sections were similar to those previously reported due to infection with virulent and vvIBDV.

Conclusions: We can concluded that virulent IBDV still circulating and cause losses in vaccinated broiler and RT-PCR is useful, rapid and accurate in diagnosis IBDV outbreaks.. It is important to look for a more suitable vaccine and vaccination programs for both breeder and broiler chicken to face and controlling very virulent IBDV mutation.

**Key Words:** Broiler chicken, IBDV, isolation and identification AGP-test, RT-PCR, histopathology

eIJPPR 2017; 7(5):29-34

**HOW TO CITE THIS ARTICLE:** Ghazi A. M. Zohair, M. M. Amer, A. EL-shemy, M.A. Bosila, Kh. M. Elbayoumi. (2017). "Diagnosis And Molecular Identification of Virulent Infectious Bursal Disease in Naturally Infected Broiler Chickens." , *International Journal of Pharmaceutical and Phytopharmacological Research*, 7(5), pp:29-34.

## INTRODUCTION

The name of Infectious bursal disease (IBD) was designated by Hitchner [1] for an acute contagious disease of young chicks reported for the first time 1957 in the

Gumboro area of Southern Delmarva, USA by Cosgrove [2]. The disease was characterized by ruffled feather, watery diarrhea, trembling and severe prostration. Morbidity was 10-25 % and mortality averaging 5% where the characteristic enlargement of

**Corresponding author:** Ghazi A. M. Zohair

**Address:** Faculty of Agr., Dept of animal production, Sana`a University, Republic of Yemen

**e-mail** ✉ gzohai@yahoo.com

**Relevant conflicts of interest/financial disclosures:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Received:** 05 January 2016; **Revised:** 19 June 2017; **Accepted:** 28 July 2017



infected bursa was described. IBDV belongs to the Avibirnavirus genus. A major antigenic difference between IBDV strains of serotype-1 isolated in different poultry producing areas in USA [3,4,5] as well as major antigenic shift of serotype-I IBDV strains [6,7]. Mortalities of up to 60 % due to acute IBD outbreaks of highly pathogenic virus in broilers [8,9,10], while El-Batrawi [11] indicated that the majority of the highly virulent pathotype, producing acute disease with severe clinical picture and high mortalities up to 70%. The vvIBDV strain was differed from standard strain (CU-1) in the viral protein 2. It was reported that AGP test is useful for detection of IBDV antigen in bursa [12, 13]. RT/PCR-RE assay is a rapid method for differentiation and identification for detection of IBD viral RNA in bursa of infected chicken with unknown field isolates [14, 15].

RT-PCR was used to identify the point mutation for amplification of 743 bp fragment of VP2 gene followed by RE analysis using BstNI, StyI and MboI that cut at bp 759, 852 and 1063 respectively. BstNI enzyme cut at 759 bp, StyI enzyme cut at 852 bp and MboI enzyme that cut at 1063 bp [16]. IBDV isolates can be differentiated into several groups using restriction enzymes (SSPI, BstNI, MboI, StyI) as most of the classic strains were BstNI and StyI double positive while most of the variant strains were BstNI and StyI double negative [14,17,18]. The reverse transcriptase / polymerase chain reaction using restriction enzyme analysis assay (RT-PCR-RE) could be used to diagnose IBDV in chickens [19].

The pathological changes induced by a highly virulent IBDV causing sever clinical signs and high mortality was recorded [20, 21,22].

This study was carried out for diagnosis of IBDV infection in natural affected 3-4weeks old broiler chickens flock with sudden high mortality based on clinical as well as isolation and molecular detection of virus with histopathological examination of collected bursae.

## MATERIAL AND METHODS

### Field history:

Fife flocks with stock number of 6,000- 40,000 broiler chickens from different broiler breeds derived from vaccinated breeder flocks. these flocks were

Primer	Primer Design	Position
VP2 upstream	5` GCGATGACAAACCTGCAAGAT 3`	93-114 bp of CU-1 strain
VP2 downstream	5` AGGTGGGAACATGTGGAGAC 3`	1470-1490 bp of CU-1 strain
HVR upstream	5` TCACCGTCCTCAGCTTAC 3`	587-604 bp of STC strain
HVR downstream	5` TCAGGATTTGGGATCAGC 3`	1212-1229 bp of STC strain

### Histopathological examination:

The fixed bursae halves with the 10% neutral formol saline, dehydrated in grade alcohol, cleared with xylene and embedded in paraffin. Sections of an average thickness of 5 micros were stained with

vaccinated with intermediate IBD vaccine at 8-12 days of age via drinking water and feed on commercial pelted ration. Chicken Flocks were aged 4 to 5 weeks showing clinical signs, high mortalities (35- 72% within 3-4 days) and lesions suggestive to naturally infected with IBDV [23].

### Samples:

Ten Bursae per flock were aseptically collected from sacrificed infected diseased chickens with recording of signs and postmortem lesions. Each bursa was divided into 2 pieces one half was homogenized and the homogenates were subjected to agar gel precipitation (AGP) test, viral isolation, passage in ECE and RT-PCR, while the 2<sup>nd</sup> half was fixed in 10% neutral formol saline for histopathological examination for diagnosis of this outbreak.

### Embryonated chicken eggs (ECE):

Specific pathogen free (SPF) ECE were obtained from Koum Osheim, El-Fayoum and used for virus isolation from suspected field bursal sample, virus passage and preparation of positive antigen through inoculation on the chorioallantoic membrane (CAM) at 10-11 days of age [24,25].

### Agar gel precipitation test (AGPT):

It was carried out according to Wood et al. [26] the test was done to demonstrate the presence of IBDV antigen against positive immune serum and positive antigen prepared from CAM homogenate of inoculated ECE with E228 vaccine [24]. Also naturally infected bursa was prepared to be antigen [24, 27].

### Preparation of bursal samples for virus isolation:

The half of collected field bursae were prepared for ECE inoculation according to Hitchner [1,28].

### Conventional RT-PCR:

RT-PCR was carried out according to Cardoso et al. [29]. RNA extraction from bursa and CAM of 2<sup>nd</sup> passage in ECE was extracted according to Cardoso et al. [30]. Primers synthesized by Metabion Company, Germany as described according to the published sequence of CU-1 and STC strains by Bayliss et al. [31] in the following tables:

hematoxyline and eosin according to Sharma et al. [32] and Bancroft et al. [33].

## RESULTS AND DISCUSSION

IBD is a serious problem in young chickens and losses occurs in face of restricted biosecurity, frequently vaccination of breeder to maintain uniform high maternal antibody titers and trials to apply more effective vaccines in broiler [23]. The disease was diagnosed based on clinical signs, pathology, and AGPT. The field cases showed the following signs, lower feed intake, ruffled feather, watery diarrhea and sudden increased morbidity (range 35- 72% within 3-4 days) out of flock number 6,000-40,000 chickens from different broiler breeds derived from vaccinated breeder flocks. Similar signs and mortifies were reported [9,10,11,20,22,34,35]. Appearance of very virulent IBDV 1987 which cause mortalities up to 100 % in vaccinated birds had been reported to increase the economic importance of such disease [8, 11, 36, 37]. This result can be supported by virulent strains of IBDV of same serotype have been reported to overcome high MDAs in commercial flocks vaccinated with vaccines developed from different variants, causing up to 60% to 70% mortality [38]. Died and sacrificed chickens showed post mortem lesions including patches of hemorrhage in breast and thigh muscles (Figure 1 a), hemorrhagic atrophied bursa (Figure 1 b), enlarged edematous bursa with hemorrhages (Figure 1 c), full hemorrhagic bursa (Figure 1 d), nephritis, petichial hemorrhages in proventriculus mucosa [20,23]

hemorrhages (Figure 1c), full hemorrhagic bursa (Figure 1d), nephritis, petichial hemorrhages in proventriculus mucosa [20,23]



**Figure 1.** Lesion in naturally infected broiler chickens.

1 a: Patches of hemorrhage in breast and thigh muscles. 1 b: hemorrhagic atrophied bursa. 1c: enlarged edematous bursa with hemorrhages. 1d: full hemorrhagic bursa.

AGP test results showed positive precipitation lines in 5/5 tested flocks (100%) between the prepared bursal homogenate or CAM of inoculated eggs used for viral passage and the positive serum [24, 26, 27].

**Table 1.** Results of AGP positive bursa (n=10) and ECE in passages of isolate IBDV(n= 5).

Sample No	Bursal samples		Passage 1		Passage 2	
	Positive	%	Positive ECE	%	Positive ECE	%
1	4/10	40	3/5	60	5/5	100
2	8/10	80	4/5	80	5/5	100
3	5/10	50	3/5	60	4/5	80
4	5/10	50	2/5	40	3/5	60
5	7/10	70	3/5	60	5/5	100

Infected embryos with AGP positive homogenates in 5 flocks showed increased both embryo lesions and mortalities with passage as hemorrhages on head and toes with sever congested liver, edematous body, greenish liver, homogenates on skin feather tract, dwarfing embryo (Table 1) [39,40]. Similar embryo mortality was reported due to veriolent IBDV [41].


The bursal homogenate and CAM of 2<sup>nd</sup> passage in ECE extract were subjected to RT/PCR products of VP2 full gene was used using internal specific primers covering the hyper variable region within the VP2 gene for confirmation of the VP2 full gene RT/PCR product and for amplification of the hyper variable region for genotyping of the IBDV local isolates. The electrophoretic pattern of amplified samples revealed the presence of specific PCR products (642 bp) at the correct expected size of the VP2 gene [29,35,42].

RFLP was carried out using 2 different restriction of endonuclease enzymes BstNI and SspI for genotyping of the IBDV isolated strains. 2 µg DNA of each sample was digested with BstNI restriction enzyme. The resulted fragments were analyzed by 2% agarose gel

electrophoresis. The samples are negative at 210 bp, 171 bp, 151 bp and 110 bp bands suggested either to be vvIBDV [16, 36].

Samples were purified from low melting agarose gel and digested with Sspi restriction enzyme analyzed on 2 % agarose gel electrophoresis. Results clarify the isolates to be very virulent subtype as the enzyme digested gave 2 bands at 390 bp and 253 bp indicating very virulent subtypes [14, 17, 18, 22].

The recorded histopathological changes in the examined bursal sections of sacrificed field cases showed vacullation of the lymphoid follicles with mononuclear leukocytes infiltration (Figure 2a). Atrophy of bursal lymphoid follicles with interfollicular edema (Figure 2 b), necrosis of medullar lymphocytes interfollicular with congestion (Figure 2 c), complete follicular necrosis with cyst like space containing eosinophilic with homogenous mass and cell debris (Figure 2 d) . The recorded bursal cells changes were similar to those previously reported due to infection with veriolant and very veriolant IBDV [22,32,43,44,45].



**Figure 2.** Bursal stained sections of naturally infected chicken with IBDV. (H&E X 100).  
2a). Vacuolation of the lymphoid follicles with mononuclear leukocytes infiltration.  
2b). Atrophy of bursal lymphoid follicles with interfollicular edema.  
2c). Necrosis of medullary lymphocytes interfollicular with congestion.  
2d). Complete follicular necrosis with cyst like space containing eosinophilic.

Clinical signs, gross lesions, histopathological findings, ECE inoculation and results of AGP test were characteristic of virulent IBDV. This result was also confirmed by results of RT-PCR.

### CONCLUSION

It can be concluded that virulent IBDV still circulating and cause losses in vaccinated broiler and RT-PCR is useful, rapid and accurate in diagnosis IBDV outbreaks.. it is important to look for a more suitable vaccine and vaccination programs for both breeder and broiler chicken to face and controlling very virulent IBDV mutation.

### REFERENCES

- [1] Hitchner S B. 1970. Infectivity of infectious bursal disease virus for embryonating eggs. *Poult. Sci.*, 49: 511-516.
- [2] Cosgrove A S. 1962. An apparently new disease of chickens-avian nephrosis. *Avian Dis.* 6 385-389.
- [3] Saif, Y M. 1984. Infectious bursal disease virus types. *Proc. 19<sup>th</sup> Natl. Mett. Poult. Hlth. Condemn.*, P. 105.
- [4] Rosenberger J K , Cloud S S. 1985. Isolation and characterization of variant infectious bursal disease virus. *J. Am. Vet. Med. Assoc.*, 198: 357.
- [5] MacNulty M S , Saif Y M. 1988. Antigenic relationship of non-serotype I turkey infectious bursal disease virus. *Avian Dis.*, 32: 374-375.
- [6] Snyder D B, Marquardt W W, Mallinson E T, Russek-Cohen E, Savage P K , Allan D C. 1986. Rapid serological profiling of ELISA, association of infectious bursal disease virus serology with broiler flock performance. *Avian Dis.* 30 :139-148.
- [7] Van Den Merel R, Snyder D B, Luettiken D. 1990. Antigenic characterization of infectious bursal disease virus field isolates by their reactivity with a panel of monoclonal antibodies. *Dtsch. Tieraztl. Wschr.*, 97 :81-83.
- [8] Van Den Berg T. P., Morales D. , Meulemans G. 1997. Use of Baculovirus expressed VP2 subunit Protein for diagnosis and control of infectious bursal disease. *Proc. XIth. Int. Cong. Wld. Vet. Asso, Budapest, Hungary.* PP. 57.
- [9] Vieltz E., Vob D. M., Conard C. 1991. Is protection against Gumboro disease Problem? *Lohmann Information* 16 : 6-13.
- [10] Gaudry D. 1993. Experiences with very virulent IBDV. *Proceedings of the 130<sup>th</sup> annual meeting of the American Veterinary Medical Association , Minneapolis, Minnesota.* P. 156 (Abstract).
- [11] El-Batrawi A. M. 1990. Studies on severe out breaks of infectious bursal disease. 1-the natural and experimental disease. *Proceed. Of 2<sup>nd</sup> scientific conf. egypt. Vet. Poult.* . 339-252.
- [12] Cullen G.A. , Wyeth P.J. 1975. Quantitation of antibodies to infectious bursal disease. *Vet. Rec.*, 97 : 315.

- [13] Kadam M, Jhala M K. 2003. Comparison of serological tests for detection of infectious bursal disease virus in bursal samples. *Indian Vet J.* 80: 723-6.
- [14] Cao Y. C, Yeung W. S, Law M, BI Y Z, Leung F.C., Lim B. L. 1998. Molecular characterization of seven Chinese isolates of infectious bursal disease virus: classical, very virulent, and variant strains. *Avian Dis.* 42: 340-351.
- [15] Wu C. C, Lin T. L, Zhang H. G, Davis V S, Boyle J. A. 1992. Molecular detection of infectious bursal disease virus by polymerase chain reaction. *Avian Dis.*, 36: 221-226.
- [16] Jackwood D. J., Jackwood R. J. , Sommer S. E. Identification and comparison of point mutations associated in classic and variant infectious bursal disease viruses. *Virus Res.* 49 (1997) 131-137.
- [17] Ture O., Saif Y. M., Jackwood D. J. 1998. Restriction fragment length polymorphism analysis of highly virulent strains of infectious bursal disease virus from Holland, Turkey and Taiwan. *Avian Dis.*, 42: 470- 479.
- [18] Amer M.M. , Nassif S. A. 2005. Studies on recent virus field variant isolates: Genomic identification and differentiation using Rt-PCR-RFLP. In proc. of 4 th. Sci. Conf. *Facult Vet. Med. Bane-suef.*, 24-27 September 134-138.
- [19] Khaliel S. A., El-Lakany H. F. Molecular detection of infectious bursal disease virus in commercially reared chickens using the reverse transcriptase\polymerase chain reaction-restriction endonuclease assay *Monuha Vet. J.* vol 1 (2000) 1: 239-296
- [20] Nunoya T., Otaki Y., Tajima M., Hiraga M., Saito Y. 1992. Occurrence of acute infectious bursal disease with high mortality in Japan and pathogenicity of field isolates in SPF chickens. *Avian Dis.* 36: 597-609.
- [21] Bayoumi A. , Mousa S. , Ibrahim M.K. , Deep S. 1984. Microscopical studies of smears from tissue and blood of birds affected with infectious bursal disease. *Assuit Vet. Med.* 11 22: 83-85.
- [22] Amer M. M., EL-Bayomi K. M., Kutkat M. A., Wafaa A. Abd EL-Ghany, Shakal M. A. , Sherein S. Abd EL-Gaied. 2007. Isolation, molecular characterization and pathogenicity studies of infectious bursal disease field virus isolates. In procc. of the 5<sup>th</sup> sci.Conf., *Facult. Vet. Med., Beni-Suef Uni.*, 6-9 November 41-51.
- [23] Saif Y.M , Fadly, A. M., Glisson J. R McDougald L. R.. Nolan L. K. and Swayne, D. E. 2008. *Infectious bursal disease in Poultry Diseases*, 12<sup>th</sup> Ed., Iowa State Press, Iowa, 195 -208.
- [24] Hirai K., Kawamoto E. , Shimakura S. 1974. Some properties of precipitating antigens associated with infectious bursal disease virus. *infection and immunity*, 10 (6)1235-1240.
- [25] Gelb, J. Eidson, C. S, Kleven, S. H. 1979. Interferon production in embryonating chickens eggs following inoculation with infectious bursal disease virus. *Avian Dis.* 23 (2) 534-538.
- [26] Wood BW, Muskett JC, Hebert CN, Thornton DH. 1979. Standardization of the quantitative agar gel precipitation test for antibodies to infections bursal disease. *Biol. Stand.* 7:89-96.
- [27] Hirai, K., Shimakura S., Hirose. M.1972. Immunodiffusion reaction to avian infectious bursal disease virus. *Avian Dis.* 16: 961-964.
- [28] Anon.1990. Laboratory manual for the isolation and identification of avian pathogens. 3<sup>rd</sup> edition , A. A. A. P. Arnold printing corporation , Itheca, New York.
- [29] Cardoso T C. , Rosa A.C.G, Astolphi R.D., Vincente R.M., Novais J. B. , Hirata K.Y. and Luvizotto M. C.R. 2008. Direct detection of infectious bursal disease virus from clinical samples by in situ reverse transcriptase-linked polymerase chain reaction. *Avian Pathol* 37 (4) 457- 461.
- [30] Cardoso T.C., Rahal P., Pilz D., Teixeira M.C.B, Arns C.W. 2000. Replication of classical infectious bursal disease virus in the chicken embryo related cell line. *Avian Pathol.* 29: 213-217.
- [31] Bayliss C. D., U. Spies K. Shaw R. W., Peters A., Papageorgiou H. Muller, Boussnell M. 1990. A comparison of the sequences of segment A of four infectious bursal disease virus strains and identification of a variable region in VP2. *J Gen Virol* 71 : 1303-1312.
- [32] Sharma J. M., Dohms J. E, Metz A. L.1989. Comparative pathogenesis of serotype-1 and variant serotype-1 isolates of infectious bursal disease virus and their effect on humeral and cellular immune competence of SPF chickens. *Avian Dis.*, 33 : 112-124.
- [33] Bancroft JD, Stevens A, Turner DR. 1996. *Theory and practice of histological techniques.* 4<sup>th</sup> Ed. New York, London: Churchill Livingston.
- [34] Sultan H. A.1994. Studies on infectious bursal disease in chickens. PH. D. Thesis, Alexandria University.
- [35] El Bayomi Kh M.2007. Studies on infectious bursal disease. Dept. of Poul. Diseases, *Facult. Vet. Med., Cairo University.*
- [36] Chettle N. J., Smart J. C., Wyeth P. J. 1989. Outbreak of virulent infectious bursal disease in east Anglia. *Vet. Rec.* 125: 271-272.
- [37] Eterradosi N., Picault J. P., Druin P., Michele Guittlet, Rolande L'hospitalier, Bennejean G. 1992. Pathogenicity and



- preliminary antigenic characterization of 6 infectious bursal disease virus strains isolated in France from acute outbreaks. *J. Vet. Med. B.*, 39: 683-691.
- [38] Etterradossi N. 2001. Major advances in infectious bursal disease virus (IBV) research. Proc of 2nd Int Symp on Infectious Bursal Disease and Chicken Infectious Anaemia, Rauschholzhausen; July 2001; Ebsdorfergrund, Germany p. 6-23.
- [39] Lukert P. D., Saif Y. M. 1991. infectious bursal disease in: Disease of boultry, Ninth ed., A. A. A. P., Iowa state university press. Ames, Iowa, USA. P. 649-663.
- [40] Macferran J. B., MacNulty M. S., McKillop E. R., Conner Y. J., McCracken R. M., Collins D. S., Allan G. M. 1980. Isolation and serological studies with infectious bursal disease virus from fowl, turkey and duck: demonstration of a second serotype. *Avian Pathol.*, 9 (3) 395-404.
- [41] Lasher H.N., Shane S.M. 1994. Infectious bursal disease. *World's Poul. Sci. J.* 133-166.
- [42] Liu H. J., Giambone J. J., Dormitorio T. 1994. Detection of genetic variations in serotype I isolates of infectious bursal disease virus using PCR and restriction endonuclease analysis. *J. Virol. Methods.* 48: 281-291.
- [43] Jhala M.K., Kher H.N., Prajapati K.S. 1990. Experimental infection of infectious bursal disease virus in chicken. *Ind. J. Anim. Sci.*, 60 (11) 1309-1310.
- [44] Dash. Bb, Verma KC, Panisup AS., Kataria JM. 1991. Pathogenicity of a field isolate of infectious bursal disease virus in chicken. *Indian J of Vet Pathol.* 15 (1) 21-25.
- [45] Inoue M., Fukuda M., Miyano K. 1994. Thymic lesions in chickens infected with infectious bursal disease virus. *Avian Dis.*, 38 (4) 839-846.