



Isolation and Identification of Genotype VII of Newcastle Disease Virus From Chicken Flocks in Six Egyptian Governorates

Hager-Magdy^{1*}, M. M. Amer², Kh. M. Elbayoumi¹, M. A. Kutkat¹

¹Dept of Poultry. Diseases, Vet. Res. Division, National Research centre, P.O. Code 12311 Dokki, Giza, Egypt

²Dept of Poult. Dis., Faculty Vet. Med., Cairo University, P.O. Code 12211 Giza, Egypt

*Corresponding Author: dr_gogo200810@yahoo.com

Abstract: Samples were collected from 89 chicken flocks suffering from respiratory signs and/or nervous signs accompanied with high mortalities, in Qalubia, Sharkia, Menofia, Fayoum, Behira and Cairo governorates at the period from August 2014 to May 2015. These samples, were trachea, liver, spleen and cecal tonsils. Virus isolation and molecular identification were occurred from these collected samples. Results revealed that, positive percentage to Genotype VII of Newcastle disease virus (NDV) in Qalubia, Sharkia, Menofia, Fayoum, Behira and Cairo governorates were 31.42%; 21.74%; 37.5% ;25%; 25% and 40%; respectively. For prevention and control of such virus, it is recommended for further protective trial using different vaccination regimes for choosing the best protocol in order to control such pathotype under our Egyptian field conditions which needs further investigations.

Key words: Isolation, identification, Genotype VII, Newcastle disease virus, chicken

INTRODUCTION

Newcastle disease virus (NDV) was considered as one of the most important poultry disease in the past till now affecting both wild and domestic poultry species (Spradbrow *et al.*, 1990; seal *et al.*, 2000), causing severe economic losses (Aldous *et al.*, 2003). The disease caused by Newcastle disease virus belonged to Paramyxoviridae family, Genus Avulavirus responsible for significant outbreaks in poultry industries with varies severity due to its different pathotypes based on their virulence in chickens which are: lentogenic, mesogenic and velogenic (Alexander, 2003; OIE, 2004; Alexander *et al.*, 2012). Nowadays, velogenic strain of NDV becomes endemic in many countries (Giambrone and Closser, 1990). Clinical disease in poultry characterized by respiratory signs including coughing, gasping, sneezing and rales, swelling around neck and eye, circling and neck twisting and sharp decrease in egg production in layers (Wakamatsu *et al.*, 2006), the disease considered zoonotic as it transmitted from infected birds to contact human causing mild conjunctivitis, influenza-like symptoms and in severe cases, it can lead to some lasting impairment of vision (Beard *et al.*, 1984). Control of ND fulfilled by strict biosecurity together with proper vaccination program which is varied from country to another (Miller and Koch, 2013). Despite of use of live attenuated NDV vaccine together with inactivated oil-emulsified vaccination for control of Newcastle disease outbreaks, these strategies were not able to completely prevent outbreaks against circulating virulent virus which may be due to emerging of highly virulent pathotype which exhibit antigenic variations that could be responsible for recent outbreaks of NDV (Wang *et al.*, 2015). In Egypt, NDV outbreaks are still frequently occurring in vaccinated poultry flocks, despite the intensive vaccination programs (Abdel-Moneim *et al.*, 2006; Mohamed

et al., 2009 and 2011; Radwan et al., 2013; Hussein, et al., 2014; Nabila et al., 2014; Awad et al., 2015; El-Bagoury et al., 2015).

From the above mentioned data, our trail is designated in order to isolate and identify the field virulent NDV that circulating in vaccinated Egyptian farms causing different outbreaks causing severe economic losses with detection of its Genotype as well as its relation with strains registered in GenBank.

Materials and Methods

Chicken flocks

Eighty-nine chicken flocks aged from 17 days to 385 days with history of previous vaccination against ND reared in 6 Egyptian governorate (Qalubia, Sharkia, Menofia, Fayoum, Cairo and Behira) in the period from August 2014 to May 2015 suffering from respiratory signs and/or nervous signs accompanied with high mortalities varied from 20– 80 % with post mortem lesions suggestive to be naturally infected with ND (Fib 1-3).

Fig:(1) Proventriculus showing peticheal and echymotic haemorrhages on the tips of glands.



Fig:(2) Enlarged spleen with focal splenic necrosis



Fig (3): Caecal tonsils showing severe necrosis with haemorrhagic ulcer



Samples:

Samples including trachea, liver, spleen and cecal tonsils were collected separately from each bird. Collected samples were labeled and transported immediately in ice tank to the laboratory for processing. Samples tissue of each flock were pooled and ground in sterile phosphate buffer saline pH 7.0–7.4 containing gentamycin (50 lg/ml) and mycostatin (1000 units/ml) in a 1:5 (w/v) dilution, centrifuged and tissue supernatant was collected and stored at -20 C till being used in virus isolation by using RT-PCR.

Virus isolation:

Method were preformed according to *OIE* (2012) and summarized as follow:

Processed samples were inoculated into allantoic sac at 9-day-old specific pathogen free (SPF) embryonated chicken eggs (ECE) then incubated at 37 C for 4 days. Candling was taken place daily for embryo viability. Allantoic fluid was harvested dead embryonated eggs after chilling eggs at 4 C overnight.

Haemagglutination (HA) test: (GRIMES, 2002):

Allantoic fluids were tested for haemagglutination (HA) using 1 % chicken RBCs. The HA-negative samples were passaged two further times in SPF eggs, and the HA titer of the allantoic fluid was again determined. HA positive allantoic fluids were assayed for NDV using RT-PCR.

Reverse Transcription-Polymerase Chain Reaction (RT-PCR) Extraction of viral RNA:

The genomic viral RNA was extracted from harvested HA positive allantoic fluid by using QIAamp viral RNA extraction Kits according to the manufacture's protocol.

Conventional RT-PCR for detection of NDV F-Protein gene (Jestin & Jestin, 1991):

One step RT- PCR was carried using QIAGEN® OneStep RT-PCR kit (QIAGEN, Valencia, CA) according to the manufacturer's instructions. RT-PCR was used for the detection of partial F-gene of vNDV using the following primers:

forward 5' -ATGGGCYCCAGACYCTTCTAC-3' and

Reverse 5' -CTGCCACTGCTAGTTGTGATAATC-3'

that flanks a 535 bp of the F gene of NDV as previously described (Seal, 1995). Thermal cycling RT-PCR conditions included a reverse transcription 50°C for 30 min. Then an initial PCR activation step 95°C for 15 min. followed by 39 cycles at 94°C for 30 sec., 55°C for 30 sec. and 72°C for 45 sec. then the final extension was performed at 72°C for 10 min.

Agarose gel electrophoresis:

The PCR products were separated in 1.5% agarose gel in TAE buffer stained with ethidium bromide and compared with molecular mass marker (100 bp DNA markers) and visualized by ultraviolet (UV) transillumination.

Purification of PCR Products from the gel and gene sequencing:

For gene sequencing, the target bands of specific size were excised from the gel and purified with the QIAquick gel extraction kit (Qiagen, Valencia, CA) according to the manufacturer instructions and The purified DNA was sequenced in an automated ABI 3730 DNA sequence (Applied Biosystems, USA).

The obtained sequences were aligned by the Clustal W method using MEGA V5.05 software. The nucleotide sequences were compared with NDV sequences available in GenBank. A phylogenetic tree of aligned sequences was constructed by Boost-trap method. The deduced amino acid sequences were determined to detect the pathotype of isolated NDV.

Results

Isolation and identification of ND virus:


Samples were collected from 89 of chicken flocks, as shown in table (1). These flocks were different in types; ages and localities, previously vaccinated against ND by different types of live and in activated vaccines.

Table (1): Result of Slide HA test after 1st passage and PCR for isolates.

Province	NO. of farms	SPF ECE Passage		Slide HA test after 1 st passage		PCR Results	
		1 st passage	2 nd passage	+ve	-ve	+ve	-ve
Qalubia	35	35	11	11	24	11	24
Sharkia	23	23	5	5	18	5	18
Menofia	8	8	3	3	5	3	5
Behira	8	8	2	2	6	2	6
Fayoum	8	8	2	2	6	2	6
Cairo	5	5	2	2	6	2	6

RT-PCR NDV isolates were detected by RT-PCR using degenerate primers for the fusion protein gene result in amplicon 535 bp as shown in fig (4).

Fig (4): Agarose Gel Electrophoresis (AGE) pictures showing 535 bp RT-PCR Products of NDV Fusion protein gene.




Phylogeny and genetic analysis of the NDV isolates strains:

The amino acid sequence of fusion protein cleavage site of the isolates carries motif ¹¹²RRQKRF¹¹⁷ that is consistent with viruses of velogenic strains.

The phylogenetic analysis of partial sequences of the selected strains F gene showed that the isolated viruses belong to genotype VII.

Fig (5): Phylogenetic tree based on a partial sequence of NDV F gene. showing the relationship between the selected Egyptian NDV isolates in the present study with vaccinal strain and reference NDV strains from gene bank. Black dots refer to viruses isolated in current study.



Discussion

Newcastle disease virus considered the most predominant avian viral diseases affecting poultry industry in Egypt causing high economical losses, although intensive vaccination programmes carried out in field (Sabra, 2013 and El Behairy *et al.*, 2016). This may be due to genetic diversity among NDV strains in last years (Miller *et al.*, 2010) resulting in field outbreaks or emerging of new pathotype resulting in severe infection (Radwan *et al.*, 2013). So, the isolation and pathotyping of NDV from outbreaks among chickens is a critical point for the control of NDV and vaccination evaluation (Awad *et al.*, 2015).

The recorded signs of natural infection in field cases under investigation was previously reported by (Abdel-Moneim *et al.*, 2006; Nabila *et al.*, 2014 and Awad *et al.*, 2015). The calculated mortality rates of natural outbreaks ranged 20-90% in vaccinated flocks (Siddique *et al.*, 2013; Abdel-Gelil *et al.*, 2014 and Abd El Aziz *et al.*, 2016). Lesions in figs 1-3 were also recorded by (Abdel-Moneim *et al.*, 2006; Nabila *et al.*, 2014 and Awad *et al.*, 2015).

Molecular identification by RT-PCR revealed that; our isolate is Newcastle field virus; this result was obtained from positive sample from clinical cases after primary isolation in specific pathogen free embryonated chicken egg (SPF-ECE). Many researchers used RT-PCR for isolation of Newcastle disease as reliable method for detection of positive cases (Wang *et al.*, 2001; Creelan *et al.*, 2002 and Liu *et al.*, 2011) which considered accurate method for detection of NDV.

For relatedness detection of isolated NDV field isolate sequencing is carried out, results revealed that our isolated strain belonged to class II genotype VIIId which considered velogenic strain causing severe outbreaks in china (Liu *et al.*, 2003) and middle east (Khan *et al.*, 2010 and Rui *et al.*, 2010).

Conclusion:

Results concluded that, NDV isolates circulating among chickens are virulent (Genotype VIId) and associated with outbreaks in poultry farms and is responsible for severe economic losses. For prevention control of such virus, it is recommended for further protective trial using different vaccination regimes for choosing the best protocol in order to control such pathotype under our Egyptian field conditions which needs further investigations.

References

- Abdel-Moneim, A. S.; El-Sawah, A. A.; and Kandil, M. A. (2006): Characterization of variant strain of Newcastle disease virus in Egypt. *BS. Vet. Med. J.*, 16: 12-17.
- Aldous EW., Mynn JK., Banks J and Alexander DJ. (2003): A molecular epidemiological study of avian paramyxovirus type 1 (Newcastle disease virus) isolates by phylogenetic analysis of a partial nucleotide sequence of the fusion protein gene. *Avian Pathol.*; 32:239-256.
- Alexander, D.J. (2003): Newcastle disease. In Y. M. Saif, Barnes, H.J., Lisson, J.R., Fadly, A.M., McDougald, L.R., Swayne, D.E. (ed). *Disease of Poultry*, pp 64-87. Iowa State University Press.
- Alexander DJ., Aldous EW and Fuller CM. The long view (2012): a selective review of 40 years of Newcastle disease research. *Avian Pathol.*; 41(4):329335.
- Awad, A. M.; Sedeik, M. E. and Abdel kariem, A. A. (2015): Isolation, Molecular Characterization and Pathotyping of Newcastle Disease Viruses from Field Outbreaks Among Broiler Flocks in Egypt from 2014-2015. *Internat. J. of Current Res.* 7(2): 12925-12934. ISSN: 0975-833X.
- Beard, C.W. and Hanson, R.P. (1984): Newcastle Disease. In Hofsad, M.S., Barnes, H.J., Clanek, B.W., Reid, W.W., Yoder, H.W. (ed). *Disease of Poultry*, 8th ed. pp452-470 Iowa State University Ames.
- El-Bagoury, G.F.; Samar F. El-Adaway, El-Habbaa, A.S. and Suzan S. El-Mahdy (2015): Isolation, identification and pathotyping of Newcastle disease virus from chickens in Egypt. *Benha Vet. Med. J.*, 29 (1) 196-204.
- Elbehairy, M.A., Bastami, M.A. and Manal, A. Afify. (2016): Some Epidemiological studies on Newcastle Disease in domesticated chicken and Turkey Flocks in different Egyptian Governorates on 2015. M.Sc. Thesis, Cairo University, Egypt.
- Giambrone JJ. and Closser J. (1990): Effect of breeder vaccination on immunization of progeny against Newcastle disease. *Avian Dis.* ;34(1):114– 119.
- Grimes, S. E. (2002): A basic laboratory manual for the small-scale production and testing of I-2 Newcastle disease vaccine. RAP publication, 136.
- Hussein, H.A., Emara, M.M. and Rohaim, M.A. (2014): Molecular characterization of Newcastle Disease Virus Genotype VIID in Avian influenza H5 N1 Infected Broiler Flock in Egypt. *Internat. J. of Virol.* 10(1): 46 – 54.
- Jestin, V. and Jestin, A. (1991): Detection of Newcastle disease virus RNA in infected allantoic fluids by in vitro enzymatic amplification (PCR). *Archives of Virol.*,118:151-161.
- Jing-Yu Wang., Wan-Hua Liu., Juan-Juan Ren., Pan Tang., Ning Wu., Hung-Yi Wu., Ching-Dong Chin and Hung-Jen Liu (2015): Characterization of emerging Newcastle disease virus isolates in China. *Virol J.* 12: 119.
- Khan S.T.; Rehmani, Rue. C.; Miller, P. and Afonso, C.L. (2010): Phylogenetic and pathological characterization of Newcastle disease virus isolates from Pakistan. *J. Clin. Microbiol.*, 48:1892–1894.
- Liu, H.; Zhao, Y.; Zheng, D.; Lv, Y.; Zhang, W.; Xu, T.;Li, J. and Wang, Z. (2011): Multiplex RT-PCR for rapid detection and differentiation of class I and class II Newcastle disease viruses, *J. of Virological Methods*, 171: 149-155 *Pathol.*, 33(5): 482-491.
- Liu, X. F.; Wan, H. Q.; Ni, X. X.; Wu, Y. T. and Liu, W. B. (2003): Pathotypical and genotypical characterization of strains of Newcastle disease virus isolated from outbreaks in chicken and goose flocks in some regions of China during 1985-2001. *Archives of Virol.*, 148: 1387–1403.
- Miller, P.J., Decanini, E.L and Afonso, C.L(2010). Newcastle disease evolution of genotypes and the related diagnostic challenges. *Infect. Genet Evol.*, 10: 26–35.

Mohamed, M.H.A.; Kumar, S.; Paldurai, A.; Megahed, M.M.; Ghanem, I.A.; Lebdah, M.A. and Samal, S.K. (2009): Complete genome sequence of a virulent Newcastle disease virus isolated from an outbreak in chickens in Egypt. *Virus Genes* (2009) 39:234–237. DOI 10.1007/s11262-009-0385-7.

Mohamed N. Abd El Aziz, Hatem S. Abd El-Hamid, Hany F. Ellkany, Soad A. Nasef, Sherif M. Nasr and Ahmed R. El Bestawy. (2016): Biological and Molecular Characterization of Newcastle Disease Virus Circulating in Chicken Flocks, Egypt, During 2014-2015. *Zagazig. Vet. J.* 44(1): p. 9-20

Mostafa Y. Abdel-Glil, Sunil K. Mor, Tamer A. Sharafeldin, Robert E. Porter, and Sagar M.Goyal. (2014): Detection and Characterization of Newcastle Disease Virus in Formalin-Fixed, Paraffin-Embedded Tissues from Commercial Broilers in Egypt. *Avian Dis.*, 58(1):118-123. 2014.

Nabila Osman, Sultan, S., Ahmed, I.A., Ibrahim, R.S. and Sabra, M. (2014): Isolation and Pathotyping of Newcastle Disease Viruses from Field Outbreaks among Chickens in the Southern Part of Egypt 2011-2012. *Global Veterinaria* 12 (2): 237-243.

Naila Siddique, Khalid Naeem, Muhammad Athar Abbas, Akbar Ali Malik, Farooq Rashid, Saba Rafique, Abdul Ghafar and Abdul Rehman. (2013): Sequence and phylogenetic analysis of virulent Newcastle disease virus isolates from Pakistan during 2009–2013 reveals circulation of new sub genotype. *Virol.* 444(2013)37–40.

Office des Epizooties [OIE International] (2004): Manual of Standards for diagnostic test and vaccines draft test for comment by experts in member countries. OIE Manual.

Patti J. Miller and Koch G. (2013): Newcastle disease. In: Swayne DE, Glisson JR, McDougald LR, Nolan LK, Suarez DLO, Nair VN, editors. *Diseases of poultry*. 13th ed. Athens: John Wiley & Sons; p.89-107.

Radwan, M.M., Darwish, S.F., El-Sabagh, I.M., El-Sanousi, A.A and Shalaby, M.A., (2013). Isolation and molecular characterization of Newcastle disease virus genotypes II and VIIId in Egypt between 2011 and 2012. *Virus Genes* 47, 311–316.

Rui,Z. ; Juan, P.; Jingliang, S.; Jixun, Z.; Xiaoting, W.; Shouping, Z.; Xiaojiao, L. and Guozhong, Z., (2010): Phylogenetic characterization of Newcastle disease virus isolated in the mainland of China during 2001-2009. *Vet. Microbiol.* 141, 246-257.

Sabra, M. H. (2013): molecular characterization of Newcastle disease virus isolated from chickens in south valley provinces. M.Sc. Thesis, South Valley University, Qena, Egypt.

Seal B.S., King D.J. and Sellers H.S. (2000): *Dev Comp Immunol*, 24, 257.

Seal, B. S. (1995): Analysis of matrix protein gene nucleotide sequence diversity among Newcastle disease virus isolates demonstrates that recent disease outbreaks are caused by viruses of psittacine origin. *Virus Genes* 11:217-224.

Spradbrow, P.B. (1990). *Village Poultry and Preventive Veterinary Medicine*. Preventive Vet. Med., 8: 305-307.

Wakamatsu, N., King, D.J., Kapozynski, D.R., Seal, B.S., Brown, C.C. (2006). Experimental Pathogenesis for Chickens, Turkeys and Pigeons of exotic Newcastle disease from an outbreak in California during 2002-2003. *Vet. Pathol.*,43: 926-933 .

Wang, Z.; Vreede, F. T.; Mitchell, J. O.; Viljoen, G. J.; (2001): Rapid detection and differentiation of Newcastle disease virus isolates by a triple one-step RT-PCR. *Onderstepoort J. Vet. Res.* 68, 131--134.