



Post-extraction volumetric analysis of alveolar ridge contour using subepithelial connective tissue graft in esthetic zone: a randomized controlled clinical trial

Nourhan Gamal¹ · Nesma Shemais¹ · Marwa Al-Nawawy¹ · Noha A. Ghallab¹

Received: 28 May 2023 / Accepted: 9 September 2023
© The Author(s) 2023

Abstract

Objectives The aim of this randomized clinical trial was to assess the alveolar ridge contour after soft tissue augmentation using subepithelial connective tissue graft (SCTG) buccal to fresh extraction sockets in patients with thin buccal bone, versus minimally-traumatic extraction followed by spontaneous healing solely.

Materials and methods Forty non-restorable maxillary teeth in the esthetic zone were randomly assigned into two groups: minimally-traumatic extraction with SCTG (test) and minimally-traumatic extraction followed by spontaneous healing (control). The outcomes assessed included linear volumetric change of buccal soft tissue contour, vertical tissue loss, gingival thickness (GT), and interdental papilla (IDP) height after 6 months.

Results The SCTG group showed a significant improvement ($P < 0.001$) in all outcomes after 6 months. The SCTG group showed a statistically significant ($P < 0.001$) gain in the buccal soft tissue volumetric change compared to the control group. The SCTG group showed a statistically significant increase in GT ($P < 0.001$) and IDP height ($P < 0.05$) after 6 months compared to the control group.

Conclusions The use of SCTG buccal to extraction sockets in the anterior maxilla might be considered as a predictable approach for preserving the alveolar ridge contour.

Clinical relevance SCTG buccal to extraction sockets might counteract post-extraction hard and soft tissue alterations in the esthetic zone.

Keywords Connective tissue graft · SCTG · Alveolar ridge contour · Pouch technique · Post-extraction dimensional changes · Linear volumetric change

Introduction

Following tooth extraction, several sequential events arise causing significant qualitative and quantitative alterations at the edentulous site. Socket healing process results in ridge dimensional changes of the underlying bone, as well as the overlying soft tissue architecture [1]. Loss of soft tissue contour following tooth extraction could persist after healing, regardless of the utilization of alveolar ridge preservation techniques. As a result, clinicians seek different surgical techniques to restore the post-extraction hard and


soft tissue volume loss [2]. Since esthetics is the main concern for most patients, soft tissue grafting is becoming a routine in the daily clinical practice [2, 3], to compensate for the diminished supra-crestal tissue dimension that usually occurs following tooth loss and implant placement [4]. Such procedures have a role in increasing tissue thickness, re-establishing an adequate width of keratinized tissue, correcting mucogingival deformities, and improving esthetics around teeth and dental implants [5–7]. Moreover, the presence of a sufficient quantity and quality of soft tissues play a chief role in the long-term maintenance of peri-implant health [8]. However, recent retrospective study reported no difference in volumetric, linear changes, and peri-implant conditions between implant sites with or without soft tissue grafting over a period of 12 years [9].

The concept of performing soft tissue augmentation on the buccal side of the extraction socket following tooth

✉ Nourhan Gamal
Nourhan.moustafa@dentistry.cu.edu.eg

¹ Oral Medicine and Periodontology, Faculty of Dentistry, Cairo University, Giza, Egypt

Fig. 1 CONSORT flowchart



extraction was adopted by many experts and clinicians, in order to stabilize the soft tissues and compensate for the buccal concavity that arises after tooth loss [10]. Meanwhile, there is an extensive body of evidence proving that sub-epithelial connective tissue grafts (SCTGs) are considered the gold standard in ridge contour augmentation procedures [11–14]. Previous review articles and systematic reviews concluded that various therapeutic approaches are built on SCTG-based procedures and that SCTG is superior in soft tissue correction and augmentation surgeries [13, 15, 16]. This is in agreement with the conclusions of the latest consensus report of group 2 of the SEPA/DGI/OF workshop [17]. The authors stated that superior esthetic outcomes were observed in the presence of a thick mucosa; the connective tissue graft remains the standard protocol of care in terms of increasing mucosal thickness.

Based on the current available literature, clinical research related to dimensional alterations post tooth extraction mainly focused on the hard tissue modeling, while the impact of soft tissue healing in post-extraction sites received a little consideration. The investigation of post-extraction soft tissue volumetric changes in future clinical trials was strongly recommended in the latest consensus report on the management of the extraction sockets and timing of implant placement [18]. Given the existing gap of knowledge, this randomized clinical trial aimed to volumetrically assess the alveolar ridge contour after soft tissue augmentation using

SCTG buccal to fresh extraction sockets in patients with thin buccal bone, versus minimally-traumatic extraction followed by spontaneous healing with no soft tissue augmentation procedure.

Materials and methods

Study population

The present randomized clinical trial was registered in the ClinicalTrials.gov (ID NCT04482127), approved by the Research Ethics Committee, Faculty of Dentistry, Cairo University (13–12-19), conducted in accordance with the Helsinki Declaration of 1975, as revised in 2013 and reported according to the CONSORT guidelines [19] presented in Fig. 1. This study included 20 non-restorable maxillary teeth indicated for extraction in the esthetic zone in 32 patients selected from the outpatient clinic, Department of Periodontology, Faculty of Dentistry, Cairo University, between January 2020 and March 2022. The sites included 5 central incisors, 4 lateral incisors, 3 canines, and 8 premolars in SCTG group and 3 central incisors, 4 lateral incisors, 3 canines, and 10 premolars in minimally-traumatic extraction followed by spontaneous healing group. All participants met the following inclusion criteria: having non-restorable maxillary teeth indicated for extraction from 2nd premolar

to 2nd premolar region, intact gingival tissue with at least 2-mm keratinized tissue, buccal bone thickness 1 mm or less assessed by CBCT¹ preoperatively, and periodontally and systemically healthy patients. Exclusion criteria included pregnant and lactating females, current and former smokers, and presence of active infection with soft tissue communication. All patients provided written informed consent to participate in this trial. Initial patient examination was performed including full-mouth supragingival scaling and 0.12% chlorhexidine HCL² mouthwash twice daily was prescribed for 2 weeks with patient motivation and oral hygiene instructions.

Randomization and blinding

Sequence generation was executed using simple randomization by a computer-generated list from numbers 1:40 by an investigator (NM) who was not involved in recruitment nor treatment. Allocation concealment was implemented by the same investigator using sequentially numbered, opaque, sealed envelopes handled to the surgeon (GN) who did not open them until the beginning of the interventions. After pretreatment phase, eligible participants were randomly assigned into two parallel groups with a 1:1 allocation ratio to receive either SCTG following atraumatic extraction (test group) or minimally-traumatic extraction followed by spontaneous healing only (control group) based on the generated sequence. Due to the difference in the intervention's techniques, both the operating surgeon and the participants could not be blinded to the procedure. The outcome assessor (SN) and statistician (GNA) were blinded.

Volumetric analysis

The primary outcome in this clinical trial was the postoperative linear volumetric change of buccal soft tissue contour. Conventional polyvinyl siloxane impressions³ were taken at baseline, 3, and 6-month postoperative for each patient. The stone casts produced from the impressions were optically scanned using lab scanner⁴ to create digital surface models in Standard Triangle Language (STL) format, all STL files were imported to a digital software.⁵ The best-fit algorithm was used to superimpose digital surface models, when comparing each area of interest (AOI) throughout out the follow-up period. Unchanged neighboring teeth

surfaces were used as a reference for proper superimposition. In each patient, the AOI was kept constant for all pairwise comparisons. The volumetric analysis software calculated a mean linear volumetric change (mm) within the AOI for each patient at baseline, 3 months, and 6 months postoperative.

Clinical parameters

Clinical parameters were recorded at baseline and 3 and 6 months postoperatively by a single examiner (SN) who was blinded, trained, and calibrated with a good intra-examiner agreement (0.82 κ value). Clinical parameters recorded were gingival thickness (GT) and interdental papilla (IDP) height as secondary outcomes. GT measurements were recorded using anesthetic needle with a stopper technique [20], at levels of 1.5 mm, 3 mm, and 4.5 mm from the free gingival margin. The IDP height was recorded using UNC15 periodontal probe from the base to the tip of the IDP; the reference was the anatomical marks of the preoperative interdental papilla and the adjacent teeth. Accordingly, postoperative measurements were done using the UNC15 periodontal probe. A 6-month postoperative evaluation period was chosen based on the recommendation that final restorative measures should not be initiated until 6 months after ridge augmentation procedures. Furthermore, qualitative and probably quantitative alterations can arise during the healing period of the augmented soft tissue [21, 22].

Treatment protocol

Minimally-traumatic extraction

In both groups, flapless minimally-traumatic extraction was performed using peristomes⁶ inserted along the root surface; apical pressure and rocking motion were applied circumferentially to cut the periodontal ligaments. After initial luxation using Harpoon luxators,⁷ bayonet forceps were used to deliver the tooth out of the socket. Socket debridement was done to make sure the socket was thoroughly clean. For the control group, minimally-traumatic extraction followed by spontaneous healing was performed without soft tissue augmentation procedure.

Recipient site preparation (pouch technique + SCTG)

Patients assigned to the intervention group received pouch technique + SCTG represented in Fig. 2. Following minimally-traumatic extraction, preparation of a split

¹ OnDemand3D software, Cybermed, Seoul, Korea.

² The Arab Drug Company for pharmaceutical and Chemical Industries CO., Cairo, Egypt.

³ Hydrorise Putty and light, additional silicone, Zhermack SpA, Italy.

⁴ Shape lab scanner E series, Copenhagen, Denmark.

⁵ Exo cad, Fraunhofer Institute, Germany.

⁶ Helmut Zepf, Germany.




Fig. 2 Overview of minimally-traumatic extraction + SCTG. **a** Pre-operative clinical photograph occlusal view. **b** Minimally-traumatic extraction. **c** Initial incision of split thickness buccal pouch. **d** Extension of pouch preparation using tunneling knife. **e** SCTG harvesting

from the underlying bone by blunt dissection. **f** Harvested SCTG. **g** SCTG fitted in the buccal pouch. **h** SCTG secured in desired position by periosteal suture

thickness buccal pouch between the buccal bone plate and the overlying gingiva of the extraction socket was performed using 15c mini blade⁷ and tunneling knives.⁸ Initial partial dissection incision was done with an angle directed towards the buccal bone to avoid perforation of the overlying flap. The pouch preparation was then extended apical to the mucogingival junction and to the neighboring teeth mesiodistally using the tunneling knife to ensure sufficient flap mobility. The SCTG was harvested from the palate with a standardized thickness of 1.5 mm using single incision technique [23], which was done approximately 2–3 mm apical to the gingival margin of the maxillary teeth and parallel to the palatal long axis. Partial thickness flap was then raised and SCTG was separated by four down-to-bone incisions and harvested from the underlying bone by blunt dissection. The SCTG was contoured to fit the prepared pouch and then inserted and secured into the recipient site by a periosteal suture from the inside of the prepared pouch, using a resorbable suture material.⁸

Postsurgical phase

Postoperative analgesics⁹ three times daily and systemic antibiotics¹⁰ twice daily were prescribed for 5 days. Patients were instructed to rinse with 0.12% chlorhexidine

HCL2 three times a day for 2 weeks and to avoid any hard brushing and trauma to the surgical site. Cold ice packs were recommended to be placed extraorally adjacent to the surgical area during the first 12 h. For the test group, sutures were removed 14 days post-surgically.

Statistical and power analysis

Sample size calculation was based on finding a mean difference of at least 0.5 mm between the two studied groups with SD of 0.5 mm based on previous data [24]. Alpha was set at 0.05 and the power was set at 0.8. This resulted in the inclusion of 17 sites per group. To compensate for possible dropouts, 20 sites were included in each group, with a total of 40 sites. The sample size was calculated by PS program.¹¹ Data was explored for normality using Kolmogorov–Smirnov test and Shapiro–Wilk test. Continuous data showed normal distribution and were described using mean, standard deviation, mean difference, and 95% confidence interval. Repeated measures ANOVA test was used for comparisons between and within groups followed by Tukey post hoc test. Tukey post hoc test was used for pairwise comparisons when ANOVA test was significant. A *P* value less than or equal to 0.05 was considered statistically significant and all tests were two tailed. Data was analyzed using MedCalc software, version 19 for windows.¹²

⁷ Kiato™ 15c surgical blade, Kiato, Germany.

⁸ Tunneling Instrument, #1, Helmut Zepf, Germany.

⁹ Ibuprofen 600 mg Abbott, Egypt.

¹⁰ Amoxicillin 500 mg Cap., Egyptian Int. Pharmaceutical Industrial Co., A. R. E.

¹¹ Power and sample size program: biostat.mc.vanderbilt.edu/twiki/bin/view/Main/Power Sample Size.

¹² MedCalc Software Ltd, Ostend, Belgium.

Table 1 Volumetric change of buccal soft tissue contour and clinical parameters of both studied groups throughout the experimental period

Level of measurement from FGM	Time interval	Volumetric change of buccal soft tissue contour (mm)		Mean difference [95% CI]	P value
		STCG Mean (\pm SD)	Minimally-traumatic extraction + spontaneous healing Mean (\pm SD)		
1.5 mm	0–3	-0.38 ± 0.31	-2.05 ± 0.81	1.66 [1.23, 2.09]	$P < 0.001^*$
	0–6	-0.53 ± 0.40	-2.17 ± 0.80	1.64 [1.20, 2.09]	$P < 0.001^*$
	3–6	-0.14 ± 0.21	-0.12 ± 0.28	1.20 [0.84, 1.55]	$P = 0.837$
3 mm	0–3	0.07 ± 0.31	-1.47 ± 0.66	1.54 [1.18, 1.91]	$P < 0.001^*$
	0–6	-0.11 ± 0.44	-1.73 ± 0.63	1.61 [1.22, 1.99]	$P < 0.001^*$
	3–6	-0.19 ± 0.24	-0.26 ± 0.41	0.06 [-0.17, 0.30]	$P = 0.580$
4.5 mm	0–3	0.30 ± 0.39	-1.12 ± 0.57	1.43 [1.09, 1.77]	$P < 0.001^*$
	0–6	0.14 ± 0.50	-1.34 ± 0.69	1.48 [1.06, 1.90]	$P < 0.001^*$
	3–6	-0.17 ± 0.30	-0.22 ± 0.34	0.05 [-0.17, 0.27]	$P = 0.639$
Level of measurement from FGM	Time interval	Gingival thickness (mm)		Mean difference [95% CI]	P value
		STCG Mean (\pm SD)	Minimally-traumatic extraction + spontaneous healing Mean (\pm SD)		
1.5 mm	Baseline	1.26 ± 0.39	1.23 ± 0.46	0.02 [-0.25, 0.30]	$P = 0.831$
	3 months	5.26 ± 1.49	5.23 ± 1.99	0.02 [-1.20, 1.25]	$P = 0.961$
	6 months	3.52 ± 1.17	1.23 ± 0.43	2.29 [1.67, 2.91]	$P < 0.001^*$
3 mm	Baseline	1.26 ± 0.53	1.5 ± 0.68	$-0.23 [-0.66, 0.19]$	$P = 0.272$
	3 months	4.88 ± 1.37	3.05 ± 1.58	1.82 [0.78, 2.86]	$P < 0.001^*$
	6 months	3.14 ± 0.99	0.88 ± 0.41	2.26 [1.73, 2.79]	$P < 0.001^*$
4.5 mm	Baseline	0.79 ± 0.39	0.82 ± 0.39	$-0.02 [-0.30, 0.24]$	$P = 0.829$
	3 months	2.58 ± 1.18	1.17 ± 0.58	1.41 [0.75, 2.06]	$P < 0.001^*$
	6 months	2.20 ± 0.88	0.67 ± 0.24	1.52 [1.07, 1.98]	$P < 0.001^*$
Site	Time interval	IDP height (mm)		Mean difference [95% CI]	P value
		STCG Mean (\pm SD)	Minimally-traumatic extraction + spontaneous healing Mean (\pm SD)		
MIP	Baseline	2.76 ± 0.43	2.67 ± 0.70	0.08 [-0.32, 0.49]	$P = 0.664$
DIP		2.47 ± 0.37	2.55 ± 0.46	$-0.08 [-0.38, 0.20]$	$P = 0.545$
MIP	3 months	2.55 ± 0.68	1.9 ± 0.42	0.61 [0.22, 1.01]	$P = 0.003^*$
DIP		2.29 ± 0.43	1.85 ± 0.38	0.44 [0.15, 0.72]	$P = 0.003^*$
MIP	6 months	2.47 ± 0.51	1.61 ± 0.37	0.85 [0.53, 1.16]	$P < 0.001^*$
DIP		2.29 ± 0.30	1.61 ± 0.33	0.67 [0.45, 0.90]	$P < 0.001^*$

*Corresponds to statistically significant difference

Results

Initial buccal bone thickness

The SCTG group presented an initial buccal bone thickness with a mean of 0.75 ± 0.13 , while the minimally-traumatic extraction followed by spontaneous healing group showed a mean of 0.77 ± 0.15 . Intergroup analysis revealed a mean difference [95% CI] of $-0.021 [-0.12, 0.07]$, with no statistically significance (P value = 0.65) between the two groups regarding the baseline bone thickness.

Volumetric analysis

Buccal soft tissue contour

Table 1 shows volumetric changes recorded for both groups throughout the study. The presented results within the SCTG group showed that there was a statistically significant difference ($P < 0.05$) between different follow-up periods. Similarly, the minimally-traumatic extraction followed by spontaneous healing group demonstrated a statistically significant difference ($P < 0.05$) in the mean




Fig. 3 Case description including preoperative, 3 months, and 6 months postoperative clinical photographs and 0–6 months linear volumetric change of test group (from **a** to **h**) and control group (from **i** to **p**). **a** Preoperative clinical photograph, occlusal view, **b** preoperative clinical photograph, frontal view, **c** 3 months postoperative clinical photograph showing soft tissue contour, **d** 3 months postoperative clinical photograph showing IDP heights and vertical tissue level, **e** 6 months postoperative clinical photograph showing soft tissue contour, **f** 6 months postoperative clinical photograph showing IDP heights and vertical tissue level, **g** 0–6 months superimposed models by best fit algorithm, **h** 0–6 months superimposed models showing

gain of 0.5 mm in buccal soft tissue contour, **i** preoperative clinical photograph, occlusal view, **j** preoperative clinical photograph, frontal view, **k** 3 months postoperative clinical photograph showing soft tissue contour, **l** 3 months postoperative clinical photograph showing IDP heights and vertical tissue level, **m** 6 months postoperative clinical photograph showing soft tissue contour, **n** 6 months postoperative clinical photograph showing IDP heights and vertical tissue level, **o** 0–6 months superimposed models by best fit algorithm, and **p** 0–6 months superimposed models showing loss of 3.14 mm in buccal soft tissue contour

change of buccal soft tissue contour between follow-up periods. Intergroup analysis between the two modalities at 1.5 mm, 3 mm, and 4.5 mm revealed statistically significant difference from 0 to 3 months and 0 to 6 months ($P < 0.001$). However, there was no statistically significant difference at the same 3 measured points from 3 to 6 months ($P > 0.05$). Interestingly, sites treated with SCTG showed a statistically significant ($P < 0.001$) gain in mean buccal soft tissue contour compared to atraumatic extraction group after 6 months (Fig. 3).

Vertical tissue loss

The present results showed that the SCTG group presented less vertical tissue loss with mean (\pm SD) of $-0.66 (\pm 0.62)$, while the minimally-traumatic extraction followed by spontaneous healing group showed $-1.75 (\pm 0.73)$ with a

statistically significant difference ($P < 0.001$) between them from 0 to 6 months postoperatively.

Clinical outcomes

Gingival thickness

Table 1 shows GT recorded for both groups throughout the study. Both groups showed the same mean baseline values for GT at 1.5 mm, 3 mm, and 4.5 mm. There was a statistically significant ($P < 0.001$) increase in the mean GT at 3 and 6 months postoperatively compared to baseline values in the SCTG group. Meanwhile, the minimally-traumatic extraction followed by spontaneous healing group presented a statistically significant ($P < 0.05$) decrease in mean GT at 3 and 6 months postoperatively compared to baseline values. The SCTG group showed a statistically significant increase in mean GT at 1.5 mm, 3 mm, and

4.5 mm ($P < 0.001$) when compared to minimally-traumatic extraction followed by spontaneous healing group after 6 months.

Interdental papilla height

Table 1 shows IDP heights recorded for both groups throughout the study. This investigation observed that there was no statistically significant difference in both mesial IDP ($P = 0.664$) and distal IDP ($P = 0.545$) heights between the two groups at baseline, yet at 3 and 6 months there was a statistically significant difference ($P < 0.05$) in both mesial and distal IDP heights between both groups.

Discussion

Several techniques were suggested for alveolar ridge contour preservation including immediate implant placement, socket grafting, guided tissue regeneration concept, platelets concentrate, or other protocols like socket seal surgery and partial extraction therapy [2, 15, 25–27]. Despite their effectiveness and predictability, there are still few drawbacks regarding the esthetic outcome, since they mainly focus on hard tissue regeneration [28–30]. Furthermore, none of the proposed protocols could entirely avoid the soft tissue remodeling phenomenon, even some cases resulted in an esthetic discrepancy with an observed buccal concavity [31–33]. With regard to the above-mentioned gap of knowledge, the present trial targeted only the impact of soft tissue augmentation by SCTG on the post-extraction volumetric changes of buccal soft tissue contour.

Soft tissue augmentation techniques, such as bilaminar technique/coronally advanced flap and the pouch procedure with CTG, have emerged to address esthetic concerns in previous techniques. Such protocols showed predictability and improved soft tissue dehiscence and volume loss around dental implants [34–36]. Moreover, Marzadori et al. [37] concluded that the pouch technique with SCTG is the ultimate choice for soft tissue augmentation, particularly in areas of high esthetic demands, maintaining color and soft tissue appearance. To the best of the authors' knowledge, this study is the first randomized controlled clinical trial comparing minimally-traumatic extraction with SCTG to minimally-traumatic extraction followed by spontaneous healing, investigating the influence of soft tissue management in fresh extraction sockets using volumetric analysis. No socket sealing material was used in the present study, in order to allow the inflammatory phase of the socket healing to begin [38], without any factors that might influence the healing process. Thus, this randomized clinical trial investigated the influence of soft tissue management solely in

fresh extraction sockets via volumetric analysis without any confounding factors.

Post-extraction volumetric buccal soft tissue change is of utmost clinical interest, since the loss of root convexity may lead to unfavorable esthetic results and requires additional augmentation procedures. Accordingly, linear volumetric change of buccal soft tissue contour was considered the primary outcome in this study together with vertical tissue loss, gingival thickness, and interdental papilla height. A 6-month postoperative evaluation period was chosen based on the recommendation that final restorative measures should not be initiated until 6 months. Furthermore, qualitative and probably quantitative alterations can arise during the healing period of the augmented soft tissue [21, 22]. The accuracy of volume measurement with optical scanning-based digital technologies showed tremendous precision and high reproducibility [39]. Consequently, digitalized volumetric analysis was used to assess the linear volumetric change of buccal soft tissue contour and vertical tissue loss in the present clinical trial.

The findings presented herein showed that minimally-traumatic extraction followed by spontaneous healing without soft tissue augmentation caused a significant decrease in the buccal soft tissue contour, vertical tissue level, gingival thickness, and interdental papillae height after 6 months compared to baseline. These findings align with previous studies investigating spontaneous socket healing [40, 41]. Schneider et al. [40] observed buccal contour loss in spontaneously healing sites compared to sites treated with ridge preservation techniques. This was also supported by Thoma et al. [41] who reported a buccal soft tissue collapse of 1.2–1.6 mm in unassisted healing sites. The current study confirms the unsatisfactory esthetic results in unassisted socket healing presented in the literature [42, 43]. However, Clementini et al. [44] reported an increase in buccal soft tissue profile and thickness in unassisted healing sites after 4 months, attributing it to spontaneous soft tissue thickening phenomena. Similarly, Chappuis et al. [45] observed spontaneous soft tissue thickening in thin bone phenotypes after 8 weeks. This discrepancy might be attributed to the fact that the current investigation evaluated the outcomes after 6 months, representing the full maturation of hard and soft tissues at the extraction site [38]. Regarding the interdental papilla height, the minimally-traumatic extraction followed by spontaneous healing group showed decreased IDP heights over time, likely due to the thin phenotype of the treated sites. Previous studies on papilla fill around implant restorations supported these results, indicating that gingival phenotype influenced papilla volume/fill [46–48].

On the other hand, results of this clinical trial showed that the use of SCTG buccal to extraction sockets significantly preserved buccal soft tissue contour, reduced vertical

tissue loss, increased gingival thickness, and maintained interdental papillae height after 6 months. These findings were consistent with previous studies showing the benefits of SCTG in improving soft tissue profiles and correcting alveolar ridge contour defects [22, 24, 34, 49–51]. Furthermore, utilizing SCTG with a pouch preparation buccal to fresh extraction sockets limited post-extraction soft tissue alterations and yielded positive outcomes. These observations were supported by a recent systematic review by Zucchelli et al. [16], which emphasized the esthetic improvement and long-term stability achieved with SCTG in managing ridge deformities in the esthetic zone. The volumetric buccal soft tissue changes from baseline to 6 months in the SCTG group revealed a mean \pm SD change of -0.53 ± 0.40 , -0.11 ± 0.44 , and 0.14 ± 0.50 mm at 1.5 mm, 3 mm, and 4.5 mm from the free gingival margin, respectively. These results were superior to the use of “saddled” connective tissue graft combined with biomaterials [34] and were inferior to those reported by De Bruyckere et al. [24] who compared SCTG with GBR to reestablish ridge profile at the buccal aspect of single implants. These differences might be attributed to the different timing of interventions and prosthetic involvement in shaping the augmented soft tissue. It is worthy to mention that the baseline assessment in this investigation was performed with the presence of the natural tooth transgingival support. Besides, no soft tissue shaping occurred during the follow-up period. Remarkably, the currently observed findings might suggest that the use of the pouch and SCTG technique might conceal the post-extraction buccal concavity.

The current statistical analysis revealed that all the outcomes measured in the minimally-traumatic extraction with SCTG group surpassed those of the minimally-traumatic extraction followed spontaneous healing group solely with no soft tissue augmentation after 6 months. These observations suggest that the presence of the SCTG might effectively increase the soft tissue volume and mask the post-extraction contour deficiencies, especially in the esthetic zone. Furthermore, the increased gingival thickness observed after the use of SCTG might imply its ability to act as a biologic filler, improve the stability of the interdental tissues, and enhance the gingival phenotype. The current data sheds light on the effectiveness of the pouch and SCTG technique in preserving alveolar ridge contour during post-extraction modeling and remodeling. Future studies are warranted to explore the influence of intraoral scanners on volumetric analysis, in addition to examining the use of prosthetic devices and their impact on soft tissue shaping after augmentation.

One of the limitations of the present clinical trial was that a third arm of bone graft placement into the socket as well as connective tissue graft should have been performed. Another limitation was that IDP measurements were not calibrated.

Conclusions

Within the limitations of this randomized clinical trial study, it can be concluded that the pouch technique utilizing a SCTG buccal to fresh extraction sockets might be considered as an effective modality for alveolar ridge contour preservation and ridge profile stabilization up to 6 months. This study paves the way for future randomized clinical trials with longer follow-up periods.

Acknowledgements The authors are immensely grateful to Dr. Mina Magdy Selim for his hard work and efforts in the volumetric analysis process.

Author contribution Nourhan Gamal performed of the surgical procedures, wrote the main manuscript text, and prepared the figures. Nesma Shemais assessed the outcomes and critically revised the main manuscript. Marwa Al-Nawawy concealed the allocation sequence and critically revised the main manuscript. Noha Ghallab performed the statistical analysis, wrote the main manuscript, and revised the tables and figures. All authors approved the final manuscript.

Funding Open access funding provided by The Science, Technology & Innovation Funding Authority (STDF) in cooperation with The Egyptian Knowledge Bank (EKB).

Data Availability Data is available by contacting the author upon request.

Declarations

Competing interests The authors declare no competing interests.

Ethics approval and consent to participate The present randomized clinical trial was approved by the Research Ethics Committee, Faculty of Dentistry, Cairo University (13-12-19), conducted in accordance with the Helsinki Declaration of 1975, as revised in 2013. All patients provided written informed consent to participate in this trial.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Cardaropoli G, Araújo M, Lindhe J (2003) Dynamics of bone tissue formation in tooth extraction sites. An experimental study in dogs. *J Clin Periodontol* 30(9):809–18. <https://doi.org/10.1034/j.1600-051x.2003.00366.x>
2. Jung RE, Ioannidis A, Hämmerle CHF, Thoma DS (2018) Alveolar ridge preservation in the esthetic zone. *Periodontol* 2000 77(1):165–75. <https://doi.org/10.1111/prd.12209>

3. Thoma DS, Mühlemann S, Jung RE (2014) Critical soft-tissue dimensions with dental implants and treatment concepts. *Periodontol* 2000 66(1):106–18. <https://doi.org/10.1111/prd.12045>
4. Imber JC, Rocuzzo A, Stähli A et al (2023) Spontaneous regeneration of keratinized tissue at implants and teeth. *J Clin Periodontol* 50(8):1064–1074. <https://doi.org/10.1111/jcpe.13820>
5. Cortellini P, Pini Prato G (2012) Coronally advanced flap and combination therapy for root coverage. Clinical strategies based on scientific evidence and clinical experience. *Periodontol* 2000 59(1):158–84. <https://doi.org/10.1111/j.1600-0757.2011.00434.x>
6. Zuhr O, Bäumer D, Hürzeler M (2014) The addition of soft tissue replacement grafts in plastic periodontal and implant surgery: critical elements in design and execution. *J Clin Periodontol* 41(Suppl 15):S123–S142. <https://doi.org/10.1111/jcpe.12185>
7. Zucchelli G, Mounssif I (2015) Periodontal plastic surgery. *Periodontol* 2000 68(1):333–68. <https://doi.org/10.1111/prd.12059>
8. Rocuzzo M, Rocuzzo A, Marruganti C, Fickl S (2023) The importance of soft tissue condition in bone regenerative procedures to ensure long-term peri-implant health. *Periodontol* 2000. <https://doi.org/10.1111/prd.12496>
9. Bienz SP, Carrera ER, Hüslér J, Rocuzzo A, Jung RE, Thoma DS (2023) Soft tissue contour changes at implant sites with or without soft tissue grafting in the esthetic zone: a retrospective case-control study with a 12-year follow-up. *Clin Oral Implants Res* 34(5):521–530. <https://doi.org/10.1111/clr.14058>
10. Zuhr O, Hürzeler M (2012) Plastic-esthetic periodontal and implant surgery, 1st edition
11. Thoma DS, Benić GI, Zwahlen M, Hämmerle CH, Jung RE (2009) A systematic review assessing soft tissue augmentation techniques. *Clin Oral Implants Res* 20(Suppl 4):146–165. <https://doi.org/10.1111/j.1600-0501.2009.01784.x>
12. Cairo F, Nieri M, Pagliaro U (2014) Efficacy of periodontal plastic surgery procedures in the treatment of localized facial gingival recessions. A systematic review. *J Clin Periodontol* 41(Suppl 15):S44–62. <https://doi.org/10.1111/jcpe.12182>
13. Zucchelli G, Felice P, Mazzotti C et al (2018) 5-Year outcomes after coverage of soft tissue dehiscence around single implants: a prospective cohort study. *Eur J Oral Implantol* 11(2):215–224
14. Tavelli L, Barootchi S, Nguyen TVN, Tattan M, Ravidà A, Wang HL (2018) Efficacy of tunnel technique in the treatment of localized and multiple gingival recessions: a systematic review and meta-analysis. *J Periodontol* 89(9):1075–1090. <https://doi.org/10.1002/jper.18-0066>
15. Thoma DS, Buranawat B, Hämmerle CH, Held U, Jung RE (2014) Efficacy of soft tissue augmentation around dental implants and in partially edentulous areas: a systematic review. *J Clin Periodontol* 41(Suppl 15):S77–91. <https://doi.org/10.1111/jcpe.12220>
16. Zucchelli G, Tavelli L, McGuire MK et al (2020) Autogenous soft tissue grafting for periodontal and peri-implant plastic surgical reconstruction. *J Periodontol* 91(1):9–16. <https://doi.org/10.1002/jper.19-0350>
17. Jung RE, Becker K, Bienz SP et al (2022) Effect of peri-implant mucosal thickness on esthetic outcomes and the efficacy of soft tissue augmentation procedures: consensus report of group 2 of the SEPA/DGI/OF workshop. *Clin Oral Implants Res* 33(Suppl 23):100–8. <https://doi.org/10.1111/clr.13955>
18. Tonetti MS, Jung RE, Avila-Ortiz G et al (2019) Management of the extraction socket and timing of implant placement: consensus report and clinical recommendations of group 3 of the XV European Workshop in Periodontology. *J Clin Periodontol* 46(Suppl 21):183–194. <https://doi.org/10.1111/jcpe.13131>
19. Moher D, Hopewell S, Schulz KF et al (2012) CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. *Int J Surg* 10(1):28–55. <https://doi.org/10.1016/j.ijsu.2011.10.001>
20. Paolantonio M, Dolci M, Esposito P et al (2002) Subpedicle acellular dermal matrix graft and autogenous connective tissue graft in the treatment of gingival recessions: a comparative 1-year clinical study. *J Periodontol* 73(11):1299–1307. <https://doi.org/10.1902/jop.2002.73.11.1299>
21. Lindhe J, Karring TP, Lang N (2008) Clinical periodontology and implant dentistry, 5th edition
22. Akcalı A, Schneider D, Ünlü F, Bıçakçı N, Köse T, Hämmerle CH (2015) Soft tissue augmentation of ridge defects in the maxillary anterior area using two different methods: a randomized controlled clinical trial. *Clin Oral Implants Res* 26(6):688–695. <https://doi.org/10.1111/clr.12368>
23. Hürzeler MB, Weng D (1999) A single-incision technique to harvest subepithelial connective tissue grafts from the palate. *Int J Periodontics Restorative Dent* 19(3):279–287
24. De Bruyckere T, Cabeza RG, Eghbali A, Younes F, Cleymaet R, Cosyn J (2020) A randomized controlled study comparing guided bone regeneration with connective tissue graft to reestablish buccal convexity at implant sites: a 1-year volumetric analysis. *Clin Implant Dent Relat Res* 22(4):468–476. <https://doi.org/10.1111/cid.12934>
25. Mendoza-Azpur G, Olaechea A, Pinazo M et al (2015) Histomorphometric evaluation of ridge preservation with and without connective tissue graft over buccal plate using different types of bone substitute: an animal study. *Implant Dent* 24(6):686–692. <https://doi.org/10.1097/ID.0000000000000307>
26. Buser D, Chappuis V, Belser UC, Chen S (2017) Implant placement post extraction in esthetic single tooth sites: when immediate, when early, when late? *Periodontol* 2000 73(1):84–102. <https://doi.org/10.1111/prd.12170>
27. Al Yafi F, Alchawaf B, Nelson K (2019) What is the optimum for alveolar ridge preservation? *Dent Clin North Am* 63(3):399–418. <https://doi.org/10.1016/j.cden.2019.02.007>
28. Araújo MG, Sukekava F, Wennström JL, Lindhe J (2005) Ridge alterations following implant placement in fresh extraction sockets: an experimental study in the dog. *J Clin Periodontol* 32(6):645–652. <https://doi.org/10.1111/j.1600-051X.2005.00726.x>
29. Araújo MG, Sukekava F, Wennström JL, Lindhe J (2006) Tissue modeling following implant placement in fresh extraction sockets. *Clin Oral Implants Res* 17(6):615–624. <https://doi.org/10.1111/j.1600-0501.2006.01317.x>
30. Araújo MG, Wennström JL, Lindhe J (2006) Modeling of the buccal and lingual bone walls of fresh extraction sites following implant installation. *Clin Oral Implants Res* 17(6):606–614. <https://doi.org/10.1111/j.1600-0501.2006.01315.x>
31. Stumbras A, Kuliesius P, Januzis G, Juodzbaly G (2019) Alveolar ridge preservation after tooth extraction using different bone graft materials and autologous platelet concentrates: a systematic review. *J Oral Maxillofac Res* 10(1):e2. <https://doi.org/10.5037/jomr.2019.10102>
32. Wu DT, Raouf S, Latimer JM, Nguyen TT (2021) Partial extraction therapy: a review of human clinical studies. *J Oral Implantol*. <https://doi.org/10.1563/aid-D-21-00095>
33. López-Pacheco A, Soto-Peñaloza D, Gómez M, Peñarrocha-Oltra D, Alarcón MA (2021) Socket seal surgery techniques in the esthetic zone: a systematic review with meta-analysis and trial sequential analysis of randomized clinical trials. *Int J Implant Dent* 7(1):13. <https://doi.org/10.1186/s40729-021-00294-2>
34. Vanhoutte V, Rompen E, Lecloux G, Rues S, Schmitter M, Lambert F (2014) A methodological approach to assessing alveolar ridge preservation procedures in humans: soft tissue profile. *Clin Oral Implants Res* 25(3):304–309. <https://doi.org/10.1111/clr.12144>
35. Kan JYK, Rungcharassaeng K, Deflorian M, Weinstein T, Wang HL, Testori T (2018) Immediate implant placement and provisionalization of maxillary anterior single implants. *Periodontol* 2000 77(1):197–212. <https://doi.org/10.1111/prd.12212>
36. Ramanauskaitė A, Obreja K, Sader R, Becker J, Schwarz F (2021) Assessment of soft and hard tissue dimensions following different

- treatment approaches of ligature-induced peri-implantitis defects. *Clin Oral Implants Res* 32(4):394–400. <https://doi.org/10.1111/clr.13709>
37. Marzadori M, Stefanini M, Mazzotti C, Ganz S, Sharma P, Zucchelli G (2018) Soft-tissue augmentation procedures in edentulous esthetic areas. *Periodontol 2000* 77(1):111–22. <https://doi.org/10.1111/prd.12210>
 38. Araújo MG, Silva CO, Misawa M, Sukekava F (2015) Alveolar socket healing: what can we learn? *Periodontol 2000* 68(1):122–34. <https://doi.org/10.1111/prd.12082>
 39. Ueno D, Kobayashi M, Tanaka K et al (2018) Measurement accuracy of alveolar soft tissue contour using a laboratory laser scanner. *Odontology* 106(2):202–207. <https://doi.org/10.1007/s10266-017-0315-4>
 40. Schneider D, Schmidlin PR, Philipp A et al (2014) Labial soft tissue volume evaluation of different techniques for ridge preservation after tooth extraction: a randomized controlled clinical trial. *J Clin Periodontol* 41(6):612–617. <https://doi.org/10.1111/jcpe.12246>
 41. Thoma DS, Bienz SP, Lim HC, Lee WZ, Hämmerle CHF, Jung RE (2020) Explorative randomized controlled study comparing soft tissue thickness, contour changes, and soft tissue handling of two ridge preservation techniques and spontaneous healing two months after tooth extraction. *Clin Oral Implants Res* 31(6):565–574. <https://doi.org/10.1111/clr.13594>
 42. Faria-Almeida R, Astramskaite-Januseviciene I, Puisys A, Correia F (2019) Extraction socket preservation with or without membranes, soft tissue influence on post extraction alveolar ridge preservation: a systematic review. *J Oral Maxillofac Res* 10(3):e5. <https://doi.org/10.5037/jomr.2019.10305>
 43. Canullo L, Pesce P, Antonacci D et al (2022) Soft tissue dimensional changes after alveolar ridge preservation using different sealing materials: a systematic review and network meta-analysis. *Clin Oral Investig* 26(1):13–39. <https://doi.org/10.1007/s00784-021-04192-0>
 44. Clementini M, Castelluzzo W, Ciaravino V et al (2020) The effect of immediate implant placement on alveolar ridge preservation compared to spontaneous healing after tooth extraction: soft tissue findings from a randomized controlled clinical trial. *J Clin Periodontol* 47(12):1536–1546. <https://doi.org/10.1111/jcpe.13369>
 45. Chappuis V, Engel O, Shahim K, Reyes M, Katsaros C, Buser D (2015) Soft tissue alterations in esthetic postextraction sites: a 3-dimensional analysis. *J Dent Res* 94(9 Suppl):187s-s193. <https://doi.org/10.1177/0022034515592869>
 46. Zetu L, Wang HL (2005) Management of inter-dental/inter-implant papilla. *J Clin Periodontol* 32(7):831–839. <https://doi.org/10.1111/j.1600-051X.2005.00748.x>
 47. Romeo E, Lops D, Rossi A, Storelli S, Rozza R, Chiapasco M (2008) Surgical and prosthetic management of interproximal region with single-implant restorations: 1-year prospective study. *J Periodontol* 79(6):1048–1055. <https://doi.org/10.1902/jop.2008.070431>
 48. Si MS, Zhuang LF, Huang X, Gu YX, Chou CH, Lai HC (2012) Papillae alterations around single-implant restorations in the anterior maxillae: thick versus thin mucosa. *Int J Oral Sci* 4(2):94–100. <https://doi.org/10.1038/ijos.2012.25>
 49. González-Martín O, Veltri M, Moráquez O, Belser UC (2014) Quantitative three-dimensional methodology to assess volumetric and profilometric outcome of subepithelial connective tissue grafting at pontic sites: a prospective pilot study. *Int J Periodontics Restorative Dent* 34(5):673–9. <https://doi.org/10.11607/prd.1808>
 50. Papace C, Büsch C, Ristow O, Keweloh M, Hoffmann J, Mertens C (2021) The effect of different soft-tissue management techniques for alveolar ridge preservation: a randomized controlled clinical trial. *Int J Implant Dent* 7(1):113. <https://doi.org/10.1186/s40729-021-00390-3>
 51. Ammar AH, Ahmed E, ElBarbary A, Ghalwash D, Ezz Elarab A (2022) Clinical comparison of the volumetric changes in single pontic site development through connective tissue grafting using modified pouch technique versus pouch technique in the maxillary esthetic zone: a randomized controlled clinical trial. *Int J Dent* 2022:1677471. <https://doi.org/10.1155/2022/1677471>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.