

Biological and serological studies on *Pea seed-borne mosaic virus (PSbMV)* on cowpea forage crop (*Vigna unguiculata L. Walp*)

Kararah M. A¹, Om-Hashem M.El-Banna¹, Salwa N. Zein² and Abd-Elrehim, A.F³

¹Plant Pathol. Dept. Fac. Agric. Cairo Univ. Giza, Egypt² Virus and Phytoplasma Res. Dept., Plant Patho.Res. Inst., Agric. Res. Center, Giza, Egypt³Central Administration of Plant Quarantine, Ministry of Agriculture and Land Reclamation, Egypt

ABSTRACT

In the present work, *Pea seed-borne mosaic* (PSbMV) was first reported on cowpea (*Vigna unguiculata* (L.) Walp). PSbMV was isolated from naturally infected cowpea plants, showing mosaic and chlorotic ring spot symptoms, which had been grown in the experimental fields of Giza Agricultural Research Experimental Station (A.R.E.S) in 2008. Identification studies based on host range, symptomatology and seed transmission through cowpea and different hosts belong to Fabaceae. The results indicated that the host range of the virus was expanded to seven different plant families. The virus was mechanically transmitted, and the percentage of seed transmission was 6%. PSbMV was partially purified for its antiserum production. The absorption spectrum of the purified virus had a min at 245 nm and a max at 260 nm. The ratios of A₂₆₀/A₂₈₀, A₂₈₀/A₂₆₀ and A_{max}/A_{min} were 1.16, 0.85, and 1.04, respectively. Yield of PSbMV was 1.7 mg/100g of infected leaves. Electron micrographs of the partially purified virus preparation revealed the presence of filamentous flexuous virus particles about 700- 750nm long. Titers of the antisera after first, second and third bleeding were 1/800, 1/1600 and 1/3200 respectively. The optimum concentrations of IgG and IgG conjugate were 1.0 µg/ml and 1/1000, respectively. The antigen dilution end point was 1:500. The produced antiserum was evaluated by ELISA and DBIA. Electron microscopy of ultrathin-section of PSbMV- infected leaf tissue revealed several morphological changes and aggregated virus particles associated with pinwheel inclusion bodies. Salicylic acid and Parahydroxy benzoic acid were used for induction of systemic acquired resistance, whereas camphor oil was used as an antiviral agent to reduce PSbMV infection. Parahydroxy benzoic acid was effective at 0.001% concentration, exhibited 88.7% reduction when had applied 2h before inoculation. Also, salicylic acid was effective as it reduced infection to 77.4% at 0.01% concentration when applied 3h before inoculation. Whereas camphor oil exhibited lesser reduction being (74.6) when used at 0.001% concentration before 2 h from inoculation.

Key words: *Pea seed borne mosaic Potyvirus*, Host range, Seed transmission, Purification, Electron microscopy, IgG, ELISA, DIBA, Salicylic acid, Parahydroxy benzoic acid, Camphor oil.

INTRODUCTION

Viral diseases in field-grown cowpea plants are the major concern for the forage crops, feed leguminous growers and the rate of production (Cuttings yield). Cowpea growing areas have been affected by several viruses (Gillaspie *et al.*, 1995 and Taiwo *et al.*, 2007).

Pea seed borne mosaic Potyvirus (PSbMV), an economically significant seed – transmitted virus of pea has commonly found in pea germplasm collection of many countries. The virus is suspected to have spread world- wide due to the exchange of infected germplasm material (Khetarpal and Maury, 1987).

Various percentage of seed transmission of PSbMV was recorded by several authors depending on different factors like growth conditions, cultivar genotype, seed size and the severity of PSbMV strain . (Hmpton and Mink, 1975 and Makkouk *et al.* 1988)

Different serological methods were reported to detect *Pea seed borne mosaic virus* including enzyme linked immune sorbent assay (ELISA), Dot-blot immunoassay (DBIA) and Tissue blot immunoassay (TBIA) on nitrocellulose membrane.

Aiming to minimize the negative effect of pesticides are been development the alternative control of plant disease, which includes the induction of resistance and the use of natural products with induction of resistance and /or with direct antimicrobial activities. In the latter include the use of extracts and essential oils from medicinal plants (Stangarlin *et al.*, 1991)

Several chemical and natural compounds including salicylic acid, benzoic acid and natural oils were reported to use for plant virus control purposes and effective results was obtained (Kobeasy & Zein, Salwa 2005).

The aim of the current study was: Firstly, isolation and identification of the causal agent found in collected cowpea samples using the traditional (biological) methods. Secondly, purification and production of ELISA reagents which can be used as a rapid serological diagnosis for PSbMV. Thirdly cytological studies using electron microscopy. Finally, the control measures using different applications such as antiviral agent and induced systemic resistance against the virus isolate.

MATERIALS AND METHODS

Part I: Isolation and identification:

1. Isolation:

Samples of naturally infected cowpea (*Vigna unguiculata* L.) plants showed mosaic and chlorotic ring spot symptoms were collected from Giza Agricultural Research experimental Station (A.R.E.S) in 2008. The collected samples were screened for viruses by ELISA test using antisera of Cucumber mosaic virus (CMV), Broad bean true mosaic virus (BBTMV), Bean yellow mosaic virus (BYMV), Pea seed borne mosaic virus (PSbMV), Pepper mild mottle virus (PMMoV) and Bean common mosaic virus (BCMV). Polyclonal antibodies obtained from Serology Lab., Virus Research Dept., Plant Pathology Research Institute were used in indirect ELISA as described by Clark and Adams (1977). The virus isolate was biologically purified by single local lesion technique (Kuhn, 1964). The virus isolate was propagated in pea plants as a main host while *Ch. amaranticolor* was used as local lesion host. Cowpea plants reacted positively only against PSbMV antiserum were served as source of virus inoculum.

2- Identification of the virus isolate:

2.1. Host range and diagnostic host reactions:

The isolated virus was mechanically inoculated in 39 selected hosts including certain diagnostic hosts belonging to *Alliaceae*, *Cucurbitaceae*, *Chenopodiaceae*, *Fabaceae*, *Graminaceae*, *Brassicaceae* and *Solanaceae*. Seedlings of each host plant were maintained four weeks in the greenhouse for symptoms development at temperature average 25-30° C. An equal number (10 seedlings) of healthy seedlings of the same age and species were left as a control. Symptomless plants were assessed biologically and serologically by indirect ELISA method.

2.2. Transmission through cowpea (*V. unguiculata* (L.) Walp) seeds:

Healthy cowpea plants (Buffalo.cv) were mechanically inoculated with PSbMV. The inoculated plants were kept inside an insect-proof greenhouse. Two weeks after inoculation, indirect ELISA was used to check the virus presence in cowpea plants. Plants, which gave positive reaction, were labelled and kept till seed formation. Two hundred seeds were collected from infected plants and tested for seed transmission as they were sown

in pots (25 cm diameter) containing sterilized clay soil. Seedlings were kept in the greenhouse. Percentage of seed transmission was calculated as number of seedling showing symptoms and gave positive reaction with Indirect ELISA divided by No. of immersed seedlings X 100.

3- Serological detection of the virus isolate:

Indirect ELISA technique was performed to confirm the identity of the virus isolate

Part II-Virus purification and ELISA reagents production:

1. Purification of the virus isolate:

The inoculated pea plants by the virus isolate were used as a source for virus purification using a modified method described by Gamal El-Din *et al.* (1997).

1.1. U.V. absorptionspectrum of the purified virus:

U.V. absorption spectrum of the purified virus preparation was estimated spectrophotometrically with Spectronic 2000 at range from 220 to 320 nm. Virus concentration was calculated using the extinction coefficient of 2.4 (purcifull, 1990).

1.2. Electron microscopy:

Partially purified virus preparations were negatively stained with 2% urinal acetate, mounted on, formvar-coated grids, examined with a JeolTEM-1400 in the Electron Microscope Unit, Faculty of Agriculture, Cairo University and virus particles were photographed (Noordam, 1973)

2. Antiserum production

2.1. Rabbit immunization and bleeding

Polyclonal antibodies were prepared by injecting New Zealand white rabbit intravenously eight times with 11mg. of purified PSbMV preparations. Serum was collected 7 days after the last injection along 3 weeks. Antiserum was separated by centrifugation at 5,000 rpm for 3 min. (Hampton *et al.*, 1990). The antisera was collected and stored at 4 C° until used.

2.2. Determination of antiserum titre:

Antiserum was cross-absorbed with healthy plant proteins using a method by Hampton *et al.* (1990). Antiserum titer was determined using indirect ELISA technique. Clarified sap of virus infected (PSbMV) and healthy tobacco (*N. tabacum*, White Burley) plants, was diluted at 1/3. PSbMV antiserum was diluted to 1/100, 1/200, 1/400,

1/800, 1/1600 and 1/6400 (Lommelet *et al.*, 1982).

2.3. Purification of the immunoglobulin G (IgG).

IgG was purified from PSbMV antiserum using the caprylic acid method described by Steinbuch and Audran (1969).

2.4. Conjugation of IgG with alkaline phosphatase:

IgG was conjugated with alkaline phosphatase (AP) according to protocol given by **Clark and Adams (1977)**. The concentrations of IgG and IgG conjugate were determined by a checkerboard test (Converse and Martin, 1990) evaluating a series of conjugate dilutions against a series of trapping globulin dilutions using infected and healthy plant preparations of the same species. IgG conjugate was diluted to 1/250, 1/500, 1/1000 and 1/2000 with the conjugate buffer, while IgG was diluted with the coated buffer to concentration of 0.5, 1.0, 2.0 and 4.0 µg/ml, respectively. Controls of healthy and infected cowpea sap were used. The reaction was done between IgG and IgG conjugate using DAS ELISA.

2.5. Determination of antigen dilution end point:

ELISA reagents specific to PSbMV were assembled from IgG and IgG conjugated with alkaline phosphatase,

positive and negative controls of the virus were diluted with the coated buffer to concentrations 1/500, 1/1000, 1/1500 and 1/2000.

2-6- Serological detection of PSbMV by DAS-ELISA and DBIA using the ELISA reagent produced:

2-6-1- Enzyme linked immunosorbent assay (ELISA):

The produced ELISA reagents were used in detection of PSbMV in seed lots of different plant species as described by Converse Martin (1990).

This experiment was carried out at Agricultural Research Experimental Station. Mature seeds were collected from lots to estimate percentage of natural infection with PSbMV through seeds of Chickpea (*Cicer arietinum*), Lentil (*Lens culinaris*), Pea (*Pisum sativum*), Faba bean (*Vicia faba*) and Peanut (*Arachis hypogaea*). ELISA reagents produced were used to detect the presence of PSbMV in the seeds after they were washed in running tap water and kept for 48 hr. in Petri dishes with wet cotton before homogenized and assayed by DAS-ELISA. Percentages of seed infection were calculated.

2-6-2- Dot- Blotting Immunobinding Assay on

nitrocellulose membrane (DBIA):

The DBIA on nitrocellulose membranes was essentially similar to those described by Hsu and Lawson (1991) and Azzam *et al.* (2007) for the serological detection of antigen using authentic antiserum for PSbMV.

Part III- cytological studies using electron microscopy (EM).

Ultra-thin sections taken from mechanically inoculated tobacco leaves as well from cowpea seedlings immersed from infected seeds. The used technique was the same as described by Bozzala and Russell (1999). This work was done in EM-Lab in Faculty of Agriculture- Cairo University Research Park (FARP).

Part IV- Virus control:

Salicylic acid, parahydroxy benzoic acid and camphor oil were used to test their inhibitory effect of PSbMV infection. Ten seedlings at 2 weeks stage for each treatment were sprayed with the compounds under investigation. The salicylic acid solution was prepared as described by Yalpani, *et al.* (1991) and Kobeasy & Salwa, Zein (2005), benzoic acid (White & Antoniw *et al.*, 1986 and Smith-Backer *et al.*, 1998) Camphor oil (Ahmed, Amal *et al.*, 2010) were used as foliar

sprays. Each of the inhibitors were applied by spraying one 1h, 2h, 3h, and 4 h before mechanical inoculation using different concentrations i.e., 0.0001, 0.001, 0.01, respectively.

Fifteen leaves of the local lesion host plant dusted with carborundum (600 mesh) for each treatment were inoculated with the virus isolate. Total number of local lesions for each treatment was counted two weeks post virus inoculation. Equal number of the plants were inoculated with the virus isolate and kept without spraying in the greenhouse as control. Efficiency of the inhibition was determined as described by Devi *et al.* (2004) using the following equation:

$$\text{Inhibition \%} = \frac{A-B}{A} \times 100$$

Where: A= Control, B= treatment

RESULTS**Part I: Isolation and identification:****1- Isolation:**

The virus isolated from infected (*V. unguiculata*) plants was biologically purified through single local lesion transfers on *C. amaranticolor*. The resulting virus was propagated on *pea plants*. It was identified by Indirect ELISA test using authentic polyclonal antisera obtained from the serological lab, Virus and Phytoplasma Research Department.

2- Identification of the virus isolate:

2-1- Host range and diagnostic host reactions:

Result in Table (1) and Fig (1) show the reaction of forty plant species and cultivar belong to seven different families to virus infection. Inoculated plants showed a great variability in symptoms expression, such as mosaic, mottling, deformed leaves, chlorotic and necrotic local lesions.

It was also indicated that the tested hosts belonging to family Solanaceae and Fabaceae susceptible to infection with PSbMV. Some hosts showed no obvious symptoms and when checked by ELISA gave positive reaction (symptomless).

It is obvious from Table (1) that species and cultivars tested can be classified into four general groups:

(1) Plants reacted only with local symptoms: *Chenopodium quinoa*, *C. amarantifolium*, *Coste & reyne Datur ainnoxia* L. and *Nicotiana glutinosa* L.

(2) Plants reacted only with systemic symptoms: *Vigna unguiculata* Cv. beef, *Pisum sativum* cv. Litel Marvel, *Vicia faba* cv. Giza 3, *Cicer arietinum* cv. Giza 531, *Lens culinaris* cv. Kordy1, and *N. tabacum* cultivars *White*, *Samsun* and *Xanthi*.

(3) Plants didn't show symptoms, but from which

PSbMV could be recovered either by back inoculation onto the indicator host plants or by ELISA test include the following: *Glycine max* cv. Crawford, *Phaseolus vulgaris* L. cv. Bronco, *Vicia faba* L. cv. Giza 3, *Cicer arietinum* cv. Giza 531, *Lens culinaris* cv. Kordy1, *Hordeum vulgare* L. cv. G123, *Triticum aestivum* cv. sods1,

(4) Plants didn't show any symptoms and PSbMV couldn't be recovered by either by back inoculation nor by ELISA, these are:

Allium cepa cv. Giza 6, *Allium sativum* cv. Baladi, *Beta vulgaris* cv. Pleno, *Glycine max* cultivars Giza 35 and Clark, *Trifolium alexandrinum* cv. Giza 6, *Zea mays* cv. T. W.C-360, *N. rustica*, *Lycopersicon esculentum* cv. Casle Rock, *Solanum melongena* cv. Baladi, *Capsicum annuum* cv. California Wonder, *Solanum tuberosum* cv. sponta, *Brassica oleracea* var. Capitata cv. Baladi, *Brassica oleracea* L. var. botrytis cv. soltany, *Brassica oleracea* var. italic, *Brassica Rapa* cv. Baladi and *Eruca sativa* cv. Baladi.




Fig (1): Symptoms of PSbMV as a result of artificial inoculation on some hosts (A,B) *V. unguiculata* systemic mosaic ; (C and D) *N. tabacum* cv **Xanthi** systemic mosaic; (E ,F,G and H) *N. tabacum* deformed leaves, (I,J, K and L) systemic mosaic ; (M and N) *N. tabacum* cv. **Samsun** deformed leaves ; (O) *C. quinoa* **L** local lesion ; (P) *C. amaranticolor* **Coste & Reyn** local lesion.

Biological and serological studies on *Pea seed-borne mosaic virus* (PSbMV) on cowpea forage crop (*Vigna unguiculata* L. Walp)

Table (1) Symptoms of *Pea seed-borne mosaic potyvirus* in different host as a result of artificial inoculation

Scientific name of Tested plant	English name	Variety	Observed symptoms	ELISA
Alliaceae				
<i>Allium cepa</i> L. cv. Giza 6	Onion	Giza 6	NS	-
<i>Allium sativum</i> L. cv. Baladi	Garlic	Baladi	NS	-
Chenopodiaceae				
<i>Beta vulgaris</i> L. cv. Pleno.	beet	Pleno	NS	-
<i>Chenopodium quinoa</i> Wild	quinoa		LL	+
<i>C. amaranticolor</i> Coste & reyne	large lambsquarters	Coste & reyne	LL	+
Cucurbitaceae				
<i>Cucumis sativus</i> L. cv. Beta Alfa	Cucumber	Beta Alfa	NS	-
<i>Arum palaestinum</i> L. cv. Baladi	Arum	Baladi	NS	-
<i>Cucurbita moschata</i> cv. winter lecsery	Pumpkin	winter lecsery	NS	-
<i>Citrullus vulgaris</i> L. cv. Giza 1	Watermelon	Giza 1	NS	-
<i>Cucumis melo</i> cv. Reticulatus	Melon	Reticulatus	NS	-
<i>Cucurbita pepo</i> L. cv. Escandrani	Squash	Escandrani	NS	-
Fabaceae				
<i>Vigna unguiculata</i> L. Cv. beef	Cowpea	beef	SM	+
<i>Glycine max</i> L. cv Giza 35	Soybean	Giza 35	NS	-
<i>Glycine max</i> L. cv. Crowford	Soybean	Crowford	NS	+
<i>Glycine max</i> L. cv. clark	Soybean	clark	NS	-
<i>Phaseolus vulgaris</i> L. cv. Bronco	Bean	Bronco	NS	+
<i>Pisum sativum</i> L. cv. Litel Marvel	Pea	Litel Marvel	SM	+
<i>Vicia faba</i> L. cv. Giza 3	Broad bean	Giza 3	SM	+
<i>Cicer arietinum</i> .cv. Giza 531	chickpea	Giza 531	SM	+
<i>Lens culinaris</i> cv. Kordyl.	lentil	Kordyl	SM	+
<i>Trifolium alexandrinum</i> L. cv. Giza 6	Berseem clover	Giza 6	NS	-
<i>Arachis hypogea</i> L. cv. Giza 5	Peanut	Giza 5	NS	+
Graminaceae				
<i>Hordeum vulgare</i> L. cv. G123	Barley	G123	NS	+
<i>Triticum aestivum</i> cv. sods1	Wheat	sods1	NS	+
<i>Zea mays</i> cv. T.W.C-360	Maize	T.W.C-360	NS	-
Solanaceae				
<i>Datura innoxia</i> L.	Sacred datura		NLL	+
<i>Nicotiana glauca</i> L.	wild tobacco		NLL	+
<i>N. rustica</i> L.	wild		NS	-
<i>N. tabacum</i> L. White Burley	tobacco	White Burley	SMD	+
<i>N. tabacum</i> L. Samsun	tobacco	Samsun	SMo	+
<i>N. tabacum</i> cv Xanthi	tobacco	Xanthi	SM	-
<i>Lycopersicon esculentum</i> L. cv. Casle Rock	Tomato	Casle Rock	NS	-
<i>Solanum melongena</i> L. cv. Balady	eggplant	Balady	NS	-
<i>Capsicum annuum</i> L. cv. California Wonder	pepper	California Wonder	NS	-
<i>Solanum tuberosum</i> L. cv. sponta	potato	sponta	NS	-
Brassicaceae				
<i>Brassica oleracea</i> var. Capitata cv. Baladi	cabbage	Baladi	NS	-
<i>Brassica oleracea</i> L. var. botrytis cv. soltany	Cauliflower	soltany	NS	-
<i>Brassica oleracea</i> var. italica	Broccoli	italica	NS	-
<i>Brassica Rapa</i> cv. Baladi.	Turnip	Baladi	NS	-
<i>Eruca sativa</i> cv. Baladi	Arugula	Baladi	NS	-

SM: Systemic mosaic, **SMo:** Systemic mottling., **SMD:** systemic mosaic and deformed leaves., **NLL:** Necrotic local lesions, **LL:** local lesions, + : **Positive**, - : **Negative** NS : NoSymptoms

2.2. Seed transmission through cowpea (*V. unguiculata* (L.) Walp):

Results showed that, PSbMV could be transmitted through cowpea (*V. unguiculata* (L.) Walp) seeds of local cultivar (buffalo). It was noticed that the percentage of transmission of PSbMV was reached 6%.

3- Serological detection:

The virus isolate was detected by indirect ELISA. Positive reaction was obtained only between the virus and its corresponding antiserum but not reacted with the other antisera reported earlier.

II- Virus purification and ELISA reagent production:

1- Purification of the virus isolate:

In the present work PSbMV was partially purified for antiserum production following the method mentioned before.

1.1. UV-absorption spectrum of the purified virus:

The absorption spectrum of the purified isolate was typical for nucleoprotein, with a maximum at 260 nm and a minimum at 245 nm. The ratios of A_{260}/A_{280} , A_{280}/A_{260} and A_{max}/A_{min} ratios were 1.16, 0.85, 1.04, respectively. Yield of the partially purified virus was 1.7 mg/100g of infected leaves.

1-2-Electron microscopy:

Electron micrographs of the partially purified virus preparation obtained from infected plants revealed the presence of numerous unaggregated filamentous virus particles (Fig. 2) of about 700-750nm long.




Fig (2): EM micrograph of partially purified virus particles of BSbMV, negatively stained with 2% uranyl acetate, pH 7.0 .(60.000 x)

2. Production of polyclonal antibodies produced against to BSbMV:

2.1.Determination of antiserum titer:

Antiserum developed against BSbMV after rabbit immunization from bleeding taken 3 times at weekly intervals after the last injection had antibody dilution titer of 1/800 in the first, rose to 1/1600 in the second bleeding (Table 3), and then to 1/3200 in the third bleeding, respectively, when infected sap was diluted to 1/10 was used in indirect

ELISA. Antiserum obtained after the 3rd bleeding (1\3200) was used in subsequent experiments.

2.2. Purification of IgG:

Purification of IgG was done using caprylic acid method as described before. Its concentration was adjusted to 1mg/ml (A₂₈₀= 1.4).

2.3. Conjugation of IgG with alkaline phosphatase:

Positive reactions obtained when purified IgG and IgG conjugated with alkaline phosphatase were tested for PSbMV- infected plants using DAS-ELISA. The optimum concentrations of IgG and IgG conjugate were 1.0 µg/ml and 1/1000, respectively according to the schematic diagram of checkerboard arrangement test (Table 3).

2.4. Determination of antigen dilution end point:

It was found that, IgG and IgG conjugated with alkaline phosphatase can be readily applied for virus detection in tobacco plants extracts at dilutions up to 1:500 (Table 4). Reading after 30 min incubation with the substrate.

2.5. Serological detection of PSbMV by DAC ELISA and DBIA using the ELISA reagents produced:

2.5.1. DAS –ELISA:

Results in Table (5) showed that, PSbMV could be transmitted through peaseeds cv. master B, lentil cv. Kordy 1 and peanut cv. Giza 5 with percentage of transmission 7%, 6% and 4% respectively. It was noticed that PSbMV was not transmitted through seeds of chickpea cv. Giza 1, and faba bean cv. sakha 1.

2.5.2. Dot- Blotting Immunobinding Assay on nitrocellulose membrane (DBIA).

PSbMV was readily detected immunologically using DBIA (Fig.3). Positive reaction was indicated by development of purplish- blue colour, whereas in negative reaction, tissues from healthy plants remain green.




Fig (3): Dot blot immunoassay (DBIA) using poly polyclonal antibodies. (1) Healthy and infected cowpea cv. Buffalo plant samples of PSbMV and (2) *N. tabacum*, cv White Burley (3) *N. tabacum* L Samsun L. Purplish blue color indicates positive reaction

Table (2): Determination of PSbMV antiserum titer in relation to time of blood collection

Antiserum dilution	ELISA reading of PSbMV					
	Antisera collected at weekly intervals (A405nm)					
	1 st week		2 nd week		3 rd week	
	Inf.	H.	Inf.	H.	Inf.	H.
1/100	<u>1.123</u>	<u>0.555</u>	0.873	0.365	0.892	0.350
1/200	0.892	0.517	<u>0.726</u>	<u>0.225</u>	0.728	0.337
1/400	0.756	0.371	0.603	0.262	0.730	0.261
1/800	0.674	0.362	0.580	0.256	0.669	0.230
1/1600	0.418	0.284	0.488	0.221	<u>0.573</u>	<u>0.205</u>
1/3200	0.390	0.216	0.297	0.232	0.424	0.212
1/6400	0.311	0.181	0.202	0.194	0.286	0.208

Reading after 30 min incubation with the substrate.

Inf. = Infected plants H. = Healthy plants

Table (3): Schematic diagram of checkerboard arrangement determination of approximate working dilutions of IgG and IgG conjugate to BSbMV for ELISA test.

Dilution of IgG conjugate	Concentration of IgG (µg/ml)							
	4.0		2.0		1.0		0.5	
	I	H	I	H	I	H	I	H
1/250	0.892	0.350	0.731	0.321	0.634	0.290	0.554	0.280
1/500	0.851	0.341	<u>0.672</u>	<u>0.290</u>	0.623	0.272	0.509	0.256
1/1000	<u>0.697</u>	<u>0.292</u>	0.604	0.259	0.587	0.232	0.461	0.236
1/2000	0.442	0.242	0.401	0.221	0.399	0.211	0.300	0.193

I = Infected plants ; H= Healthy plants

Table (4): Determination of antigen end point:

Antigen dilution	Absorbance at (405 nm)	
	ELISA reading	
	I	H
1/250	1.099	0.266
1/500	<u>0.640</u>	<u>0.169</u>
1/1000	0.408	0.255
1/2000	0.260	0.151

Table (5): The use of the produced ELISA kit in detection of PSbMV
In seed lots of different hosts .

Seedlings of	scientific name	No.infected seedlings/No.tested seedlings	% seed transmission
Chickpea	<i>Cicer arietinum</i>	0/100	0%
Lentil	<i>Lens culinaris</i>	6/100	6 %
Pea	<i>Pisum sativum</i>	7/100	7 %
Faba bean	<i>Vicia faba</i>	0/100	0%
Peanut	<i>Arachis hypogea</i>	4/100	4%

Part III- Cytological studies using electron microscopy:

Electron microscopy of ultra-thin sections prepared from infected tobacco leaf tissues revealed that almost all the inspected tissues were drastically affected with infection. The nucleus was misshapen, the nuclear membrane is irregular, and the plasma membrane is characterized with multiple invaginations (Fig.5). On the other hand, the aggregates of virus particles (VP) are found in the cytoplasm next to the chloroplast (Fig.6). The starch granules assumed to be large in the affected plastids. The thylakoids as well as intergrana lamella appeared as if not well fixed and showed different pathological anatomical symptoms (Fig.6). The phloem of the infected plants was also affected. The parenchymatous of the vessel cylinder showed unusual vesicle like structure. The membrane of the affected cells showed different degrees of degeneration.

The content of the companion cell assumed as if not fixed well (Fig4). Concerning leaf cell of cowpea

seedlings, the changes were disorganized chloroplast and its content, the envelop of plastid is not observed and the thylakoids exhibited signs of disintegration (Fig.11).

The mitochondria (M) were aggregated and degenerated, also abnormal structures in the cells were observed (Fig.9, 10) These cytological changes as described above were not observed in the cells of healthy control plants (Figs 8 and 12).

Part IV- Virus control:

Two different compounds were tested for their ability to make induce resistance against PSbMV in local lesion formed on *C. amanticolor*. Whereas camphor oil was used as antiviral agent Results demonstrated in Table (6) reveal that all these compounds induced resistance to virus

Infection, when applied to the plant as spray before virus inoculation




Fig (4): EM micrograph of vascular cylinder of tobacco leaf artificially infected with PSbMV. The parenchymatous of the vessel cylinder showed unusual vesicle like structure. The membrane of the affected cells showed different degrees of degeneration (5000X). **S**sieve tube, **V**vessel.




Fig (5): EM micrograph of ultrathin section of tobacco leaf artificially infected with PSbMV show misshaped uncles (N), the nuclear membrane (NM) is irregular and the plasma membrane also affected in the central cell compared with the right cell in the current figure (15000X). **TP**:tonoplast, **N**:nuclear, **NM**:nuclear membrane, **CW**: cell wall, **PL**:plasma membrane, **M**: mitochondria








Fig (6): EM micrograph of ultrathin section of tobacco leaf artificially infected with PSbMV. The aggregates of virus particles (VP) are found in the cytoplasm next to the chloroplast. The starch granules (SGs) assumed to be large in the affected plastids. the thylakoids as well as intergrana lamella (L) appered as if not well fixed and showed different pathological anatomical symptoms. (12000X).

	
<p>Fig (7): Magnified part of Fig .(6) showing several particles of rod shape of PSbMV(50000X)</p>	<p>Fig (8): EM micrograph of ultrathin section in healthy cell of tobacco leaf (30000X). TP: tonoplast, CW: cell wall, PL: plasma membrane, G: grana</p>
	
<p>Fig.(9): EM micrograph of ultrathin section on seedling of cowpea plant(7days after germination), growing from seeds which harvested from artificially inoculated plants with PSbMV. Note Pinwheel inclusions (Pw) in the cells .(2500X)</p>	<p>Fig:(10): EM micrograph of ultrathin section on seedling of cowpea plant 7days after germination showing natural structure of mesophyll cells in healthy cowpea leaf (2500X). Ch:chloroplast , Nu: nucleus , M: mitochondria</p>




Fig (11): EM micrograph of ultrathin section in leaf cells of cowpea plant (7days after germination), growing from seed which harvested from artificially inoculated plants with PSbMV. Note the disorganized plastid and its content, also the envelop of plastid is not observed and the thylakoids expiated signs of disintegration. (25000X).

Fig (12): EM micrograph of ultrathin section on healthy seedling of cowpea plant (7days after germination) showing natural structure of plastid in cowpea leaf (30000X).Ch:chloroplast, L: intergrana lamella.

Table (6): Effect of salicylic acid (SA) and parahydroxy benzoic acid (PHBA) and camphor oil (CO) on PSbMV infection expressed as average number of local lesion.

Chemical Concentratio n %	Percentage of inhibition after treatment with								
	Salicylic acid (SA) after			Parahydroxy benzoic (PHBA) acid after			Camphor oil (CO) after		
	1h	2h	3h	1h	2h	3h	1h	2h	3h
0.1	46.4	49.2	50.7	63.3	74.6	63.3	47.8	40.8	63.3
0.01	16.9	46.4	77.4	42.2	74.6	52.1	57.7	66.1	71.8
0.001	60.5	35.2	50.1	74.6	88.7	69.0	29.5	74.6	73.2

Average number of local lesions induced in 15 leaves with SA, PHBA and CO

Control = unsprayed Plant

88.7% reductions before 2h of inoculation than 1 h (74.6%). Also, salicylic acid was effective as it reduced infection by 77.4% when used 0.01% concentration before

3hours of infection . Moreover camphor oil gave a little like reduction as salicylic acid (74.6) when used with 0.001% concentration before 2 h.

Fortunately, all compounds were effective to inhibit PSbMV.

DISCUSSION

In this work *Pea seed borne mosaic Potyvirus* was isolated from naturally infected cowpea plants cultivated in A.R.E.S. It was identified on the bases of symptomology, host range, and serology and particles morphology. Symptoms vary depending on virus or strain, host plant, time of year and environmental conditions (Matthews,1991). The virus was found to infect several hosts resemble those produced by some other PSbMV isolates especially *Chenopodiaceae* and *Fabaceae*. These agree with the results obtained by several workers (Hampton and Mink,1975,Alconero *et al.*, 1986 and Brunt *et al.*, 1996).It was noticed that some hosts did not produced any symptoms but when checked by ELISA gave positive reactions. Apola *et al.* (1974)mentioned that non-leguminous host are infected without producing symptoms. This might be attributed to the low concentration of the virus in these plant species. Some differences in host reaction were also observed, so these hosts can be used to differentiate between our PSbM isolate (cowpea isolate) and other PSbMV isolates.

Knowledge of the ways in which viruses are transmitted from plant to plant is important for recognized particular viral disease and

development of satisfactory control measures (Matthews,1991). The virus under study was transmitted through the seeds of cowpea, lentil, pea and peanut at 6,6,7 and 4%, respectively, but not through chickpea and faba bean. Khatarpal and Maury(1987) found that PSbMV was transmitted in the lentils at frequencies of 32-44% and through a low percentage of seeds of faba bean. Whereas Shukla *et al.* (1994) reported seed transmission of the virus in lentil at range of 0.2 - 44% and in pea at range of 0.3-80%.

The use of ELISA technique greatly facilitated the identification of the virus and provide more accurate and consistent results than did symptomatology and host range (Matthew,1992). Positive reaction obtained with the virus and its corresponding antiserum using indirect ELISA provided further evidence that the virus under study is indeed PSbMV.

In the present investigation PSbMV was partially purified for antiserum production following the method described earlier. The estimated yield of the purified virus was 1.7 mg/100g of infected leaves. The yield was lower than that of the other Potyvirus (Awad,Maisa *et al.*,2005, Farag,,Azza *et al.*2005 and El-Kady *et al.*,2010). It should be pointed out that a technique which succeeds with one virus isolate may fail with other one because of such factors as host species, growing conditions and

isolation procedure (Matsubara, 1985).

Electron micrograph of the partially purified virus preparation should numerous unaggregated filamentous flexuous virus particles of about 700-750 nm long. Such length is within the range reported for PSbMV by other investigators (Alconero *et al.* 1986, Franki *et al.* 1991 and El-banna *et al.*, 2008). This type of virus particles is characteristic for Potyvirus (Bos, 1970)

In the present study polyclonal antibody raised against PSbMV was prepared after rabbit immunization from bleeding taken to 3 weeks after the last injection using indirect ELISA. The titer of the antisera was 1/800 for the 1st, 1/1600 for the 2nd and 1/3200 for the 3rd bleeding. The latter is higher than that reported by (Brunt *et al.* 1996) for PSbMV. The concentration of Ig G and IgG conjugate were 1.0 mg/ml and 1/1000, respectively. These results agree with the results obtained by Salama (1998) and EL-kady *et al.* (2010).

Our study showed that the procedure of DAS-ELISA and DBIA could be readily applied in detection of PSbMV in the either infected plants or infected seeds for routine indexing of large number of samples.

Several investigators used both techniques for detection of their isolated viruses (Salama, 1998,

Awad, Maisa *et al.* 2005 and El-kady *et al.*, 2010)

External symptoms are reflection of disturbed cell metabolism leading to modifications in tissues, cell and cell organelles (Franck, *et al.*, 1985). Ultrathin sections of infected cells showed several cytological changes, these include chloroplast which were disorganized and fragmented, grana and intergrana were not fixed well and degenerated. The mitochondria were aggregated and degenerated. Intracellular structures such as inclusion bodies or pinwheels were observed. Besides these cytological abnormalities, aggregates of virus particles are also found in the cytoplasm next to the chloroplast. PSbMV has been reported by different works to induce cylindrical cytoplasmic inclusion (CCI). CCI seen in transverse section are described as pinwheels and their presence in diseased plant is of diagnostic value as they are characteristic of Potyvirus group (Edwards *et al.* 1984, and Shukla *et al.* 1994).

Important advances in virus chemotherapy have been made during the last few years. A variety of compounds with potent and selective antiviral activity had been found. These antiviral agents affected viral synthesis. Other compounds also result in increase resistance of the treated areas to some viruses (Gupta *et al.*, 1980, and Sherwood, 1985). In this work salicylic acid (SA), parahydroxy

benzoic acid (BHBA) and camphor oil (CO) were used for reduce PSbMV infection. The three compounds were found effective in this respect. Nevertheless, application gave suppression results (88.7 %) followed by CO (74.6%) when applied 2h before inoculation, whereas SA had the lower effect (60.5%) at concentration of 0.001 before 1h of inoculation. Several investigators used SA and PHBA for their ability to stimulate immune system and offer some indirect protection against viral infection through these effects (Hooir *et al.* 1986, Kobeasy and Salwa, Zein, 2005 and El-Kady *et al.* 2010).

On the other hand essential oils such as camphor oil were mainly used to control plant pathogens viz, fungal, bacteria and viral pathogens and act as antiviral agents (Verma *et al.*, 1998, Romeilah *et al.*, 2010 and Modhusudhan *et al.*, 2011).

REFERENCES

- Aapola, A.A., Knesek, J.E., Mink, G.I. (1974). The influence of inoculation procedures on the host range of pea seed borne mosaic virus. *Phytopathology* 64: 1003-1006.
- Ahmed, Amal A., Salwa N. Zein and Eman A.H. Khatab (2010). Characterization of Celery mosaic virus isolated from some apiaceae plants. *Egyptian J. Virol.* 7: 121-136.
- Alconero, R., weeden, N.f., Gonsalves, D. and Fox, D.T. (1985) loss of genetic diversity in the pea germplasm by the elimination of individuals infected by pea seed-borne mosaic virus. *Ann. Appl. Biol.* 106: 357- 364.
- Awad, Maisa A.E., EL-Banna, Om-Hashim, M., El-Kady, M.A.S., Hassan, H.M.S. and Abdalla, Ashgan A.M. (2005). purification and production of ELISA reagents against watermelon mosaic virus. *Egyptian J. virology* 2(1):101-111.
- Azzam, Clara R., Salwa N. Zein, Salwa N. and Abbas, Salwa M. (2007). Biochemical genetic markers for levels of resistance to Cowpea Aphid Borne Mosaic Potyvirus (CABMV) in sesame (*Sesamum indicum* L.) irradiated with gamma ray. *Egypt. J. Plant Breed.* 11 (2): 861-885.
- Bos, L. (1970). The identification of three new viruses isolated from *Wisteria* and *Pisum* in the Netherlands and the problem of variation within the potato virus Y group. *Neth. J. path.* 76:8-46.
- Bozzola, J.J. & Russell, L.D. (1999) Chapter 2, Specimen preparation. *Electron Microscopy: Principles and Techniques for Biologists*, 2nd

- edition. Sudbury, Massachusetts, Jones & Bartlett. pp. 14–37
- Brunt, A.A., Crabtree, K., Dallwitz, M.J., Gibb, A.J. and Waston, L. (1996).** Viruses of plants. 2nd CAB International Wallkingford. U.K., 148pp.
- Clark, M. F. and Adams, A. N. (1977).** Characteristics of the microplate method of enzyme-linked immunosorbent assay for detection of plant viruses. *J. Gen. Virol.* 34: 475-483.
- Converse, R.H., and Martin, R.R. (1990).** ELISA methods for plant viruses. In: Serological methods for detection and identification of viral and bacterial plant pathogens. pp. 179-196. Hompton, R. O.; Ball, E. M.; and DeBoer, S.H. (eds.). The American Phytopathological Society, St. Paul, Minnesota, USA.
- Devi, P.R., Doraiswamy, S., Nakkeeran, S., Rabindran, R.R., Ganapathy, T., Ramiah M. and Mathiyazhagan, S. (2004).** Antiviral action of *Harpulia Cupanioides* and *Mirabilis jalapa* against Tomato spotted wilt virus (TSWV) infecting tomato. *Archives of Phtopathology and Plant Protecion*, 37 (4): 245-259.
- Edwardson, J. R., Christie, R. G. and Ko, N. J. (1984).** *Potyvirus* cylindrical inclusions subdivision–IV. *Phytopathology* 74:1111- 1114.
- EL-Banna Om-Hashem M., Ebrahem, A.H., Kattab Eman A.H and El-Kammar Hanan (2008).** Isolation, identification, purification and production of ELISA Reagent for pea seed-borne mosaic potyvirus affecting *Pisum sativum* L., plants Egypt. *Egyptian j. Virol.* 5 (2), 157-172.
- El- Kady, M.A.S., Badr, A.B.B., Salwa N. Zein and Khalifa, M.A.A. (2010).** Antiserum production and control measures of *Pepper Mottle Potyvirus* using different applications. *Egyptian J. Virol.* 7: 105-120.
- Farag, Azza, G., M.A.S. El-Kady, El-Banna, Om-Hashem ,M.E., Hassan, H.M.S. and Abdullah, Ashgan A.M. (2005)** Biological and molecular characterization of *Watermelon mosaic potyvirus* (WMV). *Egyptian J. virol.* 2(1) : 77-99.
- Francki, R. I. B. Milne, R. G. and Hatta, T. (1985).** Atlas of plant viruses, Vols.1, 2 CRC press, Boca Roton, Florida (c. f. Matthews *et al.*, 1991).
- Francki, R. I. B., Fauquet, C.M., Knudson, D. L., and Brown, F. (1991).** Classification and Nomenclature of Viruses: Fifth Report of the International Committee on Taxonomy of Viruses. Sprinder, Wine and

- New York Archives of Virology (Supplement2).
- Gamal El-Din A.S., El-Kady M.A.S., Shafie M.S.A and Abo-Zeid A.A. (1997)** Tuber necrotic ringspot strain of potato virus Y (PVY^{NTN}) in Egypt 8th congress of Egyptian Phytopathol. Soc., Cairo, Egypt.
- Gillaspie, A.G., Jr, H.M.S., Pinnow, D.L., and Hampton, R.O. (1995)**. Seedborne viruses in preintroduction cowpea seed lot and establishment of virus-free accession. *Plant Dis.* 79: 388-391.
- Gupta, M.D., Rao, R. and Verma, V.S. (1980)**. Inhibition of mosaic virus of *vignasinensis* with four chemicals. *Acta. Microbiol. Pol.* 2a: 65-68.
- Hampton, R.O., and Mink, G.I. (1975)**. Pea Seed-borne mosaic virus CMI\ AAB Discription of plant viruses. N^o 146.
- Hampton, R., Ball, E. and Beboer, S. (1990)**. Serological Methods for Detection and Identification of Viral and Bacterial Plant Pathogens. The American Phytopathological Society St. Paul, Minnesota, USA, 389pp.
- Hoor, R.A.M., Vanhijdsuijnen, S.W., Alblas, R.H. and Bol, J.F. (1986)**. Induction by salicylic acid of pathogenesis-related protein and resistance to alfalfa mosaic virus infection in various plants species. *J. Gen. Virol.* 67:2135-2143.
- Hsu, H. T., and Lawson, R.H. (1991)**. Direct tissue blotting for detection of tomato spotted wilt virus in impatiens. *Plant Dis.* 75: 292-295.
- Khetaral, K.R and Maury, Y. (1987)**. Pea seed – borne mosaic virus: a review. *Agronomie.* 7(4): 215-224.
- Kobeasy, M. I and Salwa N. Zein, Salwa N. (2005)**. Role of salicylic and parahydroxybenzoic acids as resistance inducer for *Barley stripe Mosaic virus* infection in barley plants. *J. Agric. Sci. Mansoura Univ.*, 30 (8): 4929 – 4943.
- Kuhn, C.W. (1964)**. Separation of cowpea virus mixture. *Phytopathology* 54: 739-740.
- Lommel, S. A., McCain, A. H., and Morris, J. T. (1982)**. Evaluation of indirect enzyme-linked immunosorbent assay for the detection of plant viruses. *Phytopathology* 72:1018-1022.
- Makkouk, K.M., Bos, L., Azzam, O.L., Kumari, S. and Rizkallah, L.R. (1988)**. Survey of viruses affecting faba bean in six Arab countries. *Arab Journal of Plant Protection* 6:53-61.

- Matsubara, A.** Kawno, S. Marita, M. Hattori, M., Uyeda, and Shikata, E. (1985). Purification and serology of Japanese isolate of barley dwarf virus. *Ann. phytopatho. Soc. Jpn.* 51:152-158.
- Matthews, R.E.F. (1991).** *Plant Virology*, 3th ed. Academic Press, London, 538pp.
- Matthews, R.E.F. (1992).** *Fundamental of Plant Virology*. Academic Press, Inc. 403pp.
- Modhusuhan, k.N.,** Vinavarani, G., Deepak, S.A., Niraniana, S.R., Prakash, H.S., Singh, G.P., Sinha, A.K. and Parsad, B.C. (2011). Antiviral activity of plant extracts and other inducers against Tobacco mosaic virus infection in bell pepper and tomato plants. *International Journal of plant pathology* 2: 35-42.
- Noordam, D. (1973).** Identification of Plant Viruses. Methods and Experiments. Center for Agric. Publishing and Documentation, Wageningen, Netherlands, 207 pp.
- Purcifull, D.E. (1990).** *Tobacco etch Potyvirus*. (537-540). In: Viruses of Tropical plants A. Brunt, K. Crabtree and A. Gibbs (eds.) 537-540pp. C.A.B. International. Wallingford.
- Romelah, R.M.,** Fadyed, S.A. and Mahmoud, G.I. (2010) Chemical composition, antiviral and antioxidant activity of seven essential oils. *Journal of Applied Science Research* 6(1):50-62.
- Salama, M.Im. (1998).** Molecular and serological studies of some faba bean (*Vicia faba* L.) viruses: Ph.D. thesis, fac. Agric., Ain-Shams Univ., Cairo. 94pp.
- Sherwood, J.L. (1985).** The association of (pathogenesis-related) proteins with viral-induced necrosis in *Nicotiana sylvestris*. *Phytopathology zeitschrift* 112:48-55.
- Shukla, D.D., Ward, C.W. and Brunt, A.A. (1994).** The potyviridae. CAB international Wallingford, U.K., 516 pp.
- Smith-Backer, J., Marois, E., Huguet, E. J., Midland, J. J. S. and Keen, N. T. (1998).** Accumulation of salicylic acid and 4-hydroxybenzoic acid in phloem fluids of cucumber during systemic acquired resistance is preceded by a transient increase in phenylalanine ammonia-lyase activity in petioles and stems. *Plant Physiol.* 116(1): 231 – 238.
- Stangarlin, J.R.,** Schwan-Estada, K.R.F., Cruz, M.E.S. and Nozaki, M.H. (1991). Medicinal plants and alternative control of phytopathogens. *Biorecologia ciencia & Desenvolvimento* 11:16-21.

- Steinbuch, M. and Audran, R. (1969).** The isolation of IgG from mammalian sera with the aid of caprylic acid. *Archives of Biochemistry and Biophysics* 134:279-284.
- Taiwo, M.A., Kareem, K.T., Nsa, I.Y., Hughes, J.A. (2007).** Cowpea viruses: Effect of single and mixed infections on symptomatology and virus concentration. *Virology Journal* 4:95.
- Verma, H.N., Baranwal, V.K. and Srivastava, S. (1998)** Antiviral substances of plant origin. In: *Plant virus disease control*, Hadidi, A., R.K. Khetarpal and Koganezawa (Eds.). APS Press, St. Paul, Minnesota: 154-162.
- White, R.F., Dumas, E., Show, P., and Antoniow, J.F. (1986).** The chemical induction of PR (b) proteins and resistance to TMV infection in tobacco. *Antiviral Res.*, 6: 177-185.
- Yalpani, N., Sliverman, P., Wilson, T. M. A., Kleier, D. A. and Raskin, I. (1991).** Salicylic acid is a systemic signal and an inducer of pathogenesis-related proteins in virus-infected tobacco. *Plant Cell*. 3: 809 – 818.