

REFERENCES

Abdel-Rahman, M. A. (1985):

Studies on the antigenicity and toxigenicity of *Clostridium welchii*
Ph. D. Thesis (Microbiology), Faculty of Veterinary Medicine, Cairo
University.

Abdel-Rahman, A. A.; Moustafa, F. A. and Hamd, N. A. (2006):

Detection of the prevalence and pathogenicity of *Clostridium perfringens*
and *Clostridium spiroforme* associated diarrhoea in rabbits.
Assiut Veterinary Medical Journal, 52(108):321-335.

Abd El-Salam, I. S. and El-Sanousi, S. M. (1992):

Proposed scheme for isolation and identification of *Clostridium*
perfringens and *Clostridium perfringens* – like organisms.
Review Elevista, 44(2):153-158.

Agnoletti, F.; Bonci, M.; Drigo, I. and Bano, L. (2010):

Antimicrobial susceptibility of *Clostridium perfringens* in breeding rabbit.
Summa Animalia da Reddito, 5(6): 43-48.

**Agnoletti, F.; Bacchin, C.; Bano, L.; Passera, A.; Favretti, M. and
Mazzolini, E. (2007):**

Antimicrobial susceptibility to zinc bacitracin of *Clostridium perfringens*
of rabbit origin
World Rabbit Science, 15:19-22.

Augustynowicz, E.; Gzyl, A. and Slusarczyk, J. (2000):

Molecular epidemiology survey of toxinogenic *C. perfringens* strain types
by multiplex PCR.

Scandinavian Journal of infectious Diseases, 32:637-641.

Awad, M. M.; Bryant, A. E.; Sterens, D. L. and Rood, J. I. (1995):

Virulence studies on chromosomal α -toxin and β -toxin mutants constructed by allelic exchange provide genetic evidence for the essential role of α -toxin in *Clostridium perfringens* – mediated gas gangrene.

Molecular Microbiology, 15:191-202.

Badagliacca, P.; Provvido, A.; Scattolini, S.; Pompei, G. and Giannatale, E. (2010):

Toxin genotyping of *Clostridium perfringens* strains using a polymerase chain reaction protocol.

Veterinaria Italiana, 46(1):107-112.

Banchroft, J. D.; Stevens, A. and Turner, D. R., (1996):

Theory and practice of histological techniques, .Fourth Ed. Churchill Livingstone, New York, London, San Francisco, Tokyo.

Bano, L.; Busani, L.; Cocchi, M.; Drigo, I.; Spigaglia, P.; Mastrantonio, P. and Agnoletti, F. (2008):

Prevalence and molecular characterization of *Clostridium difficile* isolated from rabbits and detection of its main toxins.

9th World Rabbit Congress, Verona, Italy.

Banwart, G. J. (1989):

Basic Food Microbiology, 2nd Ed.

Vannostrand, Reinhold, New York.

Barral, E.; Biet, F.; Duivon, D.; Filleul, J. P. and Prim, R. (2000):

Assessment of the extent of lesions, parasites and bacteria in rabbit farms contaminated by epizootic rabbit enterocolitis (ERE).

7th World Rabbit Congress.

Baskerville, M.; Wood, M. and Seamer, J.H. (1980):

Clostridium perfringens type E enterotoxaemia in rabbits.
Veterinary Record, 107(1):18-19.

**Borriello, S. P.; Fiona, E.; Barclay, A. R.; Welch, M. F.; Stringer, G. N.;
Watson, R. K. T.; Williams, D. V. S. and Kate, S. (1985):**

Epidemiology of diarrhea caused by enterotoxigenic *Clostridium Perfringens*.
Journal of Clinical Microbiology, 17(3):414-418.

Boyd, M. G., Logan, M. A. and Tytell, A. A. (1948):

The growth requirement of *Clostridium perfringens (welchii)*.
Journal of Biological Chemistry, 174:1013-1025.

Branton, S.L.; Reece, F. N. and Hagler, W. M. (1987):

Influence of a wheat diet on mortality of broiler chickens associated with necrotic enteritis.
Poultry Science, 66:1326-1330.

Brooks, D. L. (1979):

Endemic diarrhea of domestic rabbits in California.
Dissertation Abstracts International, 39(12): 5853.

Brynstad, S. and Granum, P. E. (2002):

Clostridium perfringens and food borne infection.
International Journal of Food Microbiology, 74:195-202.

Bullen, J. J. (1952):

Clostridium welchii type D in the alimentary tract of normal sheep.
Journal of Pathology and Bacteriology, 64: 201-210.

Buogo, C.; Capaul, S.; Hani, H.; Frey, J. and Nicolrt, J. (1995):

Diagnosis of *Clostridium perfringens* type C enteritis in pigs using a DNA amplification technique (PCR).

Journal of Veterinary Medicine (B), 42(1):51-58.

Campos, D. F. S.; Baccaro, M. R.; Ferreira, A. J. P.; Dotto, D. S.; Murakami, D. F.; Moreno, A. M.; Knobl, T. and Shinya, L. T. (2002):

Effects of *Clostridium perfringens* CPB-2 positive strains inoculation in rabbit ileal loops.

Arquivos do Instituto Biologico (Sao Paulo), 69:126-128.

Campos, D. F. S.; Baccaro, M. R.; Moreno, A. M.; Ferreira, A. J. P.; Doto, D. S.; Murakami, D. F.; Knobl, T. and Shinya, L. T. (2004):

Clostridium perfringens type-A strains carrying the CPB-2 gene associated to the lesions in rabbit ileum.

Arquivos do Instituto Biologico (Sao Paulo), 71(3):287-292.

Carter, G. R. and Cole, J. R. (1990):

"Diagnostic procedures"

Veterinary Bacteriology and Mycology, 5th Edition, Academic press, Harcourt, Boace. Jov. Publisher New York, Boston, Tokyo, Toronto.

Cato, E. P.; George, W. L. and Finegold, S. M. (1986):

"Genus *Clostridium*"

Bergey's manual of systemic bacteriology. Vol. 2. Sneath, P. H., Mair N.S., Sharpe, M.E. and Hott, J.G. (eds) Williams & Wilking Co., Baltimore pp. 1141-1200.

Chai, T.; Wang, L.; Wang, H.; Duan, H.; Muller, W. and Zucker, B. A. (2007):

Isolation and characterization of *Clostridium perfringens* from apparently healthy animals of the Shandong province of China.

Deutsche Tierärztliche Wochenschrift, 114(10): 394-396.

Cheecke, P.R., (1986):

Potentials of rabbit production in tropical and sub tropical agricultural systems.

Journal of Animal Science, 63(5):1581-1586.

Cruikshank, R.; Deguid, J. P.; Morromain, B. P. and Swaim, R. H. (1975):

Medical Microbiol. 12th ed. Vol. II, Churchill Livingstone. Edinberg, London and New York.

Diab, R. A.; El-Sehemy, M. M.; Nadia, M. E.; Fatheia Shafie and Hussein, A. Z. (2003):

Enterotoxaemia in rabbits and trials for preparing vaccine from the isolated strains.

Journal of Veterinary Medical Association, 63(2):59-64.

Downes, M. J.; Stern, L. and Andreq, J. H. (1986):

A comparison of selective media for the isolation of anaerobic bacteria from clinical material.

Pathology, 18(1):141-144.

Drigo I.; Bacchin C.; Cocchi M.; Bano L. and Agnoletti F. (2008):

PCR identification and toxin-typing of *Clostridium Spiroforme* field strains.

9th World Rabbit Congress -Verona -Italy.

El-Naenaey, E. Y. and El-Fetouh, M. S. A. (1993):

The efficacy of *Clostridium perfringens* enterotoxin in rabbits.

Veterinary Medical Journal (Giza), 41(3):17-22.

El-Rahman, M. A. and Atwa, E. I. (2006):

Studies on *Clostridial* microorganisms in rabbits and the use of ELISA for detection of *Clostridium perfringens* toxins.

Veterinary Medical Journal Giza, 54(3):671-684.

Fernandez-Miyakawa, M. E.; Marcellino, R. and Uzal, F. A. (2007):

Clostridium perfringens type A toxin production in 3 commonly used culture media.

Journal of Veterinary Diagnosis and Investigation, 19(2):184-186.

Ficken, M. D. and Wages, D. P. (1997):

Calnek (Ed.), Diseases of poultry 10th ed. P.261-264.

Ames, IA: Mosby-Wolfe.

Finzi, A. and Amici, A. (1991):

Traditional and alternative rabbit breeding systems for developing countries.

Rivista di Agricoltura Subtropicale e Tropicale, 6(1):103-125.

Frankel, E. (1893):

On the etiology of gas gangrene.

Zentblatt Pakistani, 13:13-16.

Genigeorgis, C. ; Sakaguchi, G. and Riemann, H. (1973):

Assay methods for *Clostridium perfringens* type A Enterotoxin.

Applied Microbiology, 26 (1): 111-115.

Geppert, P. and Eisgruber, H. (1995):

Assay of the phosphatase of *Clostridium perfringens*.

Archive Fur Leben Smitte Hygiene, 46(2):30-35.

Glenn Songer, J. and Ralph Meer, R. (1996):

Genotyping of *Clostridium perfringens* by polymerase chain reaction is a useful adjunct to diagnosis of *Clostridial* enteric disease in animals. *Anaerobe*, 2: 197-203.

Green Wood, P. E and Nickalin, M. S. (1996):

A guide to Chi-squared testing.
Wiley, New York.

Haghour, R. and Murad, M. (1984):

Laboratory examination of enterotoxaemic diseases in sheep in Syria. *Assuit Veterinary Medical Journal*, 11(22):171-174.

Han, S. Y.; Sang, U.; Kyoung, Y. P. and Yong, H. P. (1997):

Molecular typing and epidemiological survey of prevalence of *Clostridium perfringens* types by multiplex PCR. *Journal of Clinical Microbiology*, 28: 228-232.

Hara-Kudo, Y.; Morishita, Y.; Nagaoka, Y.; Kasuga, F. and Kumagai, S. (1996):

Incidence of diarrhea with antibiotics and the increase of *clostridia* in rabbits. *Journal of Veterinary Medical Science*. ;58(12):1181-5.

Harbola, P. C.; Kumar, S. and Datt, N. S. (1975):

A report on the laboratory examination of enterotoxaemic diseases in sheep and goats. *Indian Veterinary journal* 52(8):597-599.

Harris, I. E. and Portas, B. H. (1985):

Enterotoxaemia in rabbit caused by *Clostridium spiroforme*. *Australian Veterinary Journal*, 62(10):342-343.

Heba, M. D. (2010):

Microbiological studies on *Clostridium perfringens* affecting laboratory animals.

M.V.Sc. Thesis (Microbiology), Faculty of Veterinary Medicine, Cairo University.

Hein, H. and Timms, L. (1972):

Bacterial flora in the alimentary tract of chickens infected with *Eimeria brunetti* and in chickens immunized with *Eimeria maxima* and cross infected with *Eimeria brunette*.

Experimental Parasitology, 31: 188-193.

Hofshagen, M. and Stenwig, H. S. (1992):

Toxin production by *Clostridium perfringens* isolated from broiler chickens and capercaillies (*Tetrao urogallus*) with and without necrotizing enteritis.

Avian Diseases, 36:837-843.

Horadniceanu, T. and Sasarman, A. (1964):

Method of isolating *Clostridium perfringens*, Comparison of some media.

Archive Roum Pathology and Experimental Microbiology, 23:633-640.

Huybens, N.; Houeix, J.; Licois, D.; Mainil, J. and Marlier, D. (2011):

Epizootic rabbit enteropathy inoculum (TEC4): antibiograms and antibiotic fractionation.

Veterinary Research Communication, 35 (1):13-20.

Hughes, S.; Warhurst, G.; Turnberg, L. A.; Higgs, N. B.; Giugliano, L. G. and Drasar, B. S. (1983):

Clostridium difficile toxin-induced intestinal secretion in rabbit ileum *in vitro*.

Gut, 24: 94-98 doi:10.1136/gut.24.2.94.

Hunter, S. E.; Clarke, I. N.; Kelly, D. C. and Titball, R. W. (1992):

Cloning and nucleotide sequencing of the *Clostridium perfringens* Epsilon toxin gene and its expression in *E. coli*.

Infectious Immunology, 60:102-110.

Ivanics, E.; Glavits, R. and Hadhazy, A. (1982):

Occurrence of Tyzzer's disease in brown hares (*Lepus europaeus*).

Mayar Allatorvosok Lapja, 37(8):525-527.

Jiang XiangYu (2008):

Control of *Clostridium welchii* in rabbits.

Chinese Journal of Rabbit Farming, 8:10-11.

Joklik, W. K.; Willet, H. P. and Amos, D. B. (1980):

Zinsser Microbiology

17^{ed} Appletin, Century Crofts, New York.

Jong, A. E. I. (2002):

Optimizing sporulation of *Clostridium perfringens*.

Journal of Food Protection, 65(9): 1457-1462.

Jorge, E. V.; Bruce, A. M.; Juliann, S.; Jaquelyn P. and Francisco, A. U. (2008):

Effects of *Clostridium perfringens* Beta-Toxin on the rabbit small intestine and colon

Infection and Immunity, 76(10): 4396-4404.

Julian, I. Rood (1998):

Virulence genes of *Clostridium perfringens*.

Annual Review of Microbiology, 52:333-360.

Kadra, B.; Guillou, J. P.; Popoff, M. and Bourliox, P. (1999):

Typing of sheep clinical isolates and identification of enterotoxigenic *Clostridium perfringens* strains by classical methods and by polymerase chain reaction (PCR)

FEMS Immunology and Medical Microbiology, 24(3): 259-266.

Keel, M. K. and Songer, J. G. (2006):

The Comparative Pathology of *Clostridium difficile*-associated Disease.

Veterinary Pathology, 43: 225-240.

Koneman, E. W.; Allen, S. D.; Janda, W. M.; Schreckenberger, P. C. and Winn, W. C. (1992):

Color atlas and textbook of diagnostic microbiology.

J.B. Lippincott Company Philadelphia, Fourth Edition

Kunstyr, I.; Matthiesen, I. and Matthiesen, T. (1975):

Acute enteritis in rabbits.

Zeitschrift Versuch, 17(1):57-63.

Labbe, R. G. (2000):

Clostridium perfringens. In: Lund, Baied - Parker, T., Gould, G. (Eds.).

The microbiological safety and quality of food. Aspen Publishers, MD, USA, PP. 1110-1135.

Lee, W. K.; Fujisawa, T.; Kawamura, S.; Itoh, K. and Mitsuoka, T. (1991):

Isolation and identification of *Clostridia* from the intestine of laboratory animals.

Laboratory Animal; 25(1):9-15.

Lepas, F.; Cmndrt, P.; Rochambeaude, H. and Thebault, R. G. (1997):

The rabbit husbandry, health and production (new revised version).

FAO, Animal Production and Health Series, No. 21.

Lukefahr, S. D., Nwosu, C. V. and Mo, D. R. (1989):

Cholestrol-level of rabbit meat and trait relationships among growth, carcass and lean yield performances.

Journal of Animal Science, 67: 2009-2017.

Macfaddin, J. F. (2000):

Biochemical test for identification of medical bacteria.

3rd Ed. Lippin Cott. Willions, Washington, Philadelphia, USA.

Mackie, T. J. and McCartney, J. E. (1960):

Hand book of bacteriology, 10th edition, E. and S. living stone limited, Edinburg and London.

Mariano, E. F. ; Derek, J. F.; Rachael, P.; Sameera, S.; Vicki, A.; Julian,

I. R.; Bruce, A. M. and Francisco, A. U. (2007):

Both epsilon-toxin and beta-toxin are important for the lethal properties of *Clostridium perfringens* Type B isolates in the mouse intravenous injection model.

Infection and Immunity, 75(3):1443-1452

Markovic, L.; Asanin, R. and Dimitrijevic, B. (1992):

In-vitro effect of *Clostridium perfringens* enterotoxin on vero cells and effects *in-vivo* animals.

Acta Veterinaria (Beogard), 43(4): 191-198.

Marlier, D.; Dewrée , R.; Lassence , C.; Licois , D.; Mainil , J.; Coudert , P.;

Meulemans , L.; Ducatelle , R. and Vindevogel, H. (2006):

Infectious agents associated with epizootic rabbit enteropathy: Isolation and attempts to reproduce the syndrome
The Veterinary Journal, 172: 493-500.

Matthes, S. (1981):

Bacteriocin susceptibility of *C. perfringens* type A provisional typing system.
Applied Microbiology, 28(2):172-176.

McDonel, J. L. and Duncan, C. L. (1975):

Histopathological effects of *C. perfringens* enterotoxin in rabbit ileum.
Infectious Immunology, 12 (5):1214-1218.

McDonel, J. L. and Duncan, C. L. (1977):

Regional localization of activity of *C. perfringens* type A enterotoxin in rabbits.
Journal of Infectious Diseases, 136(5):661-666.

Mead, G. C. (1985):

Selective and differential media for *Clostridium perfringens*.
International Journal of food Microbiology, 2 (1-2): 89-98.

Mehmet, H.; Ozlem, O. and Necdet, D. (2009):

Pathological and ultrastructural studies on mucoid enteropathy in New Zealand rabbits.
Journal of Exotic Pet Medicine, 18(3): 224-228

Merck, (1998):

"*Clostridial* diseases"
Veterinary Manual, 7th Edition, Merck publishing Group. Rahway. NJ, USA, pp. 445-447.

Merck, (2006):

"Digestive system"
Veterinary Manual, 9th Edition, Merck publishing Group. Rahway. NJ,
USA, pp. 225-237.

Migula, W. (1900):

System of bacteria.
Jena, Fisher (German).

Mogollon, G. J. D.; Pena, B. N. E.; Jimeno, M. C. and Rodriguez, M. G. (1981):

Complex mucoid enteritis of rabbits. II. Causal interrelation between
aflatoxins and *Clostridium perfringens*.
Revista Instituto Colombiano Agropecuario. 16(2): 97-111.

Molloy, R. (1978):

Bacterial phospholipases.
J. Jeljaszewicz and T.Wadstrom (Ed.) Bacterial Toxins and cell
membranes, Academic press. London, P. 367-424.

Mostafa, E. M. (1992):

Studies on the incidence of *Clostridial* organisms in domestic rabbits.
M.V.Sc. Thesis (Microbiology), Faculty of Veterinary Medicine Zagazig
University.

Nagi, G. M.; Laila, A.; Ebeid, M. H. and El-Sagheer, M. (1988):

Afield study on role of *Clostridium perfringens* in rabbit's diarrhea
complex.
Veterinary Medical journal, 36(2): 221-230.

National Committee for Clinical Laboratory Standards, (1998):

Performance standards for antimicrobial disc susceptibility tests.
8th Ed. International Supplement. M-I00-S7, Vol. 18 No.1. National
Committee Clinical Laboratory Standards. Villanova, P A, USA.

Niilo, L. (1971):

Mechanism of action of the enteropathogenic factor of *Clostridium perfringens* type A.

Infectious Immunology, 3(1):100-106.

Niilo, L. (1988):

Hemolytic patterns for presumptive identification of *Clostridium perfringens* type C.

Veterinary Research, 122(2): 281.

Oakley, C. I. and Warrack G. H. (1953):

Routine typing of *Clostridium wechii*.

Journal of hygiene Cambridge, 51:102-107.

Osman, M. R.; Kamal, O. E.; El-Sehemy, M. M. and El-Helw, H. A. (2008):

Immunological and molecular studies on *C. perfringens* type A isolated from different animals and poultry.

Proceedings of the Middle East & North Africa (MENA) Region Animal Wealth Research Conference. Future of animal wealth, Cairo International Convention Center, Egypt, 21-30.

Peter, H. A.; Nicholas, S.; Elisabeth, S. and John, G. (1986):

Bergey's manual systemic bacteriology. Vol. 2, Williams and Wilkins.

Peeters, J. E.; Geeroms, R.; Carman; R. J. and Wilkins, T. D. (1986):

Significance of *Clostridium spiroforme* in the enteritis-complex of commercial rabbits.

Veterinary Microbiology, 12(1): 25-31

Petit, L.; Gilbert, M. and Popoff, M. R. (1999):

Clostridium perfringens: toxinotype and genotype.
Trends Microbiology, 7:104-110.

Prescott, J. F. (1977):

Tyzzler's disease in rabbit in Britain.
Veterinary record, 100(14):285-286.

Pritt, S.; Henderson, K. S. and Shek, W. R. (2010):

Evaluation of available diagnostic methods for *Clostridium piliforme* in laboratory rabbits (*Oryctolagus cuniculus*)
Laboratory Animals, 44:14-19.

Quinn, P. J.; Markey, B. K.; Carter, M. E.; Donnelly, W. J.; Leonard, F. C. and Maguire, D. (2002):

Veterinary microbiology and microbial disease.
2nd Ed., Blackwell Science, 84-96.

Rahman, M. (1978):

Free sponug *Clostridium welchii* in ordinary laboratory media and conditions.
Journal of Clinical Pathology, 31: 359-360.

Rahman, M. S.; Rahman, M. M. and Khan, M. A. H. N. A. (1999):

Clinicopathological observation of *Clostridium perfringens* toxins in experimental guinea pigs.
Indian Veterinary Journal, 76(6): 497-500.

Reddy, N. V., Mo, D. R. and Chen, C. P. (1977):

Comparative performance of rabbits and broilers.
Nutrition Reports International, 16(1): 133-137.

Roberts, R. S.; Guven, S. And Narrat, E. F. (1970):

Studies on *Clostridium oedematiens*.

Journal of Comparative Pathology, 80: 9-18.

Romero, C.; Nicodemus, N.; Chamorro, S.; Garcia-Ruiz, A. I.; Ibanez, M. A. and Blas, C. (2009 a):

Estimation of Epizootic Rabbit Enteropathy effect on weight loss by means of *Clostridium perfringens* concentration in soft faeces.

XXXIX Jornadas de Estudio, XIII Jornadas, sobre Produccion Animal, Zaragoza, Espana, 188-190.

Romero, C.; Nicodemus, N.; Garcia-Ruiz, A. I.; Ibanez, M. A. and Blas, J. C. (2009 b):

The use of soft faeces for the prediction of the caecal contents concentration of *Clostridium perfringens* in relation with epizootic rabbit enteropathy.

Spanish Journal of Agricultural Research, 7(4): 807-812.

Romero, C.; Nicodemus, N.; Garcia-Rebollar, P.; Garcia-Ruiz, A. I.; Ibanez, M. A. and Blas, J. C. (2009 c):

Dietary level of fiber and age at weaning affect the proliferation of *Clostridium perfringens* in the caecum, the incidence of Epizootic Rabbit Enteropathy and the performance of fattening rabbits.

Animal Feed Science and Technology, 153(1-2): 131-140.

Sambrook, J.; Fritsch, E. F. and Maniatis, T. (1989):

“Purification of closed circular DNA by equilibrium centrifugation in Cs-CL ethidium bromide gradients, Molecular cloning a laboratory manual”

New York, CSH Laboratory.

Sayed, S.; Uzal, F. A.; Fisher, D. J.; Saputo, J.; Vidal, J. E.; Chen, Y.; Gupta, P.; Rood, J. I. and McClane, B. A. (2008):

Beta toxin is essential for the intestinal virulence of *Clostridium perfringens* type C disease isolate CN3685 in a rabbit ileal loop model.
Molecular Microbiology, 67(1): 15-30.

Scharmann, W. and Wolff, D. (1985):

Occurrence and prevention of Tyzzer's disease in rabbit colony.
The contribution of laboratory animal science to the welfare of man and animal,
8th ICLAS/CALAS symposium, Vancouver, 53-57

Secasiu, V. and Pastarnac, N. (1993):

Chemosensitivity of *Clostridium welchii* (*Clostridium perfringens*) strains isolated from furbearing animals.
Revista Romana de Medicina Veterinara, 3(1): 63-69.

Sheedy, S. A.; Ingham, A. B; Ood, J. I. and Moore, R. J. (2004):

Highly conserved alpha-toxin sequences of avian isolates of *C. perfringens*.
Journal of Clinical Microbiology, 42(3):1345-1347.

Shi XiShan; Wang CunLian and Xu Tong (2008):

Diagnosis and treatment of *Clostridium welchii* in Rex rabbit.
Chinese Journal of Rabbit Farming, 3: 36-37.

Smith, L. D. S. (1955):

Introduction to the pathogenic anaerobes.
University of Chicago. Press, Chicago, Illinois.

Smith, L.D. (1975):

The pathogenic anaerobic bacteria.

2nd Ed., Charles C-Thomas publisher, U.S.A.

Smith, L. D. and Holdman, L. V.(1968):

The pathogenic anaerobic bacteria.

Charleshomas publisher, USA. Ist ed. P 201-255.

Songer, J.G. (1996):

Clostridial enteric diseases of domestic animals.

Clinical microbiology Review, 9:216-234.

Stern, M. and Batty, I. (1975):

Pathogenic *Clostridia*.

Buthogenic, London, Baston.

Sting, R. (2009):

Detection of beta 2 and major toxin genes by PCR in *Clostridium perfringens* field isolates of domestic animals suffering from enteritis or enterotoxaemia.

Berliner und Munchener Tierarztliche Wochenschrift, 122(9-10): 341-347.

Szemerédi, G.; Palfi, and Gaco, I. (1983):

Etiology of diarrhea in rabbits at weaning.

Magyar Allatrosok Lapja, 83(5): 280-283.

Timoney, J. F.; Gillespie, J. H.; Scott, F. W. and Bariough, J. E. (1988):

Hagan and Bruneries microbiology and infectious diseases of domestic animals.

8th ed., Cornell University Press, Ithaca, N.Y., 223-229.

Titball, R. W.; Naylor, C. E. and Basak, A. K. (1999):

The *Clostridium perfringens* Alpha toxin.

Anaerobe, 5 (2): 51-64.

Tong, Y. L. and Labbe, R. (2003):

Enterotoxigenicity and genetic relatedness of *C. perfringens* isolated from retail food in United States.

Applied and Environmental Microbiology, 69(3):1642-1646.

Tripathi, B. N.; Prihar, N. S. and Singh, K. P. (1992):

Pathoanatomical responses to *Clostridium perfringens* type C infections toxin in ligated jejunal loops in guinea pigs.

India Journal of Animal Science, 62(8): 711-714.

Tzika, E. D. and Saoulidis, K. (2004):

Rabbit enteritis.

Journal of the Hellenic Veterinary Medical Society, 55(2):145-155.

Uzal, F.A. and McClane, B.A. (2011):

Recent progress in understanding the pathogenesis of *Clostridium perfringens* type C infections.

Veterinary Microbiology, 21;153(1-2):37-43.

Uzal, F. A.; Plumb, J. J.; Blackell, L. L. and Kelly, W. R. (1997):

PCR detection of *Clostridium perfringens* producing different toxins in faeces of goats.

Letters in Applied Microbiology, 25(5): 339-344.

Vaikosen, E. S. and Muller, W. (2001):

Evaluating biochemical test for isolation and identification of *Clostridium perfringens* in fecal samples of small ruminants in Nigeria.

Bulletin of animal health and production in Africa, 49(4): 244-248.

Veillon, A. and Zuber, A. (1898):

Studies on the strictly anaerobic microbes and their role as pathogens.
Archive of Medical Experiment, 10:517-522.

Warren, C.A.; Calabrese, G.M.; Li, Y.; Pawlowski, S.W.; Figler, R.A.; Rieger, J.; Ernst, P.B.; Linden, J. and Guerrant, R.L. (2012):

Effects of adenosine A2A receptor activation and alanyl-glutamine in *Clostridium difficile* toxin-induced ileitis in rabbits and cecitis in mice.
BMC Infectious Diseases. 20;12(1):13.

Welch, W.H. and Nutall, G.H.F. (1892):

Diarrhea caused by enterotoxigenic *Clostridium Perfringens*.
Bulletin John Hopkins Hospital, 3:81-91.

Willis, A. T. (1977):

Anaerobic Bacteriology in Clinical Medicine.
2nd Ed., London, Butterworths.

Willis, A. T. and Hobbs, G. (1959):

Some new media for the isolation and identification of *Clostridea*.
Journal of Pathology and bacteriology, XX: 511.

Wildson, A. J. (1931):

Observation on the classification of *Bacillus Welchii*.
2nd Reproductive Dir Institute Animal Pathology Cambridge, P. 53-85.

Wilson, G. S. and Miles, A. A. (1975):

Principle of bacteriology, virology and immunology.
6th Ed., Vol. I and Vol. II, Baltimore: Williams and Wikins.

Yamagishi, T.; Sugitani, K.; Tanishima, K. and Nakamura, S. (1997):

Polymerase chain reaction for differentiation of five types of *C. perfringens*.

Microbial Immunology, 41(4):295-299.

Yao ChunYang and Zhuang JinQiu (2007):

Control of *Clostridium welchii* in rabbits in winter-spring.

Chinese Journal of Rabbit Farming, 2:8-10.