

## LIST OF TABLES

| No. | Title  | Page |
|-----|--|------|
| 1   | Primer of alpha toxin gene of <i>C. perfringens</i> used in conventional PCR   | 33   |
| 2   | Primers for the four toxins genes of <i>C. perfringens</i> used in multiplex PCR   | 35   |
| 3   | Interpretation of the results of dermonecrotic test  | 45   |
| 4   | PCR cycling protocol for alpha gene of <i>C. perfringens</i> using a conventional PCR  | 47   |
| 5   | PCR cycling protocol for the four toxins genes of <i>C. perfringens</i> using a multiplex PCR.   | 48   |
| 6   | Interpretation of inhibition zone standards of the used antibiotic discs (Oxoid).  | 50   |
| 7   | Experimental design of the experiment  | 52   |
| 8   | History of the examined rabbits at different Egyptian governorates   |      |
| 9   | The total number and the status of the examined rabbits at different Egyptian governorates   | 58   |
| 10  | Detailed history of the different types and numbers of samples at different Egyptian governorates  | 58   |
| 11  | Isolation of <i>Clostridial</i> organisms (single and mixed infections) recovered from surveyed rabbit's farms at different Egyptian governorates                  | 64   |
| 12  | Recovered <i>Clostridial</i> organisms from surveyed rabbit's farms at different Egyptian governorates   | 65   |
| 13  | Morphological and microscopical appearance of suspected <i>Clostridial</i> strains from surveyed rabbit's farms at different Egyptian governorates                 | 67   |
| 14  | Biochemical identification of suspected <i>Clostridial</i> strains from surveyed rabbit's farms at different Egyptian governorates                                 | 69   |
| 15  | Prevalence of <i>C. perfringens</i> and <i>Clostridial</i> spp. other than <i>C. perfringens</i> isolated from rabbit's samples at different Egyptian governorates | 70   |

|    |   |            |
|----|---|------------|
| 16 | Prevalence of <i>C. perfringens</i> and <i>Clostridial</i> spp. other than <i>C. perfringens</i> isolated from rabbit's samples at different Egyptian governorates                      | <b>71</b>  |
| 17 | The incidence of the isolated <i>Clostridial</i> strains at different Egyptian governorates   | <b>72</b>  |
| 18 | Prevalence of toxigenic types of <i>C. perfringens</i> recovered from surveyed rabbits farms at different Egyptian governorates   | <b>76</b>  |
| 19 | Types of isolated toxigenic <i>C. perfringens</i> recovered from surveyed rabbit's farms at different Egyptian governorates   | <b>76</b>  |
| 20 | Different types of <i>C. perfringens</i> among surveyed rabbit's farms at different Egyptian governorates   | <b>77</b>  |
| 21 | Prevalence of <i>Clostridial</i> spp. other than <i>C. perfringens</i> among surveyed rabbit's farms at different Egyptian governorates   | <b>78</b>  |
| 22 | Recovery rate of single and mixed infections of <i>Clostridial</i> spp. from surveyed rabbit's farms  | <b>80</b>  |
| 23 | <i>In-vitro</i> antibiotic sensitivity test of different types <i>C. perfringens</i> and <i>C. difficile</i> recovered from surveyed rabbit's farms at different Egyptian governorates. | <b>85</b>  |
| 24 | Daily observed mortalities in each rabbits' group during 12 days post challenge   | <b>88</b>  |
| 25 | The weekly feed consumption, weight gain and feed conversion rate of experimentally infected oral groups during observation period  | <b>100</b> |