



Isolation, Characterization and Pathogenicity of the Most Common Bacteria Associated with Gut Health in Egyptian Broiler Chicken Flocks

Ali Z. Qandoos¹, Nayera M. Alatfeehy² and Wafaa A. Abd El-Ghany^{1*}

¹Poultry Diseases Department, Faculty of Veterinary Medicine, Cairo University, 12211, Giza, Egypt

²Research Laboratory for Veterinary Quality Control, Animal Health Research Institute, Egypt

*Corresponding author: wafaa.soliman1974@gmail.com

Article History: 21-309

Received: 01-May-21

Revised: 28-May-21

Accepted: 31-May-21

ABSTRACT

This study was designed to isolate and characterize the pathogenicity of the most common bacteria causing enteric diseases in broilers in some Egyptian governorates. Enteric bacterial organisms like *Escherichia coli* (*E. coli*), *Salmonellae*, and *Clostridia perfringens* (*C. perfringens*) were isolated and identified from 100 dead and diseased broilers in Giza, El-Kalubia and El-Sharqia governorates. The samples were subjected to conventional isolation as well as biochemical and serological identification techniques. Molecular and toxigenic detection of *C. perfringens* were performed on one selected isolate by multiplex PCR. The pathogenicity test of some isolated bacterial strains was done on 80 a-day-old broiler chicks. The clinical observation, performance parameters, bacterial re-isolation and histopathological examination were carried out after bacterial challenge. The results revealed isolation of *E. coli*, *Salmonella* and *C. perfringens* in rates of 17, 11 and 39%, respectively. Serological identification of *E. coli* revealed that O78 (35.2%) and O1 (23.5%) were the highest isolated serotypes, while O117 (17.6%), O91 (11.7%), O112, and O146 (5.8% each) were the lowest ones. *Salmonella enteritidis* (*S. enteritidis*) was the most prevalent (36.3%) followed by *S. kentucky*, *S. larochelle* (27.2%, each) and *S. inganda* (9%). Multiplex PCR of *C. perfringens* strains revealed presence of both *cpa* and *cpb* genes that encoding to alpha (α) and beta (β) lethal toxins, respectively. The pathogenicity of *E. coli* O78, *S. enteritidis*, and *C. perfringens* were tested in broiler chickens. The results showed that *E. coli* O78, *S. enteritidis* and *C. perfringens* were pathogenic strains. It was concluded that enteric bacterial pathogens especially *E. coli* O78, *S. enteritidis*, and *C. perfringens* type C are widely distributed in Egyptian broiler chicken flocks and still causing severe losses because of mortalities and decreasing in production.

Key words: *E. coli*; *Clostridium*; Broiler; Egypt; *Salmonella*.

©2021 IJVS - All Rights Reserved

INTRODUCTION

Animal production especially poultry sector represents as a major source of high-quality human's food worldwide. This sector is subjected for infectious and non-infectious constrains that adversely affect the productivity. Enteric infection is regarded as one of the most important problem that causes severe damage of the digestive tract of poultry (Reynolds 2003). Poor feed efficiency parameters as well as high mortalities of birds resulting from intestinal damage due to bacterial enteric infections (Awaad et al. 2019). Enteric diseases like salmonellosis, colibacillosis, and Clostridial infection have a significant and adverse impact on poultry industry as a result of severe worldwide economic losses. These losses include poor growth performance, increased

mortality, zoonotic significance as well as increased medication costs (Hussein et al. 2013; Soliman et al. 2018).

Avian colibacillosis caused by *Escherichia coli* (*E. coli*) is one of the main widespread bacterial infection that causes losses and a decrease in the production (Ronco et al. 2017; Ellakany et al. 2019; Mehmood et al. 2020). Some strains of *E. coli* are presented as intestinal commensal of birds, but in weak immune hosts, the pathogenic or even non-pathogenic strains can cause diseases conditions (Sarba et al. 2019). Infection with *E. coli* in poultry may induce acute septicemia with high mortality or appear in subacute form with polyserositis (Calnek et al. 1997). The most prevalent serogroups of pathogenic *E. coli* in chickens are belonging to O78, O1, and O2, and to a lesser extent, O15 and O55 (Rahman et al. 2004; Ali et al. 2019).

Cite This Article as: Qandoos AZ, Alatfeehy NM and Abd El-Ghany WA, 2022. Isolation, characterization and pathogenicity of the most common bacteria associated with gut health in Egyptian broiler chicken flocks. International Journal of Veterinary Science 11(1): 7-15. <https://doi.org/10.47278/journal.ijvs/2021.069>

Salmonellosis caused by paratyphoid *Salmonella* species, is one of the major worldwide foodborne pathogens that transmit to human through consumption of raw or insufficient prepared poultry meat. In addition to the significant public health problem, paratyphoid *Salmonella* species have adverse economic impact on poultry industry that represented by high mortality and morbidity, and impaired production (Shoab et al. 2017; Alam et al. 2020). Multiple *Salmonella* serotypes induce different diseases conditions in nearly all types of avian species (Su et al. 2011). However, *Salmonella enteritidis* (*S. enteritidis*) is regarded as the most dominating serovar that affecting poultry flocks (Schlundt 2002; Afshari et al. 2018; Yasmin et al. 2020; Ahmad et al. 2020).

Necrotic enteritis is an important disease condition that is characterized by high mortality, poor feed conversion, weight losses as well as decreasing in egg production of the affected birds (Sarkar et al. 2013; la Mora et al. 2020). Coccidial affection is the main predisposing factor for initiation of necrotic enteritis (Palliyeguru et al. 2010). The disease in poultry is regards as enterotoxaemia caused by *Clostridium perfringens* (*C. perfringens*) (Williams 2005). Based on the production of major lethal toxins α , β , ϵ and iota (i). *Clostridium perfringens* can be classified into five toxin production biotypes (A, B, C, D, and E) (Uzal et al. 2014). Types A and C of *C. perfringens* are the most important types that affecting poultry. However, *C. perfringens* type A is regarded as the most common type and to a lesser extend type C. Strains of *C. perfringens* type A have gene coding α -toxin (*cpa*), while type C have both α -toxin (*cpa*) and β -toxin (*cpb*) gene encoding (Enstrom et al. 2003; Abd El-Hamid et al. 2015). Accordingly, this work was designed to isolate, characterize, and detect the pathogenicity of the most common bacteria causing enteric diseases in broiler chickens in some Egyptian governorates.

MATERIALS AND METHODS

The pathogenicity test was carried out after the approval of the Institutional Animal Care and Use Committee, Cairo University (CU-IACUC) (CU II F 3 20).

Samples Collection

This research work was conducted during the period from April till September 2019. A total of 100, 4-6-week-old diseased and freshly dead broilers with a history of diarrhea, depression, and reduced growth performance were collected from different flocks in Giza, El-Kalubia, and El-Sharkia governorates of Egypt. The examined dead birds showed severe enteritis, general congestion of the internal abdominal viscera as well as perihepatitis, pericarditis and peritonitis. Samples including heart, liver, and intestine were collected, identified, and rapidly

transported to the laboratory in an icebox for further processing.

Isolation and Characterization of the Enteric Bacterial Strains

Isolation of *E. coli* was done according to Lee and Nolan (2008). Collected samples were cultured on tryptic soy broth and incubated at 37°C for 18 hours. A loop full of incubated broth was streaked into MacConkey's agar and incubated at 37°C for 18-24 hours. The suspected *E. coli* colonies were subjected for morphological and biochemical identification. The isolates were serologically identified as Kok et al. (1996) by using rapid diagnostic *E. coli* antisera sets (Denka Seiken Co, Japan).

Besides, isolation and identification of *Salmonella* species were done according to standard methods (ISO 6579 2002). Rappaport-Vassiliadis enrichment broth was used for isolation of *Salmonella* and incubated at 37°C for 18 hours. A loop-full of culture was streaked into Xylose Dextrose (XLD) and MacConkey agar and incubated at 37°C for 18-24 hours. The morphological and bio-chemical identification were carried out on the suspected colonies, while the serotyping was done as described by Kauffman-white scheme (Kauffman 1974; Popoff 2001) for somatic (O) and flagella (H) antigens using *Salmonella* antiserum (DENKA SEIKEN Co. Japan).

For *Clostridia* isolation, the samples were inoculated into a cooked meat broth medium and incubated in an anaerobic gas pack jar for 24 hours at 37°C. A loop full of broth culture was streaked onto 10% sheep blood agar, supplemented with neomycin sulfate (200µg/ml) (Carter and Cole 1990) and incubated anaerobically at 37°C for 24 hours. Gram staining from suspected colonies were prepared for microscopical examination. Biochemical identification was applied (Koneman et al. 1988). The strains were maintained in 25% glycerol stock at -80°C until further use.

Detection of *Clostridium perfringens* Toxins using Multiplex PCR

Genomic DNA was extracted from *C. perfringens* strain by culturing on blood agar media using QIAamp Mini Kit (Cat No. 51304). The specific three primer sets design corresponding to each α , β and ϵ toxins of *C. perfringens* (Komoriya et al. 2007; van Asten et al. 2009) is shown in Table 1. The PCR was performed in a Thermal Cycler® (Bio-Rad, USA) in a total reaction volume of 50M containing 25M of 2x Master Mix (thermos scientific, 1M of 1pmol of each primer and 5M of template DNA). Cycling conditions of the primers during multiplex PCR is represented in Table 2. Then, 8M of the amplified product was electrophoresed in a 1.5% agarose gel and stained with ethidium bromide. Amplified bands were visualized and photographed under ultraviolet illumination.

Table 1: Primers used in multiplex PCR for the detection of toxin's type of *C. perfringens*

Amplicon (bp)	Primer position	Sequence	Gene	Toxin
324	663-968	GCTAATGTTACTGCCGTTGA CCTCTGATACATCGTGTAAG	<i>Plc (cpa)</i>	α
196	871-1045	GCGAATATGCTGAATCATCTA GCAGGAACATTAGTATATCTTC	<i>cpb</i>	β
655	267-862	GCGGTGATATCCATCTATTC CCACTACT TGTCCTACTAAC	<i>etx</i>	ϵ

The Pathogenicity Test

Eighty, day-old chicks obtained from local hatchery were divided into 4 equal groups consisting of 20 birds each. Each group was housed in a thoroughly cleaned and disinfectant pen, given feed and water *ad libitum* as well as vaccination programs according to the manufacturer's instructions. Group 1 was a negative control non-challenged birds, while groups 2, 3, and 4 were challenged orally at 2 weeks of age. Each bird of the challenged groups was inoculated with 1ml containing 1×10^8 Colony Forming Unit (CFU) for *C. perfringens* type C and 1×10^9 for *S. enteritidis* and *E. coli* O78 (Barrow et al. 1999; Botlhoko 2009; Johny et al. 2012; Abd El-Tawab et al. 2015b). Chickens in *C. perfringens* challenged group were orally inoculated with a mixture of *Eimeria* species oocysts (0.25ml/bird) at 12 days old to stimulate necrotic enteritis induction (Williams et al. 2003). The body weight and feed consumption of each group were taken weekly to measure the feed conversion ratio (FCR). Moreover, clinical signs, mortality rates, and post-mortem lesions of chickens in the challenged groups were recorded every day for 3 weeks later. Specimens including heart, liver, and intestine were collected from dead and euthanized birds (at the 3rd week post-challenge) for bacterial re-isolation and histopathological examination.

The Histopathological Examination

Specimens including liver, intestine and heart were collected and fixed in 10 % formalin solution, dehydrated in different grades of ethyl alcohol, and embedded in paraffin. Tissues of 5 microns thickness were stained with hematoxylin and Eosin (H&E) and microscopically examined using light microscope (Slaoui and Fiette 2011).

Statistical Analysis

Data were expressed as mean \pm SE. Statistical comparison between the mean of the different groups was made by One-Way Analysis of Variance (ANOVA) and multiple comparisons between groups (post hoc) LSD

using SPSS version 26. A probability $P \leq 0.05$ was assumed for statistical significance.

RESULTS

The results of isolation of some enteric bacterial strains in different Egyptian governorates are illustrated in Table 3. From the table, it could be observed that the isolation rate of *E. coli* from 17 out of 100 samples was 17%. Culturing of suspected *E. coli* samples on MacConkey's agar showed medium-sized, rounded, and pink colonies. Gram's staining revealed Gram-negative, non-sporulated bacilli that arranged single, pairs, and in groups. In addition, the biochemical tests of the suspected isolates showed negative oxidase, urease, citrate utilization, and Voges Proskauer tests, but positive catalase, indole, methyl red, nitrate reduction and triple sugar iron tests. The isolates showed motility in semisolid agar media. The serological identification of the suspected *E. coli* isolates in different Egyptian governorates is shown in Table 4. The results demonstrated that the most predominant serotypes were O78 (6 isolates) followed by O1 (4 isolates), O117 (3 isolates), O91 (2 isolates), and O112, O146 (one isolate; each).

Regarding the isolation of *Salmonellae*, the isolation rate of *Salmonellae* was 11% as 11 out 100 samples were positive (Table 3). Colonies of suspected *Salmonellae* cultures appeared as pink with black center on XLD agar and colorless ones on MacConkey agar, while Gram-negative medium size bacilli were seen in stained culture smears. Biochemical identification of cultures showed positive catalase, methyl red, citrate utilization, H₂S production tests, while negative oxidase, indole, urease, Voges Proskauer, and gelatin liquefaction tests. Sugar fermentation was positive for glucose, maltose, and xylose, whereas negative for lactose and sucrose. Serological identification of the suspected *Salmonella* isolates in different Egyptian governorates was *S. enteritidis* (4), *S. kentucky* (3), *S. larochelle* (3), and *S. inganda* (1) as represented in Table 5.

Table 2: Cycling conditions of the primers during multiplex PCR for the detection of toxin's type of *C. perfringens*

Primary Denaturation	Secondary Denaturation	Annealing	Extension	No. of cycles	Final Extension
95°C	94°C	55°C	72°C	35	72°C
10 min.	45 sec.	30 sec.	90 sec.		10 min.

Table 3: The isolation rate of enteric bacterial strains in different Egyptian governorates

The governorate	Status of examined birds	No. of samples	The isolated enteric bacterial isolates					
			<i>E. coli</i>		<i>Salmonella</i>		<i>Clostridia</i>	
			No. of positive samples	% of positive samples	No. of positive samples	% of positive samples	No. of positive samples	% of positive samples
Giza	Diseased	20	2	10*	1	5*	5	25*
	Dead	6	1	16.7*	1	16.7*	2	33.3*
	Total	26	3	11.5**	2	33.3**	7	26.9**
El-Kalubia	Diseased	35	5	14.3*	4	11.4*	18	51.4*
	Dead	8	3	37.5*	2	25*	3	37.5*
	Total	43	8	18.6**	6	14**	21	48.8**
El-Sharqia	Diseased	25	4	16*	0	0	9	36*
	Dead	6	2	33.3*	3	50*	2	33.3*
	Total	31	6	19.9**	3	9.7**	11	35.5**
Total		100	17	17***	11	11***	39	39***

*The percentage was calculated according to each bird's status; **The percentage according to the total number in each governorate;

***The percent according to the total number of tested samples.




Fig. 1: Agarose gel (1.5%) electrophoresis of multiplex PCR products of *C. perfringens* toxins; Lane 1: DNA marker (Gene Ruler 100 bp DNA Ladder, Thermo-scientific); Lane 2: Control Positive (mix of various toxin types); Lane 3: Control negative and Lane 4: PCR product of the isolate toxin.

Table 4: The serological identification of *E. coli* isolates in different Egyptian governorates

The governorate	Total No. of isolated <i>E. coli</i>	Serogroup	No. of each serotype
Giza	3	O78	1
		O146	1
		O91	1
El-Kalubia	8	O78	3
		O1	3
		O91	1
		O112	1
El-Sharkia	6	O117	3
		O78	2
		O1	1
Total		17	

Table 5: The serological identification of *Salmonella* isolates in different Egyptian governorates

The governorate	Total No. of isolates	Serotypes	No. of each serotype
Giza	2	<i>S. kentucky</i>	1
		<i>S. inganda</i>	1
El-Kalubia	6	<i>S. kentucky</i>	2
		<i>S. enteritidis</i>	2
		<i>S. larochelle</i>	2
El-Sharkia	3	<i>S. enteritidis</i>	2
		<i>S. larochelle</i>	1
Total		11	

The results of *C. perfringens* isolation rate that presented in Table 3 demonstrated that out of 100 examined samples, 39 (39%) were positive. Anaerobic incubation of suspected samples on sheep blood agar showed presence of small, circular, smooth, flat, glistening and greyish colonies surrounded by double zones of hemolysis. Microscopically, Gram-positive spore-forming bacilli were noticed. Biochemically, positive results for lecithinase activity on yolk agar as well as sugar fermentation of glucose, lactose, maltose, and sucrose were demonstrated. Nevertheless, negative results for indole production, catalase, and oxidase were also found. The results of multiplex PCR for detection of the toxin types for *C. perfringens* revealed presence of the gene coding α toxin (*cpa*) and the gene encoding β toxin (*cpb*) (Fig. 1). The latter results indicated presence of *C. perfringens* type C.

Regarding the pathogenicity test, the mortality rate and the post-mortem lesions in different challenged groups are seen in Table 6. No clinical signs were appeared in negative control non-challenged birds, but birds challenged with *E. coli* O78 showed depression, brownish diarrhea, nasal and ocular discharge and conjunctivitis. Greenish diarrhea and poor growth were seen in chickens challenged with *S. enteritidis*, while bloody diarrhea, and ruffled feather were noticed in birds challenged with *C. perfringens*. Control negative non challenged birds showed no mortalities while chickens challenged with *E. coli* O78, *S. enteritidis*, and *C. perfringens* revealed mortality rates of 25, 15 and 35%, respectively. The necropsy findings of dead and sacrificed bird's revealed perihepatitis, pericarditis and enteritis in group challenged with *E. coli* O78; enlarged liver with necrotic foci, perihepatitis, enlarged congested kidney and enteritis in group challenged with *S. enteritidis*, as well as friable ballooned intestine with hemorrhagic mucosa and necrotic pale liver in group *C. perfringens*.

Table 7 shows the average body weight and FCR in different groups. In comparison with control negative non-challenged group, challenged groups revealed significant decrease in body weight and feed consumption with $P \leq 0.05$ that reflected on the FCR.

The re-isolation rates of bacteria from different organs in different challenged groups are present in Table 8. The re-isolation rates of *E. coli*, *Salmonella*, and *Clostridia* from visceral organs were 85, 70, and 85% in birds challenged with *E. coli* O78, *S. enteritidis* and *C. perfringens*, respectively. In a group challenged with *E. coli* O78, the highest incidence of re-isolation was from heart (88.2%), then from liver (76.4%) and intestine (70.5%). For chickens challenged with *S. enteritidis*, the highest re-isolation was from liver (92.8%), intestine (85.7%) followed by heart (64.2%), while in birds challenged with *C. perfringens*, this rate was 88.2% from the intestine and 70.5% from the liver.

Fig. 2 to 4 show the histopathological examination results in the challenged groups of birds. The different lesions in the liver, intestine and heart of group challenged with *E. coli* O78 are demonstrated (Fig. 2A to 2D). Chickens exhibited diffuse hepatocellular necrosis which was characterized by numerous sporadic cell necrosis among the hepatocytes admixed with accumulation of diffuse edema, eosinophilic debris, and a variable number of mononuclear inflammatory cells infiltration. The intestine showed severe necrosis of the intestinal villi associated with intense inflammatory cells infiltration. The myocardial muscle revealed diffuse myocarditis which was illustrated by extensive number of mononuclear inflammatory cells infiltration.

Fig. 3A to 3D show the hepatic and intestinal microscopic alterations of *S. enteritidis* challenged chickens. The hepatic parenchyma showed multifocal edematous areas accompanied by dilation of the hepatic sinusoids. Numerous clusters of basophilic septicemic bacteria were detected in the hepatic sinusoids. The hepatocytes suffered from widespread coagulative necrosis. The intestinal sections showed severe necrosis which is characterized by atrophy of intestinal villi associated with accumulation of eosinophilic and karyorrhectic debris.

Table 6: The mortality rates and post-mortem lesions in different challenged groups

Group No.	Challenge bacteria	The mortality rate	Post-mortem lesions
1	-	0 (0)	-
2	<i>E. coli</i> (O78)	5 (25)	Perihepatitis, pericarditis and enteritis
3	<i>S. enteritidis</i>	3 (15)	Enlarged liver with necrotic foci, perihepatitis, enlarged congested kidney and enteritis
4	<i>C. perfringens</i>	7 (35)	friable ballooned intestine with hemorrhagic mucosa and necrotic pale liver

Values in parenthesis indicate percentage.

Table 7: The average body weight (g/bird) and the FCR in different groups

Group No.	Challenge bacteria	Average body weight (g/bird)					FCR
		Weeks before infection		Weeks after infection			
		W1	W2	W3	W4	W5	
1	-	160.7±3.1 ^a	422.2±10.0 ^a	853.0±19.3 ^a	1347.0±9.7 ^a	1898.0±26.0 ^a	1.673 ^a
2	<i>E. coli</i> (O78)	161.0±4.6 ^a	430.7±10.9 ^a	762.1±12.6 ^b	1223.8±15.0 ^b	1681.2±15.5 ^b	1.794 ^b
3	<i>S. enteritidis</i>	155.2±4.0 ^a	432.0±10.4 ^a	777.1±17.9 ^b	1223.6±14.1 ^b	1682.5±14.4 ^b	1.777 ^b
4	<i>C. perfringens</i>	163.25±4.4 ^a	425.2±10.0 ^a	757.5±16.8 ^b	1212.3±18.8 ^b	1676.0±20.8 ^b	1.789 ^b

W=Week; FCR=Feed conversion ratio. Mean±SE in the same column followed by the different superscripts is significantly different according to ANOVA (LSD, P<0.05).

Table 8: The re-isolation rates of bacteria from different organs in different challenged groups

Group No.	Challenge bacteria	The total re-isolation rate	The re-isolation rate from different organs		
			Liver	Heart	Intestine
1	-	0 (0)	0 (0)	0 (0)	0 (0)
2	<i>E. coli</i> (O78)	17/20 (85)	13/17 (76.4)	15/17(88.2)	12/17 (70.5)
3	<i>S. enteritidis</i>	14/20 (70)	13/14 (92.8)	9/14 (64.2)	12/14 (85.7)
4	<i>C. perfringens</i>	17/20 (85)	12/17 (70.5)	0/14 (0)	15/17 (88.2)

Values in parenthesis indicate percentage.




Fig. 2: A) Liver of *E. coli* challenged group showing sporadic cell necrosis (arrows); B) Intestine of *E. coli* challenged group showing severe necrosis of the intestinal villi (arrow); C) Intestine of *E. coli* challenged group showing intestinal necrosis admixed with accumulation of inflammatory cells and tissue debris in the submucosa (arrow) and Heart of *E. coli* challenged group showing myocarditis (arrow). H&E. Bar=25µm (A, B and D); 50µm (C).

In *C. perfringens* challenged group, variable lesions have been noticed in the liver and intestine (Fig. 4A to 4D). The liver showed multifocal to diffuse hemorrhagic areas accompanied by disorganization of the hepatic parenchyma. The hepatocytes showed marked necrobiotic changes that showed vacuolation and several stages of nuclear necrosis that including karyorrhexis, karyolysis, and pyknosis. The intestine revealed marked atrophy of the intestinal villi accompanied by enteritis that showed expansion of the submucosal layer with edema and mononuclear inflammatory cells infiltration.




Fig. 3: A) Liver of *S. enteritidis* challenged group showing congested hepatic sinusoids, edema and few inflammatory cells infiltration; B) Liver of *S. enteritidis* challenged group showing multifocal aggregation of bacterial colonies (arrows); C) Intestine of *S. Enteritidis* challenged group showing severe necrosis of the intestinal villi, crypts and intestinal glands (arrow) and D) Intestine of *S. enteritidis* challenged group showing severe necrosis of the intestinal glands admixed with mononuclear inflammatory cells infiltration in the submucosa (arrow). H&E. Bar=25µm (A, B and D); 50µm (C).

DISCUSSION

Enteric bacterial diseases are regarded as one of the most important group of diseases that affect broiler chickens and causes severe economic losses in poultry production (Hafez 2011; Abdel-Alim and Ahmed 2020). Therefore, this study is focused on isolation and characterization of some important enteric bacteria in different Egyptian governorates.




Fig. 4: A) Liver of *C. perfringens* challenged group showing hepatic hemorrhages; B) Liver of *C. perfringens* challenged group showing several stages of nuclear necrosis in the affected hepatocytes (arrows); C) Intestine of *C. perfringens* challenged group showing marked atrophy and shortening of the intestinal villi (arrow) and D) Intestine of *C. perfringens* challenged group showing expansion of the submucosa with edema and mononuclear inflammatory cell accumulation (arrows). H&E. Bar=25µm (A, B and D); 50µm (C).

Here, *E. coli* strains have been detected in 17 out of 100 samples with isolation rate of 17%. Previous Egyptian studies on *E. coli* revealed either higher isolation rates 75% (Abd El-Tawab et al. 2015a), 43.1% (Heba et al. 2012), 35% (Amer et al. 2018), 34.6% (Ellakany et al. 2019) and 27.7% (Abd El-Mongy et al. 2017), while other showed lower isolation rate 8.3% (Ahmed et al. 2017). These differences in the isolation rates may be owing to the differences in the number of samples, the age of birds, the geographic regions, the isolation methods, and previous medication. The cultural morphology, Gram staining characters and biochemical reactions of suspected *E. coli* isolates were similar to those reported by Hadiujjaman et al. (2016), Surra et al. (2018) and Ibrahim et al. (2019). Moreover, the serological identification results of this study revealed that the main serotypes of the isolated *E. coli* were O78 and O1. These findings are parallel to the studies of Rahman et al. (2004) and Ibrahim et al. (2019) who found that *E. coli* O78, O1, and O2 were the predominant serotypes of the pathogenic *E. coli* in chickens. Besides, the Egyptian study of Amer et al. (2018) declared that 25% of *E. coli* isolates were O78.

This study showed isolation of *Salmonellae* in percentage of 11 (11/100). Similar isolation rates were previously reported by Kaoud et al. (2018) and Soliman et al. (2018) who isolated *Salmonellae* from the Egyptian broiler flocks in rates of 11.33 % and 10.37%, respectively. Moreover, in Seri Lanka, Jayaweera et al. (2020) demonstrated isolation of *Salmonellae* from broiler chickens in rate of 11.6%. However, Egyptian studies of Sedeik et al. (2019) and El-Sheikh et al. (2019) showed isolation of *Salmonellae* from broiler chicks in rates of 7.5% and 16%, respectively. Regarding the colonial morphology and biochemical tests results of *Salmonella*, parallel results were also described by Akeila et al. (2013), Islam et al. (2016), Al-Mamun et al. (2017) and Mridha et al. (2020). Moreover, the serological results of the isolated *Salmonellae* showed that *S. enteritidis* and *S. kentucky* were

the most among the isolates. Similar findings were reported by Moawad et al. (2017), Al-Baqir et al. (2019) and Sedeik et al. (2019) who demonstrated that *S. enteritidis* and *S. kentucky* were the most highly prevalent strains that isolated from chicken farms in Egypt. In addition, Wang et al. (2020) demonstrated variations of *Salmonella* serotypes in chicken's flocks and found that *S. enteritidis* was the most common isolated strain.

In addition, *C. perfringens* was isolated in rate of 39%. Nearly similar isolation rate of *C. perfringens* was recorded in the study of Osman et al. (2012) who found that out of 48 broiler chicken samples, 35.4% were *C. perfringens*. Moreover, Abd-Elall and Maysa (2014) and Helal et al. (2019) declared that the prevalence of *C. perfringens* in broilers was 38.7 and 38.3%, respectively. However, higher isolation rates of *C. perfringens* (75%) and (77.4%) were observed in the studies of El-Jakee et al. (2013) and Abd El-Hamid et al. (2015), respectively. The phenotypic characterization findings of *C. perfringens* isolates were similar to others (Das et al. 2008; Malmarugan et al. 2012; Dar et al. 2017). The results of multiplex PCR of *C. perfringens* strains confirmed presence of the gene coding α -toxin (*cpa*) and the gene encoding β -toxin (*cpb*). Although *C. perfringens* type A is the predominant worldwide biotype type that isolated from chickens (Hatakka and Pakkala 2003; Merati et al. 2017), this works proved isolation of *C. perfringens* type C. Few studies either worldwide or locally showed isolation of *C. perfringens* type C from broiler chickens (Afshari et al. 2015).

Challenged groups with *E. coli* O78, *S. enteritidis*, and *C. perfringens* should variable diarrhea, respiratory signs and poor growth rate along with mortality rates of 25, 15 and 35%, respectively. Autopsy findings showed polyserositis and enteritis, liver necrosis, congested kidney and enteritis, along with friable ballooned intestine with haemorrhagic mucosa and necrotic pale liver in groups challenged with *E. coli* O78, *S. enteritidis*, and *C. perfringens*, respectively. Similar post-mortem observations in different groups were observed by Cooper and Songer (2010), Belih et al. (2016) and El-Sawah et al. (2018).

It has been observed significant ($P \leq 0.05$) differences in the average body weight and FCR in the challenged groups when compared with non-challenged group. Our results agreed with Abd El-Tawab et al. (2015b) who demonstrated depressive effect of *E. coli* on the performance parameters of chickens, Belih et al. (2016) who found significant reduction in body weight of chicken after challenge with *Salmonella*, as well as Abd El-Hamid et al. (2017) who demonstrated decrease in body weight, feed consumption and FCR in *C. perfringens* challenged chickens.

Re-isolation rates of 85%, 70%, and 85% have been recorded in chickens challenged with *E. coli* O78, *S. enteritidis* and *C. perfringens*, respectively. These results are in partial agreement with Atta et al. (2014), Abd El-Tawab et al. (2015a) and Gast et al. (2013).

The histopathological findings of chicken's liver, intestine and heart in group challenged with *E. coli* O78 are supported by the results obtained by Abalaka et al. (2017) and Shah et al. (2019). However, the microscopic changes of the liver and intestine of *S. enteritidis* challenged

chickens revealed similar results to Saha et al. (2012) and Muna et al. (2016). The microscopic findings of the intestine and liver of *C. perfringens* challenged group correspond with other authors (Abd El-Hamid et al. 2015; Park et al. 2015).

From the previous findings, it may be concluded that many Egyptian broiler chicken's flocks showed circulation of different types of enteric bacterial pathogens, particularly *E. coli* O78, *S. enteritidis*, and *C. perfringens* type C. These organisms are responsible for problems in poultry flocks resulting from increased morbidity, mortality, loss of productivity and public health significance. Periodical surveillance studies should be conducted to detect the epidemiological situation of these infections in poultry flocks. In addition, application of biosecurity measure is essential to prevent and/or eliminate these pathogens.

Author's Contribution

Ali Z. Qandoos shared in the laboratory work, the experimental study and the collection of data, Nayera M. Alatfeehy shared in the laboratory work and experimental study and collection of data and Wafaa A. Abd El-Ghany designed the work, shared in the experimental study along with writing and submission of the manuscript. All authors approved the final version.

Financial Support

This study was conducted without any financial support from any agency.

REFERENCES

- Abalaka SE, Sani NA, Idoko IS, Tenuche OZ, Oyelowo FO, Ejeh SA and Enem SI, 2017. Pathological changes associated with an outbreak of colibacillosis in a commercial broiler flock. *Sokoto Journal of Veterinary Sciences* 15: 95-102. <https://doi.org/10.4314/sokjvs.v15i3.14>
- Abd-Elall AM and Maysa AI, 2014. Toxin genotyping of *Clostridium perfringens* isolated from broiler and layer farms and their workers in Egypt. *Revue de Medecine Veterinaire* 165: 272-279.
- Abd El-Hamid HS, Ellakany HF, Bechet AA, Elbestawy AR and Bedawy S, 2015. Pathogenicity and antibiotic sensitivity of *Clostridium perfringens* isolates from chickens in El-Behera governorate, Egypt. *Life Science Journal* 12: 174-182.
- Abd El-Hamid HS, Ellakany HF, Rizk MS, Elbestawy R and Abdelfatah SL, 2017. Effect of combined *Clostridium perfringens* infection and aflatoxicosis in broiler chickens. *Alexandria Journal of Veterinary Sciences* 52: 15-27. <https://doi.org/10.5455/ajvs.239626>
- Abd El-Mongy M, Abd-El-Monem GM, Moawad AA and Mohammed ABA, 2017. Serotyping and virulence genes detection in *Escherichia coli* isolated from broiler chickens. *Journal of Biological Sciences* 18: 46-50. <https://doi.org/10.3923/jbs.2018.46.50>
- Abd El-Tawab AA, Ammar AM, Nasef SA and Reda RM. 2015a. Prevalence of *E. coli* in diseased chickens with its antibiogram pattern. *Benha Veterinary Medical Journal* 28: 224-230. <https://doi.org/10.21608/bvmj.2015.32507>
- Abd El-Tawab AA, El-Komy AA, El-Ekhnawy KI and Talaie AT, 2015b. Effect of fosfomycin on *E. coli* O78 isolated from broiler chickens *in-vitro* and *in-vivo*. *Benha Veterinary Medical Journal* 28: 294-300.
- Abdel-Alim GA and Ahmed KA, 2020. The effect of cephardine on clinicopathological pictures of experimental *Salmonella enteritidis* and *E. coli* infections in broiler chickens. *International Journal of Veterinary Science* 9: 78-83.
- Afshari A, Baratpour A, Khanzade S and Jamshidi A, 2018. *Salmonella enteritidis* and *Salmonella typhimurium* identification in poultry carcasses. *Iranian Journal of Microbiology* 10: 4550.
- Afshari A, Jamshidi A, Razmyar J and Rad M, 2015. Genotyping of *Clostridium perfringens* isolated from broiler meat in northeastern of Iran. *Veterinary Research Forum* 6: 279-284.
- Ahmad Y, Zahra R, Ali MI, Riaz MH, Khan R, Khan K, Khan MT, Ali A and Zeb MT, 2020. Molecular screening of resistant and virulent genes in *Salmonella enteritidis* and *Salmonella typhimurium* from poultry in Khyber Pakhtunkhwa. *Pakistan Veterinary Journal* 40: 343-349. <https://doi.org/10.29261/pakvetj/2020.044>
- Ahmed SAM, Ahmed IA, Osman N and El-Hamd DM, 2017. Problem of *Escherichia coli* infection in broilers in Qena province. *Assiut Veterinary Medical Journal* 63: 252-264.
- Akeila MA, Ellakany HF, Sedeik ME and Behar HM, 2013. Characterization and plasmid profiling of *Salmonella enteritidis* isolated from broiler chickens. *Alexandria Journal of Veterinary Sciences* 39: 105-111.
- Alam SB, Mahmud M, Akter R, Hasan M, Sobur A, Nazir KNH, Noreddin A, Rahman T, El Zowalaty ME and Rahman M, 2020. Molecular detection of multidrug resistant *Salmonella* species isolated from broiler farm in Bangladesh. *Pathogens* 9: 201. <https://doi.org/10.3390/pathogens9030201>
- Ali AI, Abd El-Mawgoud AM, Dahshan AA, El-Sawah AA and Nasef S, 2019. *Escherichia coli* in broiler chickens in Egypt, its virulence traits and vaccination as an intervention strategy. *Novel Research in Microbiology Journal* 3: 415-427. <https://doi.org/10.21608/NRMJ.2019.44950>
- Al-Baqir A, Hussein A, Ghanem I and Megahed M, 2019. Characterization of paratyphoid *Salmonellae* isolated from broiler chickens at Sharkia governorate, Egypt. *Zagazig Veterinary Journal* 47: 183-192.
- Al-Mamun A, Kabir SM, Islam M, Lubna M, Islam SK, Akhter AH and Hossain M, 2017. Molecular identification and characterization of *Salmonella* species isolated from poultry value chains of Gazipur and Tangail districts of Bangladesh. *African Journal of Microbiology Research* 11: 474-481. <https://doi.org/10.21608/ZVJZ.2019.10544.1028>
- Amer MM, Mekky HM, Amer AM and Fedawy HS, 2018. Antimicrobial resistance genes in pathogenic *Escherichia coli* isolated from diseased broiler chickens in Egypt and their relationship with the phenotypic resistance characteristics. *Veterinary World* 11: 1082-1088. <https://doi.org/10.14202/vetworld.2018.1082-1088>
- Atta AH, Shalaby MA and Saifan HY, 2014. Efficacy of commiphoramolmol extract against *Clostridium perfringens* experimental infection in chickens. *World Journal of Pharmacy and Pharmaceutical Sciences* 3: 365-380.
- Awaad MHH, Elmenawey MA, Afify MA, Zouelfekar SA, Mohamed FF and Demy V, 2019. The impact of high stocking density and *Saccharomyces cerevisiae boulardii* on productive performance, intestinal microbiota and gut integrity of broiler chickens. *International Journal of Veterinary Science* 8: 362-370.
- Barrow P, Lovell M, Murphy C and Page K, 1999. *Salmonella* infection in a commercial line of ducks; experimental studies on virulence, intestinal colonization and immune protection. *Epidemiology and Infection* 123: 121-132. <https://dx.doi.org/10.1017/s0950268899002605>
- Belih S, El-Hadad S, Amen G and Basyony M, 2016. Influence of sodium butyrate on *Salmonella* infection in broiler chicks. *Benha Veterinary Medical Journal* 31: 21-32. <https://doi.org/10.21608/bvmj.2016.31255>
- Bothhoko TD, 2009. Performance of *Clostridium perfringens* challenged broilers inoculated with effective

- microorganisms. MSc Agric. (Animal Science), Faculty of Natural and Agricultural Science, Pretoria University.
- Calnek BW, Barnes HJ, Beard CW, McDougald LR and Saif YM, 1997. Diseases of Poultry, 10th Ed, Iowa State University Press, Ames, IA, USA.
- Carter GR and Cole JR, 1990. Diagnostic Procedures in Veterinary Bacteriology and Mycology. 5th Ed, Academic Press, New York, USA.
- Cooper KK and Songer JG, 2010. Virulence of *Clostridium perfringens* in an experimental model of poultry necrotic enteritis. *Veterinary Microbiology* 142: 323-328. <https://doi.org/10.1016/j.vetmic.2009.09.065>
- Dar PS, Wani SA, Wani AH, Hussain I, Maqbool R, Ganaie MY, Kashoo ZA and Qureshi S, 2017. Isolation, identification and molecular characterization of *Clostridium perfringens* from poultry in Kashmir valley, India. *Journal of Entomology and Zoology Studies* 5: 409-414.
- Das A, Mazumder Y, Dutta BK, Shome BR, Bujarbaruah KM and Kumar A, 2008. *Clostridium perfringens* type A from broiler chicken with necrotic enteritis. *International Journal of Poultry Science* 7: 601-609. <https://doi.org/10.3923/ijps.2008.601.609>
- El-Jakee J, Ata NS, El Shabrawy MA, Abu Elnaga ASM, Hedia RH, Shawky NM and Shawky HM, 2013. Characterization of *Clostridium perfringens* isolated from poultry. *Global Veterinaria* 11: 88-94. <https://doi.org/10.5829/idosi.gv.2013.11.1.1138>
- Ellakany HF, Abd-Elhamid HS, Ibrahim MS, Mostafa NS, Elbestawy AR and Gado AR, 2019. Isolation, serotyping, pathogenicity, and antibiotic sensitivity testing of *Escherichia coli* from broiler chickens in Egypt. *Alexandria Journal of Veterinary Sciences* 61: 45-51. <https://doi.org/10.5455/ajvs.10916>
- El-Sawah AA, Dahshan AHM, El-Nahass E and El-Mawgoud AA, 2018. Pathogenicity of *Escherichia coli* O157 in commercial broiler chickens. *Beni-Suef University Journal of Basic and Applied Sciences* 7: 620-625. <https://doi.org/10.1016/j.bjbas.2018.07.005>
- El-Sheikh M, Abdeen E and Ammar A, 2019. Molecular detection of some virulence genes of *Salmonella* serotypes isolated from poultry in Egypt. *Journal of Current Veterinary Research* 1: 86-93. <https://doi.org/10.21608/JCVR.2019.36570>
- Enstrom BE, Fermer C, Lindberg A, Saarinen E, Baverud A and Gunnarsson A, 2003. Molecular typing of isolates of *Clostridium perfringens* from healthy and diseased poultry. *Veterinary Microbiology* 94: 225-235. [https://doi.org/10.1016/s0378-1135\(03\)00106-8](https://doi.org/10.1016/s0378-1135(03)00106-8).
- Gast RK, Guraya R, Jones DR and Anderson KE, 2013. Colonization of internal organs by *Salmonella Enteritidis* in experimentally infected laying hens housed in conventional or enriched cages. *Poultry Science* 92: 468-473. <https://doi.org/10.3382/ps/pey541>
- Hadiujjaman M, Rahman MM, Ahasan MD, Banu MA, Khatun MM and Islam MA, 2016. Isolation and identification of *Escherichia coli* from apparently healthy chicken of selected areas of Bangladesh. *International Journal of Natural and Social Sciences* 3: 15-23.
- Hafez HM, 2011. Enteric diseases of poultry with special attention to *Clostridium perfringens*. *Pakistan Veterinary Journal* 31: 175-184.
- Hatakka M and Pakkala P, 2003. Incidence of Food Borne Infections in Northern Europe. In: *Food Science and Technology*. Marcel Dekker, New York, USA, pp: 669-684. <https://doi.org/10.1201/9780203912065.pt2>
- Heba R, Soad A and Mohamed R, 2012. Incidence of *E. coli* in chickens and ducks in different governorates in Egypt. Conference: First Conference of Animal Health Research Institute Association, pp: 420-426.
- Helal S, Khalaf. N, El Menisy A and Lebdah M, 2019. *Clostridium perfringens* type A causing necrotic enteritis outbreaks among chickens in Egypt. *Zagazig Veterinary Journal* 47: 398-407. <https://doi.org/10.21608/ZVJZ.2019.14139.1055>
- Hussein AH, Ghanem IAI, Eid A, Ali MA, Sherwood JS, Li G, Nolan LK and Logue CM, 2013. Molecular and phenotypic characterization of *Escherichia coli* isolated from broiler chicken flocks in Egypt. *Avian Disease* 57: 602-611. <https://doi.org/10.1637/10503-012513-Reg.1>.
- Ibrahim RA, Cryer TL, Lafi SQ, Basha EA, Good L and Tarazi YH, 2019. Identification of *Escherichia coli* from broiler chickens in Jordan, their antimicrobial resistance, gene characterization and the associated risk factors. *BMC Veterinary Research* 15: 159. <https://doi.org/10.1186/s12917-019-1901-1>.
- ISO 6579, 2002. Microbiology of food and animal feeding stuffs – Horizontal method for the detection of *Salmonella*. ISO 6579: 2002 (E) International Standards Organization, Geneva.
- Islam MJ, Mahbub-E-Elahi A, Ahmed T and Hasan MK, 2016. Isolation and identification of *Salmonella* spp. from broiler and their antibiogram study in Sylhet, Bangladesh. *Journal of Applied Biology and Biotechnology* 4: 046-051. <https://doi.org/10.7324/JABB.2016.40308>
- Jayaweera TS, Ruwandeepika HA, Deekshit VK, Vidanarachchi JK, Kodithuwakku SP, Karunasagar I and Cyril HW, 2020. Isolation and identification of *Salmonella* spp. from broiler chicken meat in Sri Lanka and their antibiotic resistance. *The Journal of Agricultural Sciences - Sri Lanka* 15: 395-410. <https://doi.org/10.4038/jas.v15i3.9031>
- Johny AK, Mattson T, Baskaran SA, Amalrajou MA, Babapoor S, March B, Valipe S, Darre M, Hoagland T, Schreiber D, Khan MI, Donoghue A, Donoghue D and Venkitanarayanan K, 2012. Reduction of compounds trans-cinnamaldehyde and eugenol. *Applied and Environmental Microbiology* 78: 2981-2987. <https://doi.org/10.1128/AEM.07643-11>
- Kaoud HA, El-Babbly MA, El-Iraqi KG and Khalil MM, 2018. Prevalence of *Salmonella* spp. in some broiler farms in different Egyptian governorates. *Journal of Veterinary Medical Research* 25: 164-173. <https://doi.org/10.21608/jvmr.2017.43315>
- Kauffman G, 1974. Kauffman White Scheme. *Journal of Acta Pathologica Microbiologica Science* 61: 385.
- Kok T, Worswich D and Gowans E, 1996. Some serological techniques for microbial and viral infections. In: *Practical Medical Microbiology*. Lee J, Fraser A, Marmion B and Simmons A (eds), 14th Ed. Edinburgh, Churchill Livingstone, UK.
- Komoriya T, Hashimoto A, Shinozaki A, Inoue M and Kohno H, 2007. Study on partial purification of α -toxin produced from obligate anaerobe *Clostridium perfringens*. Report of the Research Institute of Industrial Technology, Nihon University 88: 1-11.
- Koneman EW, Auen SD, Dowell VR and Sommers HM, 1988. *Color Atlas and Textbook of Diagnostic Microbiology*. 2nd Ed. JB Lip Co, New York, USA.
- la Mora ZVD, Macías-Rodríguez ME, Arratia-Quijada J, Gonzalez-Torres YS, Nuño K and Villarruel-López A, 2020. *Clostridium perfringens* as foodborne pathogen in broiler production: Pathophysiology and potential strategies for controlling necrotic enteritis. *Animals* 10: 1-28. <https://doi.org/10.3390/ani10091718>
- Lee MD and Nolan KL, 2008. Chapter 3: Colibacillosis. In: *A Laboratory Manual for the Isolation and Identification of Avian Pathogen*. Zavala LD, Swayne DE, Glisson JR, Jack wood MW, Pearson JE and Reed WM (eds). 5th Ed, American Association of Avian Pathologists, USA.
- Malmarugan S, Boobalan A and Dorairajan N, 2012. Necrotic enteritis in broiler and layer farms in Tamil Nadu, India. *International Journal of Agro Veterinary and Medical Sciences* 6: 241-249. <https://doi.org/10.5455/ijavms.126>

- Mehmood K, Bilal RM and Zhang H, 2020. Study on the genotypic and phenotypic resistance of tetracycline antibiotic in *Escherichia coli* strains isolated from free ranging chickens of Anhui Province, China. *Agrobiological Records* 2: 63-68. <https://doi.org/10.47278/journal.abr/2020.014>
- Merati R, Temim S and Mohamed AAA, 2017. Identification and characterization of *Clostridium perfringens* isolated from necrotic enteritis in broiler chickens in Tiaret, Western Algeria. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 23: 595-601.
- Moawad AA, Hotzel H, Awad O, Tomaso H, Neubauer H, Hafez HM and El-Adawy H, 2017. Occurrence of *Salmonella enterica* and *Escherichia coli* in raw chicken and beef meat in northern Egypt and dissemination of their antibiotic resistance markers. *Gut Pathogens* 9: 57. <https://doi.org/10.1186/s13099-017-0206-9>
- Mridha D, Uddin MN, Alam B, Akhter AH, Islam SK, Islam MS, Khan MSR and Kabir SM, 2020. Identification and characterization of *Salmonella* spp. from samples of broiler farms in selected districts of Bangladesh. *Veterinary World* 13: 275-283. <https://doi.org/10.9775/kvfd.2017.17431>
- Muna EA, Salih MH, Zakia AM, Halima MO, Abeer AM, Ameera MM, Ali HO and Idris SB, 2016. Pathology of broiler chicks naturally infected with *Salmonella enteritidis* (*S. enteritidis*) and *Salmonella typhimurium* (*S. typhimurium*) during an outbreak in Sudan. *Journal of Scientific Research and Reports* 10: 1-8. <https://doi.org/10.9734/jsrr/2016/23431>
- Osman KM, Soliman YA, Amin ZMS and Aly MAK, 2012. Prevalence of *Clostridium perfringens* type A isolates in commercial broiler chickens and parent broiler breeder hens in Egypt. *Revue Scientifique et Technique (International Office of Epizootics)* 31: 931-941. <https://doi.org/10.20506/rst.31.3.2169>
- Palliyeguru MW, Rose SP and Mackenzie AM, 2010. Effect of dietary protein concentrates on the incidence of NE and growth performance of broiler chickens. *Journal of Poultry Science* 89: 34-43.
- Park JY, Kim S, Oh JY, Kim HR, Jang I, Lee HS and Ki YK, 2015. Characterization of *Clostridium perfringens* isolates obtained from 2010 to 2012 from chickens with necrotic enteritis in Korea. *Poultry Science* 94: 1158-1164. <https://doi.org/10.3382/ps/pev037>
- Popoff MY, 2001. Antigenic formulas of the *Salmonella* servars, WHO Collaborating Center for Reference and Research on *Salmonella*, Paris. 8th Ed. Porter RE Jr, 1998. Bacterial enteritides of poultry. *Poultry Science* 77: 1159-1165. <https://doi.org/10.1093/ps/77.8.1159>
- Rahman M, Samad M, Rahman MJ and Kabir S, 2004. Bacteriopathological studies on salmonellosis, colibacillosis and pasteurellosis in natural and experimental infections in chickens. *Bangladesh Journal of Veterinary Medicine* 2: 1-8. <https://doi.org/10.3329/bjvm.v2i1.1926>
- Reynolds DL, 2003. Multicausal enteric diseases. In: *Diseases of poultry* (ed. YM Saif), Iowa State University Press, Ames, IA, USA, pp: 1169-1171.
- Ronco T, Stegger M., Heidemann R, Sekse C, Bang A, Pohjanvirta T, Lilje B, Ulrike L, Skytt P and Pedersen K, 2017. Spread of avian pathogenic *Escherichia coli* ST117 O78:H4 in Nordic broiler production. *BMC Genomics* 18: 13. <https://doi.org/10.1186/s12864-016->
- Saha AK, Sufian MA, Hossain MI and Hossain MM, 2012. Salmonellosis in layer chickens: pathological features and isolation of bacteria from ovaries and inner content of laid eggs. *Journal of Bangladesh Agriculture University*, 10: 61-67. <https://doi.org/10.3329/jbau.v10i1.12095>
- Shah SA, Mir MS, Wani BM, Kamil SA, Goswami P, Amin U, Shafi M, Rather MA, Beigh AB, 2019. Pathological studies on avian pathogenic *Escherichia coli* infection in broilers. *Pharma Innovation* 8: 68-73.
- Sarba E, Kelbesa K, Bayu M, Gebremedhin E, Borena B and Teshale A, 2019. Identification and antimicrobial susceptibility profile of *Escherichia coli* isolated from backyard chicken in and around ambo, Central Ethiopia. *BMC Veterinary Research* 15: 85. <https://doi.org/10.1186/s12917-019-1830-z>
- Sarkar M, Ray JP, Mukhopadhyay SK, Niyogi D and Ganguly S, 2013. Study on *Clostridium perfringens* type A infection in broilers of West Bengal, India. *IIOAB Journal* 4: 1-3.
- Schlundt J, 2002. New directions in foodborne disease prevention. *International Journal of Food Microbiology* 78: 3-17.
- Sedeik ME, El-shall NA, AAM Elfeky, Abd El-Hack ME, Hussein EOS, Alowaimer AN and Swelum AA, 2019. Isolation, conventional and molecular characterization of *Salmonella* spp. From newly hatched broiler chicks. *AMB Express* 9: 136. <https://doi.org/10.1186/s13568-019-0821-6>
- Shoab M, Dasti JI, Shah MAA, Zafar MA, Hasan MU, Riaz A, Rehman SU and Khan MA, 2017. Salmonellosis in poultry, new prospects of an old disease: a review. *Pakistan Journal of Science* 69: 361-368.
- Slaoui M and Fiette L, 2011. Histopathology procedures: from tissue sampling to histopathological evaluation. *Methods in Molecular Biology* 691: 69-82. https://doi.org/10.1007/978-1-60761-849-2_4
- Soliman S, Seida AA, El-Fakar SZ, Youssef YI and El-Jakee J, 2018. *Salmonella* infection in broiler flocks in Egypt. *Bioscience Research* 15: 1925-1930.
- Su YC, Yu CY, Lin JL, Lai JM, Chen SW, Tu PC and Chu C, 2011. Emergence of *Salmonella* Enterica sero-var Potsdam as a major serovar in waterfowl hatcheries and chicken eggs. *Avian Disease* 55: 217-222. <https://doi.org/10.1637/9420-060910-Reg.1>
- Surra G, Dereje T and Chaluma N, 2018. Isolation and identification of *Escherichia coli*, *Salmonella* and *Pasteurella* from holding grounds of live-bird markets at Addis Ababa, Ethiopia. *African Journal of Microbiology Research* 12: 754-760.
- Uzal FA, Freedman JC, Shrestha A, Theoret JR, Garcia J, Awad MM, Adam SV, Moore RJ, Rood JI and McClane B, 2014. Towards an understanding of the role of *Clostridium perfringens* toxins in human and animal disease. *Future Microbiology* 9: 361-377. <https://doi.org/10.2217/fmb.13.168>
- van Asten AJAM, van der Wiel CW, Nikolaou G, Houwers DJ and Grone A, 2009. A multiplex PCR for toxin typing of *Clostridium perfringens*. *Veterinary Microbiology* 136: 411-412. <https://doi.org/10.1016/j.vetmic.2008.11.024>
- Wang J, Li J, Liu F, Cheng Y and Su J, 2020. Characterization of *Salmonella enterica* isolates from diseased poultry in northern China between 2014 and 2018. *Pathogens* 9: 95. <https://doi.org/10.3390/pathogens9020095>
- Williams RB, 2005. Intercurrent coccidiosis and necrotic enteritis of chickens: Rational, integrated disease management by maintenance of gut integrity. *Avian Pathology* 34: 159-180. <https://doi.org/10.1080/03079450500112195>
- Williams RB, Marshall RN, La Ragione RM and Catchpole J, 2003. A new method for the experimental production of necrotic enteritis and its use for studies on the relationships between necrotic enteritis, coccidiosis and anticoccidial vaccination of chickens. *Parasitological Research* 90: 19-26. <https://doi.org/10.1007/s00436-002-0803-4>
- Yasmin S, Nawaz M, Anjum AA, Ashraf K, Basra MAR, Mehmood A, Khan I and Malik F, 2020. Phytochemical analysis and in vitro activity of essential oils of selected plants against *Salmonella enteritidis* and *Salmonella gallinarum* of poultry origin. *Pakistan Veterinary Journal* 40: 139-144. <https://doi.org/10.29261/pakvetj/2019.110>