

A Convenient Synthesis of 1-Amino-4-methyl-4*H*-3-thia-4,5*a*,10-triazacyclopenta[*a*]fluoren-5-ones and Some New 2,3-Dihydro-1,3,4-thiadiazoles

Abdou O. Abdelhamid

Department of Chemistry, Cairo University, Giza, Egypt

Ahmed H. Elghandour

Department of Chemistry, Cairo University (Beni-Suef Branch), Beni-Suef, Egypt

Nora A. Rateb

Department of Chemistry, Cairo University, Giza, Egypt

Ahmed M. Awad

Department of Chemistry, Cairo University (Beni-Suef Branch), Beni-Suef, Egypt

1-amino-4-methyl-4H-3-thia-4,5a,10-triazacyclopenta-[a]fluoren-5-one and 6-methyl-6H-9H-thia-4b,6,9,11,12-pentaazaindeno[1,2-a]fluorene-5,8-dione derivatives were prepared from 2-methyl-1-oxo-3-thioxo-2,4,9b-trihydropyrimidino[1,6-a]benzimidazole-4-carbonitrile. Also, 2,3-dihydro-1,3,4-thiadiazoles were synthesized via a reaction of hydrazonoyl chlorides with 3-(methylamino)-2-substituted 3-thioxopropanenitrile. Structures of newly synthesized compounds were elucidated on the basis of elemental analyses, spectral data, and alternative methods synthesis whenever possible.

Keywords Halo ketones; isothiocyanate; nitrilimine; pyrimido[1,6-*a*]benzimidazole

INTRODUCTION

Isothiocyanates remain very important starting materials for the construction of heterocycles.¹ Several benzimidazoles are known² to possess antimicrobial, anti-inflammatory, and anticancer activities. Also, 1,3,4-thiadiazole and its derivatives have become a very useful compound in medicine, agriculture, and in many fields of technology.³ As

Received May 10, 2005; accepted August 29, 2005.


Address correspondence to Abdou O. Abdelhamid, Cairo University, Department of Chemistry, Faculty of Science, Giza 12613, Egypt. E-mail: Abdou_abdelhamid@yahoo.com

an extension of our study,^{4–9} we report herein the synthesis of 1-amino-4-methyl-4H-3-thia-4,5a,10-triazacyclopenta[*a*]fluoren-5-one and 2,3-dihydro-1,3,4-thiadiazole derivatives.

RESULTS AND DISCUSSION

The treatment of 2-(1-ethoxycarbonyl)benzimidazolylacetonitrile (**1**)¹⁰ with methyl isothiocyanate (**2**) in the presence of potassium hydroxide followed by acidification gave one isolable product according to TLC, which is formulated as 2-methyl-1-oxo-3-thio-1,2,3,4-tetrahydrobenzimidazo[1,2-*c*]pyrimidine-4-carbonitrile (**3**) (Scheme 1) based upon elemental analysis, spectral data, and chemical transformations.

Thus, the ¹H NMR spectrum of the product showed signals at $\delta = 3.72$ (s, 3H), 3.81 (s, 1H), 7.41–7.45 (m, 3H), and 8.20–8.24 (d, 1H). The ¹³C NMR showed signals at $\delta = 34$ (N–CH₃), 44 (CH₂), 115, 115, 116 (CN), 123, 123, 131, 139, 141, 150, and 196 (CO). Its IR spectrum revealed bands at 3076, 2987 (CH), 2217 (CN), 1706 (CO), and 1634 (C=N)




SCHEME 1

The treatment of **3** with chloroacetone in *N,N*-dimethylformamide in the presence of potassium hydroxide afforded 2-acetyl-1-amino-4-methyl-4H-3-thia-4,5a,10-triazacyclopenta[*a*]fluoren-5-one (**5a**) (Scheme 1). Structure **5a** was elucidated by elemental analysis and spectral data. Thus, the ¹H NMR spectrum showed signals at $\delta = 2.23$ (s, 3H), 3.45 (s, br., 2H), 7.38–7.49 (m, 2H), 7.73–7.77 (d, 1H), and 8.22–8.26 (1H). The ¹³C NMR spectrum showed signals at $\delta = 28$ (N–CH₃), 33 (CH₃CO),

115, 115, 123, 123, 129, 129 131, 136, 139, 141, 143, 159, and 191 (CO). Its IR spectrum revealed bands at 3444, 3322 (NH₂), 1710, 1635 (CO's), and 1621 (C=C). No absorption band between 2000–2300 was observed due to the absence of the CN group.¹¹ The reaction seemed to proceed through dehydrochlorination to give intermediate **4a**, which underwent cyclization via the addition of –SCH₂-hydrogens to a nitrile function to give the final product **5a**.

Similarly, compound **3** was reacted with each of phenacylbromide, ethyl chloroacetate, or chloroacetonitrile to give 1-amino-4-methyl-2-substituted 4H-3-thia-4,5a,10-triazacyclopenta[*a*]fluoren-5-one **5b–d**, respectively (Scheme 1).


Also, compound **3** was reacted with either 3-chloropentan-2,4-dione or ethyl 2-chloro-3-oxobutanoate in *N,N*-dimethylformamide containing potassium hydroxide to afford products identical (m.p., mixed m.p., and spectra) with **5a** and **5b**, respectively (Scheme 2).




SCHEME 2

Compound **5d** was reacted with each of formic acid and formamide under reflux to afford the corresponding 6-methyl-6H-7-thia-4b,6,9,11,12-pentaazaindeno[1,2-*a*]fluorene-5,8-dione (**12a**) and 8-amino-6-methyl-6H-7-thia-4b,6,9,11,12-pentaazaindeno[1,2-*a*]fluorene-5-one (**12b**), respectively (Scheme 3).

The treatment of methyl isothiocyanate (**2**) with 2-(cyanomethyl)-1-methylbenzimidazole (**13**)¹² in *N,N*-dimethylformamide containing potassium hydroxide afforded the non-isolable intermediate **14**, which converted to thioamide **15** by the acidification with hydrochloric acid (Scheme 4). The structure of **15** was confirmed by elemental analysis, spectral data, and chemical transformation. Its ¹H NMR spectrum



SCHEME 3




SCHEME 4

showed signals at $\delta = 2.18$ (s, 3H), 3.20 (s, 3H), 6.80 (s, br., 1H), 7.18–7.34 (m, 4H), and 14.08 (s, br., 1H).

Thus, the treatment of **15** with the appropriate hydrazoneyl halides^{13,14} **16a,b** afforded 2,3-dihydro-1,3,4-thiadiazoles **18a**¹⁵ and **18b**,¹⁶ respectively (Scheme 4). Structure **18** was confirmed on the basis of elemental analysis, spectral data, and alternative synthesis. The ¹H NMR spectrum of **18a** showed signals at $\delta = 1.4$ (t, $J = 7$ Hz, 3H), 3.0 (s, 3H), 4.4 (q, $J = 7$ Hz, 2H), and 7.12–8.0 (m, 9H). Its IR spectrum revealed bands at 2220 (CN) and 1713 (CO).

In the light of the foregoing results, the mechanism outlined in Scheme 4 seems to be the most plausible pathway for the formation of **18** from the reaction of **15** with **16**. The reaction involves an initial formation of thiohydrazone **17**, which undergoes intermolecular cyclization as soon as it is formed to yield **18** via the elimination of methylamine, and structures **19** and **20** were ruled out. Also, the treatment of methyl carbodithioate **21** with **16a** afforded a product identical in all respects (m.p., mixed m.p., and spectra) with **18a**.

Similarly, the treatment of 2-cyanomethylbenzothiazole (**23**) with methyl isothiocyanate (**2**) in *N,N*-dimethylformamide containing potassium hydroxide gave the non-isolable intermediate **24**, which reacted with each hydrazoneyl chloride **16a** and **16b** to give the corresponding 2,3-dihydro-1,3,4-thiadiazoles **25a** and **25b**, respectively (Scheme 5). Also, compound **26** was reacted with the appropriate hydrazoneyl chlorides **16a,b** in ethanolic triethylamine afforded products identical in all respects (m.p., mixed m.p., and spectra) with **25a,b**, respectively.



SCHEME 5

EXPERIMENTAL

All melting points were determined on an electrothermal apparatus and are uncorrected. IR spectra were recorded (KBr discs) on a Shimadzu FT-IR 8201 PC spectrophotometer. ^{13}C NMR and ^1H NMR spectra were recorded in $(\text{CD}_3)_2\text{SO}$ solutions on a Varian Gemini 300 MHz spectrometer, and chemical shifts are expressed in δ units using TMS as an internal reference. Elemental analyses were carried out at the Microanalytical Center of the Cairo University, Egypt.

The Synthesis of 2-Methyl-1-oxo-3-thioxo-2,4,9b-trihydropyrimidino[1,6-a]benzimidazole-4-carbonitrile (**3**)

A mixture of 2-(1-ethoxycarbonyl)benzimidazolylacetonitrile (**1**), methylisothiocyanate, and potassium hydroxide (5 mmol each) in *N,N*-dimethylformamide (15 mL) was stirred for 6 h. The reaction mixture was diluted with water (10 mL) and acidified with hydrochloric acid; then the resulting solid was collected and recrystallized from *N,N*-dimethylformamide to give **3** (Tables I and II).

The Synthesis of 1-Amino-4-methyl-2-substituted 4H-3-thia-4,5a,10-triazacyclopenta[*a*]fluoren-5-one Derivatives **5a–c**. General Procedure

Equimolar amounts of 2-(1-ethoxycarbonyl)benzimidazolylacetonitrile (**1**), methylisothiocyanate, and potassium hydroxide (5 mmol each) in *N,N*-dimethylformamide (15 mL) were stirred for 6 h. The appropriate chloroacetone (or 3-chloro-2,4-pentanedione), ethyl chloroacetate (or ethyl 2-chloro-3-oxobutanoate), or chloroacetonitrile (5 mmol) was added while stirring, and the reaction mixture was stirred for 2 h. The resulting solid was collected by filtration and recrystallized from *N,N*-dimethylformamide to give **5a–c** (Tables I and II).

The Synthesis of 6-Methyl-6H-7-thia-4b,6,9,11,12-pentaazaindeno[1,2-*a*]fluorene-5,8-dione (**12a**) and 8-amino-6-methyl-6H-7-thia-4b,6,9,11,12-pentaazaindeno[1,2-*a*]fluorene-5-one (**12b**)

Equimolar amounts of **5c** (0.5 g) was boiling in formic acid (10 mL, 99%) or formamide (2 mL) in *N,N*-dimethylformamide (10 mL) for 7 h. The reaction mixture was poured onto ice-cold water (40 mL); the resulting solid was collected and recrystallized from *N,N*-dimethylformamide to give **12a** and **12b**, respectively (Tables I and II).

TABLE I Characterization Data of the Newly Synthesized Compounds

Compound no.	M.P. °C (Solvent)	Color yield (%)	Mol. Formula (Mol. Wt.)	Elemental analysis [Calcd./Found (%)]			
				C	H	N	S
3	308–310	Colorless 55	C ₁₂ H ₈ N ₄ SO	56.24	3.15	21.86	12.51
	DMF		256.29	56.42	3.01	21.68	12.70
5a	272–273	Colorless 50	C ₁₅ H ₁₂ N ₄ SO ₂	57.68	3.87	17.94	10.27
	DMF		312.35	57.80	3.65	18.10	10.40
5b	241–243	Colorless 57	C ₁₆ H ₁₄ N ₄ SO ₃	56.13	4.12	16.36	9.37
	DMF		342.38	56.00	4.20	16.40	9.45
5c	243	Yellow 60	C ₂₀ H ₁₄ N ₄ SO ₂	64.16	3.77	14.97	8.56
	DMF		374.42	64.20	3.90	15.12	8.65
5d	334–336	Pale Brown 62	C ₁₄ H ₉ N ₅ SO	65.94	3.07	23.71	10.86
	AcOH		295.32	66.10	2.90	23.90	10.60
12a	>300	Colorless 57	C ₁₅ H ₉ N ₅ SO ₂	55.72	2.81	21.66	9.92
	DMF		323.34	55.60	2.90	21.80	10.20
12b	>300	Brown 52	C ₁₅ H ₁₀ N ₆ SO	55.89	3.13	26.07	9.95
	DMF		322.35	55.90	3.10	26.20	10.20
15	238–240	Brown 55	C ₁₂ H ₁₂ N ₄ S	58.99	4.95	22.93	13.12
	EtOH		244.32	58.80	4.70	22.80	13.10
18a	237–238	Yellow 60	C ₂₁ H ₁₇ N ₅ SO ₂	62.52	4.25	17.36	7.95
	EtOH		403.47	62.50	4.10	17.20	7.90
18b	274–276	Orange 57	C ₂₀ H ₁₅ N ₅ SO	64.33	4.05	18.75	8.59
	DMF		373.44	64.10	4.15	18.80	8.95
21	248–250	Yellow 82	C ₁₂ H ₁₁ N ₃ S ₂	55.15	4.24	16.08	24.54
	DMF		261.37	55.30	4.10	15.80	24.30
25a	240–242	Yellow 52	C ₂₀ H ₁₄ N ₄ S ₂ O ₂	59.10	3.47	13.78	15.78
	EtOH		406.49	59.20	3.80	13.80	15.80
25b	248–250	Yellow 59	C ₁₉ H ₁₂ N ₄ SO	60.62	3.21	14.89	17.03
	EtOH		373.44	60.50	3.10	14.90	17.20

The Synthesis of 3-(Methylamino)-2-(1-methylbenzoimidazole-3-thioxopropanenitrile (15)

Equimolar amounts of 2-(1-methylbenzimidazol-2-yl)ethanenitrile, methyl isothiocyanate, and potassium hydroxide (5 mmol) in *N,N*-dimethylformamide (15 mL) were stirred for 3 h at r.t. and then acidified with hydrochloric acid (1N). The resulting solid was collected and recrystallized from ethanol to give **15**.

The Synthesis of 2-(1-Methylbenzoimidazol-2-yl)-3-methylthio-3-thioxopropanenitrile (21)

Equimolar amounts of 2-(1-methylbenzimidazol-2-yl)ethanenitrile, carbon disulfide, and potassium hydroxide (5 mmol) in *N,N*-dimethylformamide (15 mL) were stirred for 3 h at r.t. Methyl iodide

TABLE II Spectra of Some Selected Synthesized Compounds

Compound no.	¹ H NMR and IR Spectra
3	¹³ C NMR: $\delta = 34, 44, 115, 115, 116, 123, 123, 131, 139, 141, 150, 196$. ¹ H NMR: $\delta = 3.72$ (s, 1H), 3.82 (s, 1H), 7.41–7.45 (m, 3H), and 8.20–8.24 (d, 1H). IR: 3076 (CH, aromatic), 2987 (CH, aliphatic), 2217 (CN), 1706 (CO), and 1634 (C=N).
5a	¹³ C NMR: $\delta = 28, 33, 115, 115, 123, 123, 129, 129, 131, 136, 139, 141, 143, 159, 191$. ¹ H NMR: $\delta = 2.23$ (s, 3H), 3.45 (s, br., 2H), 7.38–7.49 (m, 2H), 7.73–7.77 (d, 1H), and 8.22–8.26 (1H). IR: 3444, 3322 (NH ₂) and 1710, 1635 (CO's).
5b	¹³ C NMR: $\delta = 14, 61, 34, 109, 115, 115, 123, 123, 129, 130, 131, 138, 141, 142, 159, 161$. ¹ H NMR: $\delta = 1.37$ (t, 3H), 3.64 (s, 3H), 4.30 (q, 2H), 6.82 (s, br., 1H), 7.41–7.50 (m, 1H), 7.73–7.78 (d, 2H) and 8.33–8.38 (d, 1H). IR: 3449, 3339 (NH ₂), 3061 (CH), 2971 (CH), 1706, 1666 (CO's), and 1633 (C=N).
5c	3.64 (s, 3H), 6.82 (s, 2H) and 7.27–8.31 (m, 9H). IR: IR: 3444, 3322 (NH ₂), 2174 (CN), and 1710, 1635 (CO's)
5d	¹ H NMR: $\delta = 3.54$ (s, 3H), 7.19 (s, br., 2H), 7.42–7.54 (m, 2H), 7.77–7.80 (d, 1H), and 8.26–8.30 (d, 1H). IR: 3444, 2360 (NH ₂), 2172 (CN) and 1707, 1635 (CO's).
12a	¹ H NMR: $\delta = 3.70$ (s, 3H), 7.47–8.37 (m, 5H) and 12.82 (s, br., 1H). IR: 3450 (OH), 17407, 1662 (CO's), and 1625 (C=N).
12b	¹ H NMR: $\delta = 3.70$ (s, 3H), 3.06 (s, 1H), 6.94 (s, br., 2H), 7.34–8.37 (m, 2H), 7.70–7.74 (d, 1H), and 8.21–8.25 (d, 1H). IR: 3434, 3305 (NH ₂), 1732 (CO) and 1625 (C=N).
15	¹ H NMR: $\delta = 2.18$ (s, 3H), 3.20 (s, 3H), 6.80 (s, br., 1H), 7.18–7.34 (m, 4H) and 14.08 (s, br., 1H). IR: 3307 (NH), 2179 (CN), and 1625 (C=N).
18a	¹ H NMR: $\delta = 1.4$ (t, J = 7 Hz, 3H), 3.0 (s, 3H), 4.4 (t, J = 7 Hz, 2H), and 7.0–8.0 (m, 9H). IR: 2200 (CN) and 1710 (CO).
18b	¹ H NMR: $\delta = 2.6$ (s, 3H), 3.0 (s, 3H), and 7.0–8.0 (m, 9H). IR: 2200 (CN) and 1680 (CO).
21	¹ H NMR: $\delta = 2.25$ (s, 3H), 2.47 (s, 1H), 3.36 (s, 3H), 7.38–7.49 (m, 2H), 7.73–7.77 (d, 1H), 8.22–8.26 (1H). IR: 3327 (NH) and 2216 (CN).
25a	¹ H NMR: $\delta = 1.4$ (t, J = 7 Hz, 3H), 4.4 (t, J = 7 Hz, 2H) and 7.2–8.1 (m, 9H). IR: 2195 (CN) and 1720 (CO).
25b	¹ H NMR: $\delta = 2.5$ (s, 3H) and 7.2–8.1 (m, 9H). IR: 2188 (CN) and 1701 (CO).

(0.71 g, 0.32 mL) was added with stirring; the resulting solid was collected and recrystallized from *N,N*-dimethylformamide to give **21** as yellow crystals (Tables I and II).

The Synthesis of 2,3-Dihydro-1,3,4-thiadiazole Derivatives **18a,b** and **25a,b**

Method A

A mixture of the appropriate **15**, the appropriate hydrazoneoyl chlorides **16a,b** (5 mmol each), and triethylamine (0.75 mL, 0.005 mol) in ethanol (20 mL) was stirred for 30 min. The resulting product was collected and recrystallized from the proper solvent to give 2,3-dihydro-1,3,4-thiadiazoles **18a,b** and **24a,b**, respectively (Tables I and II).

Method B

An equimolar amount of the appropriate hydrazoneoyl chlorides **16a,b**, methylcarbodithioate **21** (or **26**¹⁷), and triethylamine (5 mmol, each) in ethanol (15 mL) was stirred at r.t. for 1 h. The resulting solid was collected and recrystallized from ethanol to give compounds **18a**, **18b**, **25a**, and **25b**, respectively.

REFERENCES

- [1] A. K. Mukerjee and R. Ashare, *Chem. Rev.*, **91**, 1 (1991).
- [2] A. V. Dolzhenko, W. Chui, and L. Chan, *J. Fluorine Chemistry*, **126**, 759 (2005).
- [3] D. I. Kornis, In *Comprehensive Heterocyclic Chemistry*, Vol. 4, A. R. Katritzky, C. W. Rees, and E. F. V. Scriven, Eds., p. 407 (Pergamon, UK, 1996).
- [4] A. O. Abdelhamid, H. F. Zohdi, and M. M. Zada, *Indian J. Chemistry*, **39B**, 202 (2002).
- [5] A. O. Abdelhamid, H. F. Zohdi, and M. M. Zada, *Indian J. Chemistry*, **40B**, 284 (2000).
- [6] A. O. Abdelhamid, M. M. M. Sallam, and S. A. Amer, *Heteroatom*, **12**, 468 (2001).
- [7] A. O. Abdelhamid, S. M. Abdelgawad, and S. F. El-Sharnobey, *Phosphorus, Sulfur, and Silicon*, **177**, 2699 (2002).
- [8] A. O. Abdelhamid, B. A. M. Abdel-Wahab, and A. A. Al-Atoom, *Phosphorus, Sulfur, and Silicon*, **179**, 601 (2004).
- [9] A. O. Abdelhamid, A. H. El-Ghandour, A. M. Hussien, and Y. H. Zaki, *Sulfur J. Chemistry*, **25**, 329 (2004).
- [10] J. Slouka, *Tetrahedron Lett*, 4007 (1968).
- [11] D. H. Williams and I. Fleming, Eds., *Spectroscopic Methods in Organic Chemistry* (2nd. ed.), (McGraw-Hill, UK, 1973).
- [12] R. A. B. Copeland and A. R. J. Day, *Am. Chem. Soc.*, **56**, 1072 (1943).
- [13] G. Favrel and J. Chrz, *Bull. Soc. Chim. France*, **41**, 1603 (1927).
- [14] N. F. Eweiss and A. Osman, *J. Heterocycl. Chem.*, **17**, 1713 (1980).

- [15] H. M. Hassaneen, A. E. Harhash, N. M. Abounada, T. A. Abdallah, and M. S. Algharib, *J. Chem. Res. (S)*, 194 (1993).
- [16] H. M. Hassaneen, A. S. Shawali, N. M. Elwan, and T. A. Abdallah, *Sulfur Lett.*, **15**, 103 (1992).
- [17] A. O. Abdelhamid, N. M. Rateb, and K. M. Dawood, *Phosphorus, Sulfur, and Silicon*, **167**, 251 (2000).