



# Biochemical and Anthropometric Nutritional Assessment in Children Infected with COVID-19: A Cross-sectional Study

Hoda Atef Abdelsattar Ibrahim<sup>1\*</sup>, Eatematd Helmy<sup>2</sup>, Aya Amin<sup>3</sup>, Dina Mahmoud<sup>4</sup>

<sup>1</sup>Pediatric Clinical Nutrition Unit, Department of Pediatrics, Faculty of Medicine, Cairo University, Cairo, Egypt; <sup>2</sup>Department of Pediatrics, El-Matria Teaching Hospital, General Organization for Teaching Hospitals and Institutes, Cairo, Egypt; <sup>3</sup>Department of Cancer Epidemiology and Biostatistics, National Cancer Institute, Cairo University, Cairo, Egypt; <sup>4</sup>Department of Clinical Pathology, Faculty of Medicine, Cairo University, Cairo, Egypt

## Abstract

**Edited by:** Ksenija Bogoeva-Kostovska  
**Citation:** Ibrahim HAA, Helmy E, Amin A, Mahmoud D. Biochemical and Anthropometric Nutritional Assessment in Children Infected with COVID-19: A Cross-sectional Study. OpenAccessMacedJMedSci.2022Jul24; 10(B):1818-1826. https://doi.org/10.3889/oamjms.2022.9782

**Keywords:** Malnutrition; COVID-19; Children; Hypokalemia; Obesity

\***Correspondence:** Hoda Atef Abdelsattar Ibrahim, Pediatric Clinical Nutrition Unit, Pediatric department, Faculty of Medicine, Cairo University, Egypt. E-mail: hodaibrahim424@cu.edu.eg

**Received:** 13-Apr-2022

**Revised:** 15-Jun-2022

**Accepted:** 14-Jul-2022

**Copyright:** © 2022 Hoda Atef Abdelsattar Ibrahim, Eatematd Helmy, Aya Amin, Dina Mahmoud

**Funding:** This research did not receive any financial support

**Competing Interests:** The authors have declared that no competing interests exist

**Open Access:** This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**BACKGROUND:** Severe acute respiratory syndrome has led to a pandemic of coronavirus disease 2019 (COVID-19). Malnutrition either biochemically or anthropometrically is a well-known risk factor for COVID-19 and may be the vice versa

**AIM:** The objectives of this study were to investigate the prevalence of malnutrition in children infected with COVID-19 through evaluating the nutritional biomarkers such as serum electrolytes, serum albumin, and hemoglobin together with the anthropometric assessment.

**METHODS:** A cross-sectional study that was conducted at El-Matria Teaching Hospital for all children admitted with confirmed COVID-19 for 6 months from February 1, 2021 to the end of July, 2021. Nutritional biochemical evaluation included serum electrolytes particularly the potassium and other nutritional biomarkers such as serum albumin and hemoglobin. Nutritional anthropometric evaluation depended on body mass index, the height/length, weight for length, and weight for height. The prevalence of malnutrition esp. hypokalemia was the main outcome.

**RESULTS:** Hypokalemia was present in 21.8% of the study participants. Other nutritional biomarkers were found as hyponatremia, hypocalcemia, hypophosphatemia, and hypomagnesemia were detected in 49.1%, 38.2%, 21.8%, and 34.5% of the study subjects, respectively. Anthropometric malnutrition was present in most of the enrolled children with COVID-19 in the study (65.5 % [n = 36]) through which overweight and obese children occupied a greater percentage.

**CONCLUSION:** Malnutrition either biochemically or anthropometrically could be linked to COVID-19 in children. COVID-19 could have negative outcomes on the nutritional status such as electrolytes disturbances. Both malnutrition and COVID-19 are considered synergistic associations

## Introduction

The novel coronavirus severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is resulting in a current worldwide pandemic of coronavirus disease 2019 (COVID-19). It has become a globe health problem and disturbed the main health determinants [1], [2], [3]. Nutritional status has been utilized as resilience toward the destabilization during the COVID-19 pandemic. Optimal nutrition can improve the immune system; thus, a sustainable approach to survive in the current context is to reinforce the immune system [4], [5].

Malnourished children are among the reported patients with worst outcomes and higher morbidities and mortalities during COVID-19 pandemic. Diagnosis, prevention, and treatment of malnutrition should therefore be routinely involved in the management of children with COVID-19 to improve the short and long-term prognoses [6], [7], [8], [9].

Childhood malnutrition is recognized to be a worldwide concern as it is problematical with poor growth and declined educational outcomes of children and can have undesirable effects during the adulthood [10], [11]. It is among the most serious health matters facing the developing countries especially Egypt. Malnutrition does not only affect the health status of children, but it also affects the economic growth as described by a former study, in which a robust statistically significant negative association between the economic growth and outcomes of child malnutrition, namely, underweight, wasting, and stunting in Egypt for 1992–2008 was found [12]. In children, the most serious risks resulted from COVID-19 crisis are not those of the COVID-19 itself, but its associated damages. These can include insufficient nutrition statuses with a risk of both underweight and overweight [13].

Particularly, COVID-19 can be associated with vomiting and diarrhea affecting the child's food intake and absorption [14]; therefore, a good nutritional status is a healthy benefit for children at risks for severe COVID-19.

In a recent review about potential interventions for pandemics, authors recommended that the nutritional status of infected patients should be assessed before management with the general treatments [15].

Malnutrition states excess, deficiencies, or imbalances in a child's intake of energy and/or nutrients. Malnutrition includes three main groups. First group is the undernutrition group that addresses wasting, stunting, and underweight. Second group is the micronutrient-related malnutrition group that addresses the micronutrient deficiencies (a lack of important minerals or vitamins) or micronutrient excess. Third group is the overweight and obesity group [16]. A deficiency of micronutrients can lead to weakened immune responses resulting in an improper cytokines release, alterations in the secretory antibody response, and the antibody affinity. Micronutrients deficiency can be treated by proper nutritional supplements, prescribed after assessing the child's nutritional status [17].

One of the major micronutrients affected is the potassium electrolyte. Correction of hypokalemia is a challenge due to the continuous potassium loss from kidneys that results from the degradation of angiotensin-converting enzyme 2 (ACE2). The high prevalence of hypokalemia among children with COVID-19 proposes the presence of disordered activity of the rennin-angiotensin system that increases as a result of diminished counteractivity of ACE2 that is bound by SARS-CoV-2 [18].

The European Society for Clinical Nutrition and Metabolism has set recommended guidelines aiming at providing experts statements and applied recommendations for the nutritional management of COVID-19 patients. One of these recommendations states that identification of malnutrition risks should be an initial step in the general evaluation of all COVID-19 patients considering that obesity should be screened and investigated as it is a great comorbidity [6].

There are hypothesized links between obesity and the COVID-19 existence. It is well-recognized that dysbiosis has a strong association with the development of obesity and metabolic syndrome. Dysbiosis is characterized by the low diversity of microbes including a low abundance of *Bifidobacterium* spp., *Lactobacillus* spp., and *Faecalibacterium prausnitzii*. Consequently, children experiencing these disorders are much more prone to COVID-19 infections, primarily due to the existing disturbances within their gut microbiota [19]. An extended link is the link of dysbiosis to the malnutrition. To illustrate, altered bacterial interactions – seen in dysbiosis – can result in reduced productions of several essential amino acids with further development of malnutrition [20].

The objective of our study was to assess the pattern of malnutrition focusing on the potassium

and body mass index (BMI), respectively. Besides, addressing the prevalence of malnutrition in these children was another objective.

## Methods

### *The current study design*

This study was a cross-sectional analytical study.

### *Criteria of the infected children in the study*

#### *Inclusion criteria*

Admitted children within the first 2 days in El-Matria Teaching Hospital along a period of 6 months-both males and females-with confirmed COVID-19, and whose parents or legal guardians approved to be enrolled in the study.

#### *Exclusion criteria*

Children who were not confirmed to be COVID-19 and whose parents or caregivers did not approve to be enrolled in the study. In addition, admitted children with periods of admission which was more than 2 days were excluded to avoid bias that could be produced from the adverse outcomes of hospitalization as the child may become malnourished during the hospital stay due to a poor appetite.

### *Sample technique*

A convenient sample of all pediatrics with COVID-19 (n = 55) who were admitted at El-Matria Teaching hospital was recruited for the current study along the study period.

### *Sample size estimation*

The primary objective of the present study was to detect the prevalence of biochemical disturbances in COVID-19 children, especially hypokalemia which is the one of the major and most serious complications of COVID-19. De Carvalho *et al.* reported that hypokalemia was found in 15.1% COVID-19 patients [21]. Using Epi Info™ software [22], and by setting an acceptable margin of error of  $\pm 0.10$  and confidence level of 95%, the minimum required that sample size for the present study was 49 patients

### Operational definitions

1. Biochemical nutritional levels: Definitions of normal, deficiency, and excess were identified using the reference ranges that differ according to the age and sex
2. Anthropometric nutritional assessment using the World Health Organization (WHO) charts:
  - A. Overweight: When the weight for the length and the weight for the height in birth to 2 years and 2–5 years, respectively, is > the mean by two standard deviations (SDs) of the WHO Child Standards for growth or BMI z score is >1 SD of the WHO Child Standards for BMI 5–19 years charts.
  - B. Obesity: When the weight for the length and the weight for the height in birth to 2 years and 2–5 years, respectively, is > the mean by three SDs of the WHO Child Standards for growth or BMI z score is >2 SD of the WHO Child Standards for BMI 5–19 years charts
  - C. Wasting: When the weight for the length and the weight for the height in birth to 2 years and 2–5 years, respectively, is < the mean by two SDs of the WHO Child Standards for growth or BMI z score is <2 SD of the WHO Child Standards for BMI 5–19 years charts.
  - D. Underweight: When the weight for age is < the mean by two SD of the WHO Child Standards for growth.
    - WHO recommends BMI as the best indicator or measure after the age 10 as the age of puberty is variable. Tracking the weight alone is not recommended. Underweight and wasting among the adolescents (10–19 year) are defined as a BMI-for age  $\leq 2$  Z-scores of a reference.
    - Weight-for-age represents body weight relative to the child's age. This indicator addresses whether a child is underweight or severely underweight, but it is not used to address a child as overweight or obese [23].
  - E. Stunting: When the length/height for age is < the mean by two SDs of the WHO Child Standards for growth.
3. Confirmation of the diagnosis of COVID-19 was done by a positive reverse transcription-PCR (RT-PCR) targeting the different genes of COVID-19 on nasopharyngeal swabs.

### Ethical statement

The study was revised and approved by the Scientific Ethical Research Committee of General Organization for Teaching Hospitals and Institute (GOTHI), ethical clearance number HM000139, and the study was done in accordance with the laws for the human research. An informed consent was obtained.

### Statistical methods

Statistical calculations were conducted by means of SPSS (Statistical Package for the Social Science; SPSS Inc., Chicago, IL, USA) version 25. Quantitative data were tested for normality through exploring the variable histogram, Q-Q plots, mean, median, SDs, coefficient of variation, and skewness as well as performing the normality tests (Kolmogorov–Smirnov and Shapiro–Wilk tests). Normally distributed numerical variables were described in terms of mean, SD, and range. Non-normally distributed quantitative variables were statistically described as median, interquartile range, and range. Categorical variables were summarized as numbers and percentages.

### Efforts to address and avoid potential sources of bias

1. This study was carried out on the admitted children in the first one or 2 days to avoid the impact of the prolonged hospital stay due to the possible malnutrition or weight loss that could occur due to the possible appetite loss.
2. Pediatrics with associated chronic illnesses were excluded as they might have additional external reasons for malnutrition due to the chronic disease effect.

## Results

Fifty-five children were enrolled. Regards the demographic criteria, male patients represented 52.7 % of the study group (n=29). Female patients represented 47.3 % of the study group (n=26). Male-to-female ratio was 1.1:1 (Table 1).

**Table 1: Demographic criteria of the study participants**

Characteristics	Number (n = 55)	Percent
Age groups (years)		
<2	28	50.9
2–<6	13	23.6
6–13	14	25.5
Median (IQR)*	1.5 (0.5–6)	
Range	0.1–13	
Gender		
Male	29	52.7
Female	26	47.3
Male: Female ratio	1.1	

\*IQR: Interquartile range.

Malnutrition was evaluated in the enrolled children. Biochemical nutritional assessment was dependent on the electrolytes levels (Table 2). Moreover, other biochemical nutritional evaluations included serum albumin (Table 3) and hemoglobin (Table 4).

Hypokalemia was observed in 21.8% (n = 12), while hyperkalemia was observed in 5.5% (n = 3) of the study groups. About 72.2% of the study participants (n = 40) had normal serum potassium. Other electrolytes

observed were sodium, calcium, magnesium, and phosphorus.

**Table 2: Electrolytes assessment of the study participants**

Electrolytes	Number (n = 55)	Percent	95% CI**
<b>K</b>			
Hypokalemia	12	21.8	11.8–35.0
Normal K level	40	72.7	59.0–83.9
Hyperkalemia	3	5.5	1.1–15.1
Mean ± SD*	4.1 ± 0.8		3.9–4.3
Range	2.3–5.8		
<b>Na</b>			
Hyponatremia	27	49.1	35.4–62.9
Normal Na level	26	47.3	33.7–61.2
Hypernatremia	2	3.6	0.04–12.5
Mean ± SD*	134.35 ± 6.5		132.6–136.1
Range	111–156		
<b>Ca</b>			
Hypocalcemia	21	38.2	25.4–52.3
Normal Ca level	34	61.8	47.7–74.6
Mean ± SD*	1.1 ± 0.2		1.1–1.2
Range	0.4–1.4		
<b>Mg</b>			
Hypomagnesemia	19	34.5	22.2–48.3
Normal Mg level	31	56.4	56.4–42.3
Hypermagnesemia	5	9.1	3.0–20.0
Mean ± SD*	1.8 ± 0.3		1.8–1.9
Range	1.4–2.5		
<b>PO<sub>4</sub></b>			
Hypophosphatemia	12	21.8	11.8–35.0
Normal PO <sub>4</sub> level	43	78.2	65.0–88.2
Mean ± SD*	4.8 ± 0.8		4.5–5.0
Range	3.1–6.6		

\*SD: Standard deviation, \*\*95% CI: Confidence interval.

Hyponatremia was detected in 49.1% (n = 27), while hypernatremia was observed in 3.6% (n = 2) of the study groups. About 47.3% of the children in the study (n = 26) had normal serum sodium. As regards the calcium level, most of the enrolled pediatrics with COVID-19 (61.8% of the study participants [n = 34]) had normal serum calcium. Only 38.2% (n = 21) of the study participants had hypocalcaemia.

**Table 3: Albumin assessment in the study group**

Albumin	Number (n = 55)	Percent	95% CI**
Hypoalbuminemia	41	74.5	61.0–85.3
Normal albumin level	14	25.5	14.7–39.0
Mean ± SD*	3.4 ± 0.7		3.2–3.5
Range	1.9–5.0		

\*SD: Standard deviation, \*\*95% CI: Confidence interval

Another affected electrolyte in the course of COVID-19 infection was the phosphorus. To illustrate, hypophosphatemia was observed in 21.8% (n = 12) in the infected children. However, 78.2% of the study participants (n = 43) had normal serum phosphorus.

Levels of magnesium in the serum were also noticed. Hypomagnesemia was observed in 34.5% (n = 19), while hypermagnesemia was observed in 9.1% (n = 5) of the study groups. About 56.4% of the study participants (n = 31) had normal serum magnesium.

As serum albumin is an important nutritional biomarker, it was included in the evaluation. Interestingly, it was affected in most of the cases. Hypoalbuminemia was present in 74.5 % of the study groups. (n = 41). Only 25.5 % (n = 14) of the enrolled children had normal serum albumin.

**Table 4: Anemia in children with COVID-19**

Hb level	Number (n = 55)	Percent	95% CI**
Anemia	43	78.2	65.0–88.2
Normal Hb level	12	21.8	11.8–35.0
Mean ± SD*	11.0 ± 1.6		10.6–11.4
Range	6.4–15.4		

\*SD: Standard deviation, \*\*95% CI: Confidence interval

Another vital nutritional biomarker is the hemoglobin. Therefore, it was evaluated in our current study. The results were exciting. To explain, anemia was a result in 78.2% (n = 43) in children with COVID-19, enrolled in the present study.

**Table 5: Anthropometric nutritional assessment of the study participants**

Malnutrition	Number (n = 55)	Percent*
Malnutrition	36	65.5%
Overweight	10	18.2
Obese	10	18.2
Underweight	4	7.3
Wasted	11	20.0
Stunted	4	7.3

\*Some cases showed more than one malnutrition disorder

As a trial to complete the nutritional assessment in the course of COVID-19 infection for the admitted children, anthropometric measures were addressed. About 65.5% (n = 36) of the children were anthropometrically malnourished (Table 5). In more depth, types of this malnutrition were identified. Overweight and obesity represented 55.6% (n = 20) of all cases of malnutrition, wasted children with COVID-19 were 30.6% (n = 11), underweight children were 11.1% (n = 4), and stunted children were 11.1% (n = 4) (Table 6).

**Table 6: Percentage of each type malnutrition in relation to all malnourished cases enrolled**

Malnutrition	Number (n = 55)	Percent
Yes	36	65.5
No	19	34.5
Type of malnutrition	Number (n = 36)	Percent*
Overweight or Obese	20	55.6
Underweight	4	11.1
Wasted	11	30.6
Stunted	4	11.1

\*Some cases showed more than one malnutrition disorder.

## Discussion

Balanced nutritional status during the course of any illness is important for improving the health outcomes [24], [25]. Indeed, malnutrition is a great burden in Egypt with a considerable prevalence of short stature, underweight, overweight, and obesity [26]. However, our study did not only include malnourished children, but it also included the healthy ones. This helps address the role of the nutritional status in the course of COVID-19 infection. Therefore, bias that can develop from the prevalence of malnutrition in Egypt that can result in expressing the malnutrition rate in it as the malnutrition rate in our study can be slightly avoided. Moreover, the estimated prevalence of malnutrition in Egypt is much lower than the estimated prevalence of malnutrition in our study addressing the impact of COVID-19 on the nutritional state in the present study [26]. In addition, our study enrolled the biochemical nutritional assessment, not only the anthropometric one.

Male patients occupied the higher percentage in our study. This finding agrees with a previous study,



in which it was found that, of the 48 pediatrics admitted with COVID-19, 25 (52%) were male [27]. This gender factor, besides the higher incidences in males for most of the diseases, might correlate with a shorter life expectancy in males – which is a general demographic fact – compared to females in the world [28], [29]. Another hypothesis of this gender difference is that males are more vulnerable to infections, whereas females have a stronger antigenic response to infections and vaccines at the cost of the higher prevalence of autoimmune diseases [30].

Malnutrition was present in a greater percent in our study. This finding was consistent with the results of one previous study, which indicated that malnourished patients or patients at risks of malnutrition are at a risk of acquiring COVID-19 [24]. Significantly, regions with an increased burden of malnutrition have a greater risk of acquiring COVID-19 as stated in a former study. That study showed increased risks of fatal COVID-19 at populations with increased indicators of malnutrition [31]. Malnutrition can result in immunosuppression – which further enhances acquiring COVID-19 – through different means, including the contribution of leptin and the hypothalamic-pituitary-adrenal axis. Malnutrition decreases leptin concentrations and rises serum levels of stress hormones, that is, glucocorticoids. Therefore, it is likely that the hypothalamic-pituitary-adrenal axis occupies a crucial role in malnutrition-related immune deficiency [32].

Children infected with COVID-19 are at greater risks for fluid and electrolyte disturbances and complications. This was figured by a former article, in which hyponatremia, hypokalemia, and hypocalcemia were the most common electrolytes disorders detected in COVID-19 patients [33].

Hypokalemia was an electrolyte disorder in a considerable percentage of children with COVID-19 in our study. In contrast to the general people, COVID-19 children can experience three risk factors for hypokalemia. First, respiratory alkalosis due to the hypoxia-driven hyperventilation that can stimulate transcellular shifts with augmented intracellular uptakes. Second, anorexia, as a result of continuative use of face masks can result in a decrease in the potassium intake. Third, diarrhea due to the cytopathic effect of COVID-19 on the gastrointestinal cells may be a recurrent cause of potassium losses from the gastrointestinal tract [34].

Another investigated cause of hypokalemia is through the angiotensin system. To illustrate, the virus invades the cells through the ACE2 on the membrane. Angiotensin converting enzyme 2 is the main counter-regulatory mechanism for the key axis of the rennin angiotensin system (RAS) that is important in the control of the blood pressure and electrolyte balance. COVID-19 binds ACE2 and augments the degradation of ACE2 and, so, decreases the counteraction of ACE2 on RAS. This results in increasing the reabsorption of

sodium and water, thus increasing blood pressure and excretion of potassium that leads to hypokalemia [18].

Interestingly, there was a considerable percentage of hyperkalemia in our study. Furin has a role in this disorder. To explain, furin is required for cleaving both COVID-19 spike protein and epithelial sodium channels (ENaC) subunits. As the furin is hijacked by COVID-19, the reduced activity of ENaC would be anticipated, that causes retention of potassium ions with subsequent hyperkalemia [35].

From the perspective that the occurrence of hypokalemia is greater than hyperkalemia in COVID-19 children in our study, it seems that mechanisms of hypokalemia predominated.

Hyponatremia occupied the higher percentage of the sodium imbalance in the present study. This result is similar to the result of another study, in which it was found that sodium imbalance disorders, particularly hyponatremia, are a common abnormality among COVID-19 patients [36]. The pathophysiological mechanisms of hyponatremia in COVID-19 patients are different, including syndrome of inappropriate antidiuretic hormone secretion, gastrointestinal loss of sodium ions, or lowered sodium ion intakes. Hyponatremia may also be a negative predictive factor in COVID-19 patients [37].

A small percentage of the study group had hypernatremia. This is probably due to the increased angiotensin II activity secondary to the SARS-CoV-2-induced downregulation of ACE2 receptors as stated by a previous study [38].

In the context of higher percentage of hyponatremia encountered in the study, it appears that the mechanisms of hyponatremia overcame those of hypernatremia in the enrolled children.

Magnesium has a strong link to the immune system and the immunological functions are disturbed in case of magnesium deficiency. It has been reported that free basal magnesium concentration has an essential role in regulating the cytotoxic immune function. By that, intracellular free magnesium concentration contributes considerably to the antiviral immunity. Therefore, decreased resistance against infection with COVID-19 in case of magnesium deficiency can be assumed [39]. This may be the cause why hypomagnesemia occupied a higher percentage among the magnesium imbalance in our study. In a preceding study, it was concluded that magnesium should be observed and, in case of the imbalanced magnesium homeostasis, an appropriate nutritional regimen or supplementation could contribute to protect against SARS-CoV-2 infection, reduce severity of COVID-19 infection, and enhance the recovery after the acute phase [40], [41].

A fewer percentage of the enrolled children was noticed to have hypermagnesemia. A link between hypermagnesemia among patients admitted with COVID-19 and the increased mortality has been identified by a prior research. Although the precise

mechanism of this relationship remains unclear, this result potentially represents increased cell turnover and the severity of illness that is frequently associated with more severe forms of acute kidney injuries [42]. In fact, hypomagnesemia could be a predisposing factor for COVID-19 infection, while hypermagnesemia could be a result found in the severe COVID-19 infection form. A detailed analysis of magnesemia and the associated factors in a cohort of 300 patients was performed. Among these, 48% had a magnesemia below 0.75 mmol/L, involving 13% of severely hypomagnesemic patients (0.65 mmol/L)<sup>1</sup>; on the other hand, a fewer proportion of patients (9.6%) was hypermagnesemic (>0.95 mmol/L). In this report, in moderate cases of COVID-19, serum magnesium concentrations were considerably lower and the prevalence of hypomagnesemia was significantly higher than in the critical cases, while the prevalence of hypermagnesemia was significantly increased in the ICU cases [43].

It was recommended that more care and medical attention should be provided to COVID-19 patients with hypophosphatemia at the admission as hypophosphatemia is greatly linked to increased mortalities in COVID-19 patients [44]. Hypophosphatemia observed in the children with phosphorus imbalance in our study that could be the result of malnutrition, extensive energy consumption and uncontrolled losses from the intestinal tract and kidneys, and respiratory alkalosis [45], [46], [47], [48]. Besides, hypocalcemia observed in the present study could be a direct effect of COVID-19 or could be the consequence of an imbalance in parathyroid hormone and/or 25-hydroxyvitamin D. In general, calcium is needed for the virus structure formation, its entry, its replication, and release [49].

Hypoalbuminemia can be considered as a severity indicator of epithelial–endothelial damage in COVID-19 children. There are suspicions that pulmonary capillary leak syndrome has a significant role in the pathogenesis of COVID-19 infection and could be a potential therapeutic target [50]. Our finding is very near to a recent report that pointed to reduced albumin levels in pediatrics with mild/moderate COVID-19 [51]. Importantly, hypoalbuminemia is a negative acute phase reactant that has been associated with the inflammatory response and undesirable outcomes in the infectious diseases such as COVID-19. Serum albumin concentration can address children who are at high risks of developing potential life-threatening conditions and death, improving risk stratification in COVID-19 children [52]. Another retrospective preceding research has found that hypoalbuminemia at the presentation independently increased the risks of deaths in COVID-19 patients by at least 6-fold and recommended evaluation of the potential therapeutic value of albumin in COVID-19 patients [53]. Similarly, a former study has postulated the possible value of the nutritional support therapy to patients with hypoalbuminemia in the course of COVID-19 infection as that study has yielded that hypoalbuminemia was

common in COVID-19 patients and might be the result of a combination of inflammation and malnutrition in these patients. These patients with hypoalbuminemia tend to have more severe clinical courses and more abnormal results of the biochemical tests that may lead to poorer clinical outcomes. Hypoalbuminemia might indicate a poorer prognosis and the necessity for more diverse therapies [54]. Furthermore, hypoalbuminemia can guide the clinicians in the risk stratification of COVID-19 patients as concluded by a prior article, in which authors demonstrated the link between hypoalbuminemia and the severe COVID-19 infection. A low albumin serum level can potentially result in an early recognition of severe courses and assist clinicians in making informed stratifications and subsequent decisions for their patients [55].

According to the conclusion of a previous research, the BMI of patients with COVID-19 was significantly higher than those without [56]. This conclusion goes in line with the current study that yielded a high prevalence of overweight and obesity among those of malnourished children with COVID-19. Our study is further closely located to a previous report that indicated increased risks of acquiring COVID-19 infection among pediatrics with medical comorbidities and certain disorders such as obesity [57]. Obesity should be recruited as individual risk factors for higher COVID-19 severity and mortality outcomes [58]. Obesity is a well-known risk factor for acquiring COVID-19 and infections in general [59]. The regulated expression of miRNAs (microRNA) is crucial in the immunomodulation and more than 23 circulating miRNAs are deregulated in obese individuals. These miRNAs have valuable effects on the immune cell development, the T lymphocyte generation, the lipid metabolism, and the macrophage pro-inflammatory responses [60].

## Conclusion

The present study has identified the prevalence of the biochemical and the anthropometrical malnutrition in pediatrics infected with COVID-19. Large-scale prospective studies are needed to prove further hypotheses underlying different pathophysiological mechanisms that link COVID-19 infection and the malnutrition to each other's. Timely diagnosis and treatment of children with malnutrition or those at nutritional risks are recommended.

## Data Availability

The datasets analyzed during the present study are available from the corresponding author on reasonable request.

## Authors' Contributions

Hoda Atef Abdelsattar Ibrahim., the main author: Conceptualization, visualization, methodology, interpretation of data, writing—original draft preparation, reviewing and editing, software and validation, supervision; Eatematd Helmy: Acquisition of data; Aya Ahmed Amin.: Formal analysis; Dina Mahmoud Nabil: Investigation; all the authors approved for the final version.

## References

- Mallah SI, Ghorab OK, Al-Salmi S, Abdellatif OS, Tharmaratnam T, Iskandar MA, *et al.* COVID-19: Breaking down a global health crisis. *Ann Clin Microbiol Antimicrob.* 2021;20(1):35. <https://doi.org/10.1186/s12941-021-00438-7> PMID:34006330
- Raman S, Harries M, Nathawad R, Kyeremateng R, Seth R, Lonne B, International society for social P, child health CWG. Where do we go from here? A child rights-based response to COVID-19. *BMJ Paediatr Open.* 2020;4(1):e000714. <https://doi.org/10.1136/bmjpo-2020-000714> PMID:32577537
- Ibrahim HA, Mahmoud D. Blood groups as biomarkers for Covid-19 in pediatrics review article. *Annal Roman Soc Cell Biol.* 2021;25(6):19733-40. Available: from <https://www.annalsofscb.ro/index.php/journal/article/view/9837>
- Aman F, Masood S. How nutrition can help to fight against COVID-19 Pandemic. *Pak J Med Sci.* 2020;36(COVID19-S4):S121-3. <https://doi.org/10.12669/pjms.36.COVID19-S4.2776> PMID:32582329
- López-Mejía L, Núñez-Barrera I, Bautista-Silva M, Vela-Amieva M, Guillen-Lopez S. Nutrition therapy in children with COVID-19. *Acta Pediatr Mex.* 2020;41(4):109-20. <https://doi.org/10.18233/apm41no4s1pps109-s1202059>
- Barazzoni R, Bischoff SC, Krznaric Z, Pirlich M, Singer P, *et al.* Endorsed by the ESPEN council, espen expert statements and practical guidance for nutritional management of individuals with sars-cov-2 infection. *Clin Nutr.* 2020;39(6):1631-8. <https://doi.org/10.1016/j.clnu.2020.03.022> PMID:32305181
- Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, *et al.* ESPEN guideline on clinical nutrition in the intensive care unit. *Clin Nutr.* 2019;38(1):48-79. <https://doi.org/10.1016/j.clnu.2018.08.037> PMID:30348463
- Gomes F, Schuetz P, Bounoure L, Austin P, Ballesteros-Pomar M, Cederholm T, *et al.* ESPEN guideline on nutritional support for polymorbid internal medicine patients. *Clin Nutr.* 2018;37(1):336-53. <https://doi.org/10.1016/j.clnu.2017.06.025> PMID:28802519
- Volkert D, Beck AM, Cederholm T, Cruz-Jentoft A, Goisser S, Hooper L, *et al.* ESPEN guideline on clinical nutrition and hydration in geriatrics. *Clin Nutr.* 2019;38(1):10-47. <https://doi.org/10.1016/j.clnu.2018.05.024> PMID:30005900
- Ibrahim HA, Abdel-Raouf R, Zeid AS, Elsebaie EH, Abdalaleem S, Amin AA, Aboulghar H. Development of a simple and valid nutrition screening tool for pediatric hospitalized patients with acute illness [version 1; peer review: 2 approved, 1 approved with reservations, 1 not approved]. *F1000Research* 2021, 10:173 <https://doi.org/10.12688/f1000research.51186.1>
- Ibrahim HA, Nasr RA, Salama AA, Amin AA. Childhood malnutrition and hypo mineralized molar defects; a cross sectional study, Egypt [version 2; peer review: 3 approved]. *F 1000Research* 2022, 10:1307 <https://doi.org/10.12688/f1000research.74557.2>
- Rashad AS, Sharaf MF. Economic growth and child malnutrition in Egypt: New evidence from national demographic and health survey. *Soc Indic Res.* 2018;135(2):1-27. <https://doi.org/10.1007/s11205-016-1515-y>
- Zemrani B, Gehri M, Masserey E, Knob C, Pellaton R. A hidden side of the COVID-19 pandemic in children: The double burden of under nutrition and over nutrition. *Int J Equity Health.* 2021;20(1):44. <https://doi.org/10.1186/s12939-021-01390-w> PMID:33482829
- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, *et al.* Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: A descriptive study. *Lancet.* 2020;395(10223):507-13. [https://doi.org/10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7)
- Short KR, Kedzierska K, Van de Sandt CE. Back to the future: Lessons learned from the 1918 influenza pandemic. *Front Cell Infect Microbiol.* 2018;8:343. <https://doi.org/10.3389/fcimb.2018.00343> PMID:30349811
- Available from: <https://www.who.int/news-room/fact-sheets/detail/malnutrition> [Last accessed on 2021 Jun 09].
- Pallath MM, Ahiwar AK, Tripathi SC, Asia P, Sakarde A, Gopal N. COVID-19 and nutritional deficiency: A review of existing knowledge. *Horm Mol Biol Clin Investig.* 2021;42(1):77-85. <https://doi.org/10.1515/hmbci-2020-0074> PMID:33544528
- Chen D, Li X, Song Q, Hu CH, Su F, Dai J, *et al.* Assessment of hypokalemia and clinical characteristics in patients With COVID-19 in Wenzhou, China. *JAMA Netw Open.* 2020;3(6):e2011122. <https://doi.org/10.1001/jamanetworkopen.2020.11122> PMID:32525548
- Vodnar DC, Mitrea L, Teleky BE, Szabo K, Calinoiu LF, Nemes SA, *et al.* Coronavirus disease (COVID-19) caused by (SARS-CoV-2) infections: A real challenge for human gut microbiota. *Front Cell Infect Microbiol.* 2020;10:575559. <https://doi.org/10.3389/fcimb.2020.575559> PMID:33363049
- Kumar M, Ji B, Babaei P, Das P, Lappa D, Ramakrishnan G, *et al.* Gut microbiota dysbiosis is associated with malnutrition and reduced plasma amino acid levels: Lessons from genome-scale metabolic modeling. *Metab Eng.* 2018;49:128-42. <https://doi.org/10.1016/j.ymben.2018.07.018> PMID:30075203
- De Carvalho H, Richard MC, Chouihed T, Goffinet N, Le Bastard Q, Freund Y, *et al.* Electrolyte imbalance in COVID-19 patients admitted to the emergency department: A case control study. *Intern Emerg Med.* 2021;16(7):1945-50. <https://doi.org/10.1007/s11739-021-02632-z> PMID:33484453
- Dean AG, Arner TG, Sunki GG, Friedman R, Lantinga M, Sangam S, Fagan RF. *Epi Info™, a Database and Statistics Program for Public Health Professionals.* Atlanta, USA: Centers for Disease Control and Prevention; 2011.
- De Onis M, Onyango AW. WHO child growth standards. *Lancet.* 2008;371(9608):204. [https://doi.org/10.1016/S0140-6736\(08\)60131-2](https://doi.org/10.1016/S0140-6736(08)60131-2)



- PMid:18207015
24. Kurtz A, Grant K, Marano R, Arrieta A, Grant K Jr., Feaster W, *et al.* Long-term effects of malnutrition on severity of COVID-19. *Sci Rep.* 2021;11(1):14974. <https://doi.org/10.1038/s41598-021-94138-z>  
PMid:34294743
  25. Nuzhat S, Hasan ST, Palit P, Afroze F, Amin R, Alam A, *et al.* Health and nutritional status of children hospitalized during the COVID-19 pandemic, Bangladesh. *Bull World Health Organ.* 2022;100(2):98-107. <https://doi.org/10.2471/BLT.21.285579>  
PMid:35125534
  26. El-Shafie AM, Kasemy ZA, Omar ZA, Alkalash SH, Salama AA, Mahrous KS, *et al.* Prevalence of short stature and malnutrition among Egyptian primary school children and their coexistence with Anemia. *Ital J Pediatr.* 2020;46(1):91. <https://doi.org/10.1186/s13052-020-00855-y>  
PMid:32600418
  27. Shekerdeman LS, Mahmood NR, Wolfe KK, Riggs BJ, Ross CE, McKiernan CA, *et al.* Characteristics and outcomes of children with coronavirus disease 2019 (COVID-19) infection admitted to US and Canadian pediatric intensive care units. *JAMA Pediatr.* 2020;174(9):868-73. <https://doi.org/10.1001/jamapediatrics.2020.1948>  
PMid:32392288
  28. Jin JM, Bai P, He W, Wu F, Liu XF, Han DM, *et al.* Gender differences in patients with COVID-19: Focus on severity and mortality. *Front Public Health.* 2020;8:152. <https://doi.org/10.3389/fpubh.2020.00152>  
PMid:32411652
  29. Parums DV. Editorial: COVID-19 and Multisystem inflammatory syndrome in children (MIS-C). *Med Sci Monit.* 2021;27:e933369. <https://doi.org/10.12659/MSM.933369>  
PMid:34075014
  30. Ahnstedt H, McCullough LD. The impact of sex and age on T cell immunity and ischemic stroke outcomes. *Cell Immunol.* 2019;345:103960. <https://doi.org/10.1016/j.cellimm.2019.103960>  
PMid:31519365
  31. Mertens E, Peñalvo JL. The burden of malnutrition and fatal COVID-19: A global burden of disease analysis. *Front Nutr.* 2021;7:619850. <https://doi.org/10.3389/fnut.2020.619850>  
PMid:33553234
  32. Schaible UE, Kaufmann SH. Malnutrition and infection: Complex mechanisms and global impacts. *PLoS Med.* 2007;4(5):e115. <https://doi.org/10.1371/journal.pmed.0040115>  
PMid:17472433
  33. Pourfridoni M, Abbasnia SM, Shafaei F, Razaviyan J, Heidari-Soureshjani R. Fluid and electrolyte disturbances in COVID-19 and their Complications. *Biomed Res Int.* 2021;2021:6667047. <https://doi.org/10.1155/2021/6667047>  
PMid:33937408
  34. Alfano G, Ferrari A, Fontana F, Perrone R, Mori G, Ascione E, *et al.* Hypokalemia in patients with COVID-19. *Clin Exp Nephrol.* 2021;25(4):401-9. <https://doi.org/10.1007/s10157-020-01996-4>  
PMid:33398605
  35. Noori M, Nejadghaderi SA, Sullman MJ, Carson-Chahhoud K, Ardalan M, Kolahi AA, *et al.* How SARS-CoV-2 might affect potassium balance via impairing epithelial sodium channels? *Mol Biol Rep.* 2021;48(9):6655-61. <https://doi.org/10.1007/s11033-021-06642-0>  
PMid:34392451
  36. Hu W, Lv X, Li C, Xu Y, Qi Y, Zhang Z, *et al.* Disorders of sodium balance and its clinical implications in COVID-19 patients: A multicenter retrospective study. *Intern Emerg Med.* 2021;16(4):853-62. <https://doi.org/10.1007/s11739-020-02515-9>  
PMid:33064253
  37. Gheorghe G, Ilie M, Bungau S, Stoian AP, Bacalbasa N, Diaconu CC. Is there a relationship between COVID-19 and hyponatremia? *Medicina (Kaunas).* 2021;57(1):55. <https://doi.org/10.3390/medicina57010055>  
PMid:33435405
  38. Zimmer MA, Zink AK, Weißer CW, Vogt U, Michelsen A, Priebe HJ, *et al.* Hyponatremia a manifestation of COVID-19: A case series. *A A Pract.* 2020;14(9):e01295. <https://doi.org/10.1213/XAA.0000000000001295>  
PMid:32909725
  39. Ravell J, Chaigne-Delalande B, Lenardo M. X-linked immunodeficiency with magnesium defect, Epstein-Barr virus infection, and neoplasia disease: A combined immune deficiency with magnesium defect. *Curr Opin Pediatr.* 2014;26(6):713-9. <https://doi.org/10.1097/MOP.0000000000000156>  
PMid:25313976
  40. Goshua G, Pine AB, Meizlish ML, Chang CH, Zhang H, Bahel P, *et al.* Endotheliopathy in COVID-19-associated coagulopathy: Evidence from a single-centre, cross-sectional study. *Lancet Haematol.* 2020;7(8):e575-82. [https://doi.org/10.1016/S2352-3026\(20\)30216-7](https://doi.org/10.1016/S2352-3026(20)30216-7)  
PMid:32619411
  41. Trapani V, Rosanoff A, Baniyasi S, Barbagallo M, Castiglioni S, Guerrero-Romero F, *et al.* The relevance of magnesium homeostasis in COVID-19. *Eur J Nutr.* 2021;61(2):625-36. <https://doi.org/10.1007/s00394-021-02704-y>  
PMid:34687321
  42. Stevens JS, Moses AA, Nickolas TL, Husain SA, Mohan S. Increased mortality associated with hypermagnesemia in severe COVID-19 illness. *Kidney 360.* 2021;2(7):1087-94. <https://doi.org/10.34067/KID.0002592021>  
PMid:35368359
  43. Quilliot D, Bonsack O, Jaussaud R, Mazur A. Dymagnesemia in Covid-19 cohort patients: Prevalence and associated factors. *Magnes Res.* 2020;33:114-122. <https://doi.org/10.1684/mrh.2021.0476>  
PMid:33678604
  44. Wang R, He M, Kang Y. Hypophosphatemia at admission is associated with Increased Mortality in COVID-19 Patients. *Int J Gen Med.* 2021;14:5313-5322. <https://doi.org/10.2147/IJGM.S319717>  
PMid:34526806
  45. Sabbagh Y, Giral H, Caldas Y, Levi M, Schiavi SC. Intestinal phosphate transport. *Adv Chronic Kidney Dis.* 2011;18(2):85-90. <https://doi.org/10.1053/j.ackd.2010.11.004>  
PMid:21406292
  46. Levi M, Gratton E, Forster IC, Hernando N, Wagner CA, Biber J, *et al.* Mechanisms of phosphate transport. *Nat Rev Nephrol.* 2019;15(8):482-500. <https://doi.org/10.1038/s41581-019-0159-y>  
PMid:31168066
  47. Biber J, Hernando N, Forster I. Phosphate transporters and their function. *Annu Rev Physiol.* 2013;75:535-50. <https://doi.org/10.1146/annurev-physiol-030212-183748>  
PMid:23398154
  48. Datta BN, Stone MD. Hyperventilation and hypophosphataemia. *Ann Clin Biochem.* 2009;46(Pt 2):170-1. <https://doi.org/10.1258/acb.2008.008199>  
PMid:19225028
  49. Pal R, Ram S, Zohmangaihi D, Biswas I, Suri V, Yaddanapudi LN, *et al.* High prevalence of hypocalcemia in non-severe



- COVID-19 patients: A retrospective case-control study. *Front Med (Lausanne)*. 2021;7:590805. <https://doi.org/10.3389/fmed.2020.590805>  
PMid:33490095
50. Wu MA, Fossali T, Pandolfi L, Carsana L, Ottolina D, Frangipane V, *et al*. Hypoalbuminemia in COVID-19: Assessing the hypothesis for underlying pulmonary capillary leakage. *J Intern Med*. 2021;289(6):861-72. <https://doi.org/10.1111/joim.13208>  
PMid:33411411
51. Loffredo L, Oliva A, Paraniñfi A, Ceccarelli G, Orlando F, Ciacci P, *et al*. An observed association between conjunctivitis and severity of COVID-19. *J Infect*. 2021;83(3):381-412. <https://doi.org/10.1016/j.jinf.2021.06.006>  
PMid:34118275
52. Viana-Llamas MC, Arroyo-Espliguero R, Silva-Obregón JA, Uribe-Heredía G, Núñez-Gil I, García-Magallón B, *et al*. Hypoalbuminemia on admission in COVID-19 infection: An early predictor of mortality and adverse events. A retrospective observational study. *Med Clin (Engl Ed)*. 2021;156(9):428-36. <https://doi.org/10.1016/j.medcle.2020.12.015>  
PMid:33969222
53. Huang J, Cheng A, Kumar R, Fang Y, Chen G, Zhu Y, *et al*. Hypoalbuminemia predicts the outcome of COVID-19 independent of age and co-morbidity. *J Med Virol*. 2020;92(10):2152-8. <https://doi.org/10.1002/jmv.26003>  
PMid:32406952
54. Chen C, Zhang Y, Zhao X, Tao M, Yan W, Fu Y. Hypoalbuminemia an indicator of the severity and prognosis of COVID-19 patients: A multicentre retrospective analysis. *Infect Drug Resist*. 2021;14:3699-710. <https://doi.org/10.2147/IDR.S327090>  
PMid:34526790
55. Aziz M, Fatima R, Lee-Smith W, Assaly R. The association of low serum albumin level with severe COVID-19: A systematic review and meta-analysis. *Crit Care*. 2020;24(1):255. <https://doi.org/10.1186/s13054-020-02995-3>  
PMid:32456658
56. Cai SH, Liao W, Chen SW, Liu LL, Liu SY, Zheng ZD. Association between obesity and clinical prognosis in patients infected with SARS-CoV-2. *Infect Dis Poverty*. 2020;9(1):80. <https://doi.org/10.1186/s40249-020-00703-5>  
PMid:32600411
57. Kompaniyets L, Agathis NT, Nelson JM, Preston LE, Ko JY, Belay B, *et al*. Underlying medical conditions associated with severe COVID-19 illness among children. *JAMA Netw Open*. 2021;4(6):e2111182. <https://doi.org/10.1001/jamanetworkopen.2021.11182>  
PMid:34097050
58. Amin MT, Fatema K, Arefin S, Hussain F, Bhowmik DR, Hossain DS. Obesity, a major risk factor for immunity and severe outcomes of COVID-19. *Biosci Rep*. 2021;41(8):BSR20210979. <https://doi.org/10.1042/BSR20210979>  
PMid:34350941
59. Atef H. Nutritional fat modulation as a non-pharmacological approach to children infected with COVID-19, a challenge in food biochemistry. *Egypt J Chem*. 2022;65(3):689-98. <https://doi.org/10.21608/ejchem.2021.87893.4234>
60. Ritter A, Kreis NN, Louwen F, Yuan J. Obesity and COVID-19: Molecular mechanisms linking both pandemics. *Int J Mol Sci*. 2020;21(16):5793. <https://doi.org/10.3390/ijms21165793>  
PMid:32806722