

# Silver nanoparticles enhance oxidative stress, inflammation, and apoptosis in liver and kidney tissues: Potential protective role of thymoquinone

**Basma Salama**

Mansoura University

**Khalid J. Alzahrani**

Taif University

**Khalid S. Alghamdi**

General Administration of Laboratories and Blood Banks, Ministry of Health

**Osama Al-Amer**

University of Tabuk

**Khalid E. Hassan**

Taif University

**Mohamed A. Elhefny**

National Cancer Institute, Cairo University

**Alaa Albarakati**

Umm Al-Qura University

**Fahad Alharthi**

Taif University

**Hussam A. Althagafi**

Al-Baha University

**Hassan Al Sberi**

National Organization for Drug Control and Research

**Hatem K. Amin**

Galala University

**Maha S. Lokman**

Prince Sattam bin Abdul Aziz University

**Khalaf F. Alsharif**

Taif University

**Ashraf Albrakati**

Taif University

**Ahmed E. Abdel Moneim**

Helwan University

**Rami B. Kassab** (✉ [rami.kassap@yahoo.com](mailto:rami.kassap@yahoo.com))

## Research Article

**Keywords:** Silver nanoparticles, thymoquinone, oxidative damage, inflammation, apoptosis, hepatorenal toxicity

**Posted Date:** July 13th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1828575/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

Silver nanoparticles (AgNPs) are the most common nanomaterials in consumer products. Therefore, it has been crucial to control AgNPs toxicological effects to improve their safety and increase the outcome of their applications. This work investigated the possible protective effect of thymoquinone (TQ) against AgNPs-induced hepatic and renal cytotoxicity in rats. Serum markers of liver and kidney functions as well as liver and kidney oxidative stress status, pro-inflammatory cytokines, apoptosis markers, and histopathology were assessed. TQ reversed AgNPs-induced elevation in serum liver and kidney function markers, including aspartate transaminase, alanine transaminase, urea, and creatinine. Moreover, TQ co-administration with AgNPs alleviates hepatic and renal oxidative insults by decreasing MDA and NO levels with a significant increase in the activity of antioxidant enzymes (superoxide dismutase, catalase, and glutathione recycling enzymes peroxidase and reductase) compared to AgNPs-treated rats. Besides, TQ upregulated hepatic and renal Nrf2 gene expression in AgNPs intoxicated rats. Furthermore, TQ co-administration decreased the hepatic and renal pro-inflammatory mediators represented by, IL-1 $\beta$ , TNF- $\alpha$ , TGF- $\beta$ , and NF- $\kappa$ B levels. Besides, TQ co-administration decreased apoptotic protein (Bax) levels and increased the anti-apoptotic protein (Bcl-2) levels. These findings were confirmed by the histopathological examination of hepatic and renal tissues. Our data affirmed the protective effect of TQ against AgNPs cytotoxicity and proposed a possible mechanism of TQ antioxidant, anti-inflammatory, and anti-apoptotic effects. Consequently, we could conclude that using TQ might control AgNPs toxicological effects, improve their safety, and increase the outcome of their applications.

## Introduction

Recently, engineered nanomaterials have acquired immense attention in technological advancements. The unique nanoparticle (NPs) characteristics are based on their morphology, size, surface charge, and coating. These different characteristics are responsible for their effects on the biological systems [1]. Metallic NPs such as silver, gold, and iron are widely used in medicine and industry owing to their unique thermal, optical, catalytic, and electrical characteristics [2].

Silver nanoparticles (AgNPs) are used in the manufacture of many medical products like antibacterial agents, diagnostic devices, deodorant sprays, wound dressings, surgical instruments, and drug delivery systems [3]. Due to their widespread use, it is mandatory to evaluate their toxicological effects to improve their safety and increase their application [4]. Previous *in vitro* and *in vivo* studies have reported the toxic effect of AgNPs [5–6]. Non-coated AgNPs showed cytotoxicity toward different cell lines such as macrophages, alveolar epithelial cells, hepatocytes, and embryonic kidney cells [7]. AgNPs were proven to enhance reactive oxygen species (ROS), apoptosis, and inflammation; authors reported that these effects are dependent on size, concentration, and route of administration [8–9].

Thymoquinone (TQ) is the primary active ingredient in *Nigella sativa* oil with numerous biological and pharmaceutical activities, including antioxidative, anti-apoptotic, anti-inflammatory, anticancer, renoprotective, and hepatoprotective agent [10–11]. The antioxidant capacity of TQ is due to scavenging

free radicals such as superoxide anion, hydroxyl, hydrogen peroxide, and peroxy nitrite radicals, in addition to enhancing antioxidant enzymes such as superoxide dismutase, catalase, and glutathione peroxidase [12]. Moreover, accumulative studies reported the ability of TQ to inhibit the production of pro-inflammatory cytokines and chemokines such as tumor necrosis factor-alpha (TNF- $\alpha$ ), interleukin-1 beta (IL-1 $\beta$ ), and cyclooxygenase 2 (COX-2) [13].

Based on the previous studies and suggestions, the present work aim is to investigate the possible protective effect of TQ against AgNPs-induced hepatorenal damage by evaluating the redox hemostasis, apoptosis, and inflammatory response in the liver and kidney tissues of rats.

## Materials And Methods

### Nanoparticles

AgNPs (size less than 40 nm) were obtained from Sigma-Aldrich (730793; St. Louis, MO, USA). According to the manufacturer instructions, AgNPs characterization was performed using transmission electron microscopy (TEM), dynamic light scattering (DLS), and UV/Visible spectral analysis to ensure that the monodisperse AgNPs are free from agglomeration with a density of 0.986 g/mL at 25 °C; refractive index  $n_{20/D}$  1.333; fluorescence— $\lambda_{em}$  401 nm) (<https://www.sigmaaldrich.com/EG/en/product/aldrich/730793>).

### Experimental design

Twenty-eight adult male Wistar albino rats aged 3-4 months old and weighed 180–200 grams, (VACSERA, Cairo, Egypt) were kept on a standard diet and tap water *ad libitum* for one week. After acclimatization, animals were randomly divided into four equal groups (n=7):

1- Control group: received normal physiological saline 0.9% NaCl for 28 days.

2- TQ group: treated orally with TQ (10 mg/kg /day) [11].

3- AgNPs group: injected intraperitoneally by AgNPs (50 mg/kg/day) for 28 days [14].

4- TQ-AgNPs group: treated orally with TQ (10 mg/kg/day), and then 2 hours later, the rats were injected intraperitoneally with AgNPs (50 mg/kg/day) for 28 days.

The study protocol was reviewed and approved by the institutional animal care and use committee, Faculty of Science, Helwan University (HU/2020/Z/AEN0120-01), following the European Community Directive (86/609/EEC).

At the end of the experiment, rats were anesthetized by sodium pentobarbital intraperitoneal injection (200 mg/kg) and subjected to a complete autopsy. Blood samples were collected and incubated at room temperature for 10 minutes to clot, then centrifuged at 3000 x *g* for 10 minutes to collect the serum

samples for further analysis. The liver and kidney were immediately dissected and divided into three parts. one part was homogenized with (10% w/v) ice-cold 50 mM Tris-HCl buffer (pH 7.4), and centrifuged at  $3000 \times g$  for 10 min at 4 °C then the supernatant was stored at -20 °C for biochemical analysis. Another part was stored at -80 °C for gene expression analysis. Finally, the third part was preserved in 10% of neutral-buffered formalin for histopathological examination.

### **Assessment of liver and kidney functions**

The levels of alanine transaminase (ALT), aspartate aminotransferase (AST), urea, and creatinine were assayed in serum samples using standard kits (Biodiagnostic, Giza, Egypt) according to the manufacturer's protocol.

### **Oxidative stress assays**

Glutathione (GSH), which is considered a non-enzymatic antioxidant marker, was assessed with the colorimetric method following Elaman's protocol [15] in liver and kidney samples. Malondialdehyde (MDA), which is considered a lipid peroxidation marker, was measured following the method of Ohkawa et al. [16]. Superoxide dismutase (SOD) enzyme activity was assessed depending on the inhibition of the reduction of nitroblue tetrazolium dye by SOD and depending on the  $H_2O_2$  decomposition rate following the method of Aebi [17]. Glutathione peroxidase (GPx) and reductase (GR) enzymes activities were assessed by measuring NADPH oxidation and reduction at 340 nm in the presence of glutathione, following the method of Paglia and Valentine [18] and Factor et al. [19], respectively.

### **Inflammatory marker assessment**

Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and interleukin-1 $\beta$  (IL-1 $\beta$ ) were measured using ELISA kits obtained from Thermo Fisher Scientific, USA for TNF- $\alpha$  (Cat. no. BMS607-3) and IL-1 $\beta$  (Cat. no. BMS6002). Transforming growth factor-beta (TGF- $\beta$ ) was measured by ELISA kits obtained from R&D System, Minneapolis, MN, USA (Cat no: SB100C). Nuclear factor-kappa B (NF- $\kappa$ B, Novus Biologicals, Centennial, CO, USA; Cat. No. NB100-2176) was measured by ELISA kits following the manufacturer's instructions.

### **Apoptotic biomarkers determination**

Bax, which is a pro-apoptotic marker, and Bcl-2, which is an anti-apoptotic marker, were measured using ELISA kits from BioVision, Inc. for (rat Bax; Cat. No.: E4513) and Cusabio (rat Bcl-2; Cat. No.: CSB-E08854r).

### **PCR analysis**

After homogenizing the liver and kidney tissues, total RNA was extracted using RNeasy Plus Mini kits. The extracted RNA (100 ng) was then reverse transcribed into cDNA using a Script<sup>TM</sup> cDNA synthesis kit

(Bio-Rad, CA). qRT-PCR was applied using an Applied Biosystems 7500 Instrument (Applied Biosystems, USA) to estimate the relative expression of the Nrf2 gene using Power SYBR® Green. B-actin was applied as a housekeeping gene.

### Histopathological examination

Fixation of Liver and kidney specimens is done with 10% neutral buffered formalin. After that samples were dehydrated, embedded in paraffin wax, and cut into 5 µm-thick sections. In the next step, liver and kidney sections were deparaffinized, stained with hematoxylin and eosin, and examined under a Nikon Eclipse E200-LED microscope (Nikon Corporation, Tokyo, Japan) for histopathological changes.

### Statistical analysis

Data were expressed as mean values ± SE (standard error), and one-way ANOVA statistically analyzed the significant differences among treatment groups. The criterion for statistical significance was set at  $p < 0.05$  for the biochemical data. All statistical analyses were performed using SPSS statistical version 21 software package (SPSS® Inc., USA).

## Results

### TQ restored AgNPs-induced elevation in serum liver and kidney functions markers

To investigate AgNPs induced liver and kidney injury and the potential antagonistic effect of TQ, serum ALT, AST, urea, and creatinine levels were assessed. As shown in table 1, liver and kidney function markers were significantly ( $P < 0.05$ ) increased in AgNPs treated rats compared to the control group. On the other hand, TQ co-administration significantly reduced both liver and kidney function markers compared to AgNPs treated group. However, TQ alone didn't show any significant effect compared to the control group.

**Table 1: Effect of TQ on AgNPs-induced hepatorenal toxicity**

	Control	TQ	AgNPs	TQ+AgNPs
<b>ALT</b>	48.21 ± 6.29	49.24 ± 5.72	103.81 ± 13.02 <sup>#</sup>	71.19 ± 8.58 <sup>#</sup> \$
<b>AST</b>	74.81 ± 12.23	78.78 ± 13.95	172.51 ± 22.42 <sup>#</sup>	101.37 ± 20.06 <sup>#</sup> \$
<b>Urea</b>	35.17 ± 17	32.68 ± 5.30	78.50 ± 13.29 <sup>#</sup>	56.47 ± 8.15 <sup>#</sup> \$
<b>Creatinine</b>	0.39 ± 0.06	0.37 ± 0.06	0.87 ± 0.15 <sup>#</sup>	0.56 ± 0.09 <sup>#</sup> \$

Data are presented as mean ± SD (n = 7). # and \$ means significance difference,  $p < 0.05$ , from control and AgNPs-treated group, respectively.

## **TQ ameliorates AgNPs-induced oxidative stress in hepatic and renal tissues**

To investigate AgNPs-induced oxidative stress and the potential antioxidant effect of TQ, the levels of MDA, NO, and GSH were evaluated in the liver and kidney (Fig. 1) tissues. AgNPs significantly ( $P<0.05$ ) increased MDA and NO levels and decreased GSH levels in liver and kidney tissues. However, TQ co-administration significantly attenuated AgNPs-induced oxidative stress by decreasing the elevated MDA and NO levels and increasing GSH levels in liver and kidney tissues.

Moreover, the activity of the antioxidant enzymes; SOD, CAT, GPx, and GR were evaluated in the liver and kidney (Fig. 2) tissues. AgNPs significantly ( $P<0.05$ ) suppressed the activity of the hepatic and renal antioxidant enzymes compared to the control group. On the other hand, TQ treatment for AgNPs-intoxicated rats significantly ( $P<0.05$ ) restored the activity of the hepatic and renal antioxidant enzymes compared to the AgNPs-treated group.

To elucidate the molecular mechanism underlying TQ antioxidant effect, mRNA expression of Nrf2 in the hepatic and renal tissue (Fig. 3) was determined. Nrf2 is a transcription factor that plays a vital role in the antioxidant and subsequent anti-inflammatory cellular response. AgNPs induced downregulation of hepatic and renal Nrf2 expression compared to the control group. TQ treatment significantly ( $P<0.05$ ) upregulated Nrf2 expression in AgNPs intoxicated rats. Our gene expression analysis data demonstrated the role of the transcription factor, Nrf2, in the antioxidant and subsequent anti-inflammatory molecular mechanisms of TQ against AgNPs oxidative and inflammatory response.

## **TQ ameliorates AgNPs-induced inflammation in hepatic and renal tissues**

To investigate the anti-inflammatory effect of TQ in response to AgNPs exposure, the pro-inflammatory cytokines levels, IL-1b, TNF- $\alpha$ , and TGF-b were evaluated in the liver and kidney (Fig. 4) tissues. AgNPs significantly ( $P<0.05$ ) increased hepatic and renal IL-1b, TNF- $\alpha$ , and TGF-b levels compared to the control group. On the other hand, TQ co-administration significantly ( $P<0.05$ ) reversed the elevated hepatic and renal pro-inflammatory cytokines levels compared to AgNPs treated group.

To get more information about the mechanism underlying TQ anti-inflammatory effect, NF- $\kappa$ B level was estimated in the liver and kidney (Fig. 5) tissues. NF- $\kappa$ B plays a role in the expression of pro-inflammatory cytokines genes. AgNPs significantly ( $P<0.05$ ) increased hepatic and renal NF- $\kappa$ B levels compared to the control group. In contrast, TQ co-administration significantly decreased hepatic and renal NF- $\kappa$ B levels in AgNPs treated rats. These data elucidate the role of NF- $\kappa$ B in TQ anti-inflammatory effect.

## **TQ ameliorates AgNPs-induced apoptosis in hepatic and renal tissues**

To investigate the anti-apoptotic activity of TQ against cell loss induced by AgNPs, Bax and Bcl2 levels were estimated in liver and kidney tissue (Fig. 6). Bax regulates apoptosis as a pro-apoptotic protein and Bcl-2 as an anti-apoptotic protein. AgNPs significantly ( $P<0.05$ ) increased hepatic and renal Bax levels with decreased Bcl2 levels. Conversely, TQ co-administration ameliorated the apoptotic effect of AgNPs

by restoring Bax level toward control value and significantly increasing Bcl2 level compared to AgNPs treated group.

### **TQ prevents histopathological changes in hepatic and renal tissues following the exposure to AgNPs**

The control and TQ groups showed normal hepatocytes with central vein structure. However, AgNPs-exposed rats exhibited inflamed, degenerated, and apoptotic hepatocytes associated with prominent Kupffer cells (Fig. 7). These histological changes are obviously mitigated following TQ pretreatment (Figs. 7). Moreover, the control and TQ groups showed normal glomeruli with renal tubule structure. Nevertheless, AgNPs-challenged rats exhibited congested glomeruli, degenerated renal tubules accompanied by severe apoptosis, inflammatory cells infiltration into the intratubular spaces, and debris in the lumen of renal tubules (Fig. 7). These pathological alterations were prominently improved following TQ pretreatment (Fig. 7).

## **Discussion**

AgNPs are widely used in numerous industrial and medicinal products such as cosmetics, paints, biosensors, food packaging, wound dressings, and antimicrobial agents [20]. Despite these promising advantages, assessment and decreasing AgNPs toxicity are crucial. In this study, we investigated the protective effect of TQ against AgNPs toxicity by observing the molecular, biochemical, and histopathological changes in liver and kidney tissues.

The small size of the NPs beside the route of administration defines AgNPs biodistribution and target organs [21]. NPs size is one of the essential factors that play a role on the NPs uptake and cellular distribution. Previous research papers confirmed the accumulation of Ag<sup>+</sup> in specific tissues, including lungs, spleen, liver, and kidneys of rats dosed with AgNPs, suggesting that the small “Nano-size” of AgNPs facilitates its accumulation into specific target organs where they may further generate Ag<sup>+</sup> [22]. AgNPs enter the body through inhalation, ingestion, transdermal and parenteral injection. Some studies reported that liver and kidney tissues are the main targets of AgNPs accumulation in rats after oral administration and intravenous injection [23–25]. AgNPs accumulation leads to tissue injury, oxidative stress, inflammation, and apoptosis [26].

In our study, liver tissue injury induced by AgNPs treatment is evidenced by elevated serum levels of liver function biomarkers, AST and ALT, indicating cell membrane leakage, membrane permeability disturbance, and structural integrity loss. AgNPs-induced kidney tissue injury is evidenced by increased serum urea and creatinine levels, indicating a disruption in kidney clearance function. Moreover, histopathological examination of liver and kidney tissues confirms that AgNPs induced hepatorenal injury

Oxidative stress could result from an imbalance between pro-oxidants and antioxidants [27]. In this study, AgNPs-induced oxidative stress is confirmed by increased generation of NO and MDA and the exhaustion of antioxidant enzymes, namely SOD, CAT, GPx, and GR, along with the decreased level of GSH. Several



reports support our findings as they reported the potency of AgNPs in the induction of oxidative stress by increasing ROS and reducing the activity of antioxidant enzymes [28–30]. Li et al. [31] suggested an explanation for enhanced ROS generation by confirming the accumulation of AgNPs in the mitochondria, leading to reduced mitochondrial membrane potential and mitochondrial impotence that may increase ROS production.

Recently it's been well accepted that AgNPs induce oxidative stress either by targeting the gene expression of antioxidant enzymes or directly interacting with enzyme surfaces, especially SOD and CAT [32]. Wei et al. [33] proved the surface interaction between AgNPs and SOD and CAT, forming an NPs-protein enzyme complex. Subsequently, NPs-enzyme interaction leads to conformational changes and enzyme activity inhibition. Moreover, AgNPs exhibit a strong affinity for GSH thiol groups; this could explain the decreased GSH levels in liver and kidney cells [34].

Our data showed similar results of oxidative stress status in kidney tissues. These results are in accordance with previous reports [35, 24]. These studies suggested that AgNPs accumulate in kidney tissues and other target organs such as liver tissues, then dissociation into Ag<sup>+</sup> might lead to oxidative stress and inflammation.

TQ, the main constituent of *Nigella Sativa*, has been investigated for its immunomodulatory, antioxidant, anti-inflammatory, and anticarcinogenic effects [36]. These advantageous effects prompted us to assume that TQ could control AgNPs toxicity. Therefore, in this work, we investigated the possible hepatic and renal protective effect of TQ against AgNPs-induced oxidative, inflammatory, and apoptotic effects.

TQ co-administration significantly improved the levels of serum liver and kidney function markers in AgNPs intoxicated rats. These results confirmed the protective effect of TQ. The antioxidant property is the main effect of TQ. Previous studies attributed the capability of TQ to protect against several conditions such as hepatic cancer and ischemia-induced liver injury to oxidative stress suppression as TQ could decrease NO synthesis through direct suppression of nitric oxide synthases (iNOS and eNOS) besides TQ's ability to normalize the depleted GSH level and antioxidant enzymes [37]. Moreover, previous studies confirm the potency of TQ to protect against oxidative stress induced by several hepatotoxic agents such as CB 1954 [38], cisplatin [39], cyclophosphamide [40], Aflatoxins [41] and carbon tetrachloride [42] through decreasing lipid peroxidation and NO synthesis along with increased expression of antioxidant enzymes. Our data revealed that TQ could relieve the oxidative stress induced by AgNPs in hepatic and renal tissues. TQ decreased the levels of MDA and NO and increased the levels of GSH and antioxidants enzyme activities in AgNPs intoxicated rats.

To get more information about the ability of TQ to ameliorate AgNPs-induced oxidative stress, the expression level of Nrf2 was estimated using qRT-PCR in hepatic and renal tissues. Nrf2 is a key transcription factor that regulates the transcription of the antioxidant enzymes by binding to the antioxidant response element (ARE) in the DNA [43]. Our data revealed that AgNPs intoxication leads to downregulation of Nrf2 expression in both hepatic and renal tissues. These findings may clarify the

reason behind the decreased activity of antioxidant enzymes in liver and kidney tissues. On the other hand, TQ treatment normalized the expression of hepatic and renal Nrf2 in AgNPs intoxicated rats. Our findings reveal the ability of TQ to mitigate the oxidative stress induced by AgNPs in hepatic and renal tissues.

Once the oxidative balance is impaired, the inflammatory response and mitochondria-related cell death will follow. Increased ROS production promotes inflammatory signals that lead to increased production of pro-inflammatory cytokines such as IL-1 $\beta$  and TNF- $\alpha$  and TGF- $\beta$ . These pro-inflammatory cytokines are commonly used as a marker for toxicant-induced inflammation [44]. Herein, we evaluated the inflammatory effect of AgNPs and the possible anti-inflammatory effect of TQ. Our data confirmed the inflammatory impact of AgNPs represented by increased hepatic and renal pro-inflammatory cytokines, IL-1 $\beta$ , TNF- $\alpha$ , and TGF- $\beta$  and increased NF- $\kappa$ B level. The activation of NF- $\kappa$ B regulates the expression of pro-inflammatory cytokines genes [45].

In line with our results, previous reports concluded the inflammatory effect of different sizes of AgNPs on various organs. Shehata et al. [14] demonstrated that oral administration of AgNPs resulted in increased inflammatory cytokines in liver and kidney tissues of rats. Choia et al. [46] confirmed the hepatotoxic and inflammatory effect of AgNPs in adult zebrafish. Other research groups revealed the toxic and inflammatory effect of AgNPs manifested by increased inflammatory cytokines such as IL-1 $\beta$  and TNF- $\alpha$  and TGF- $\beta$  in mice [47], guinea pig [48], and in vitro studies on hepatic cells [49].

On the other hand, our data showed that TQ pretreatment prevented AgNPs induced hepatic and renal inflammatory responses, as evidenced by the decreased pro-inflammatory cytokines and NF- $\kappa$ B levels. It is well documented that TQ possesses an anti-inflammatory effect that protects several disorders such as asthma, arthritis, diabetes, and neuroinflammatory diseases. The noted anti-inflammatory effect of TQ is proved by decreased pro-inflammatory cytokines and suppression of the NF- $\kappa$ B pathway [50].

Bax regulates apoptosis as a pro-apoptotic protein and Bcl-2 as an anti-apoptotic protein; therefore, we investigated the effect of AgNPs and TQ on these protein levels. Our data support previous reports [51, 5] about the apoptotic effect of AgNPs in both hepatic and renal tissues manifested by elevated Bax and decreased Bcl-2 levels. Besides, immunohistochemistry revealed increased hepatic and renal caspase-3 levels. These data demonstrate the mechanism of AgNPs-induced apoptosis. Our results showed that TQ significantly reversed the changes in Bax and Bcl-2, confirming the protective anti-apoptotic effect of TQ against AgNPs toxicity. These results agreed with previous studies that show the anti-apoptotic effect of TQ [52–53].

## Conclusion

Our study concluded the protective effect of TQ against AgNPs induced hepatorenal toxicity in rats. Moreover, as shown in Fig. 8, we proposed the possible mechanism underlying TQ antioxidant, anti-inflammatory, and anti-apoptotic effects. Based on the recorded results, we recommend the usage of TQ to control AgNPs toxicity, improve AgNPs safety and increase the outcomes of AgNPs application.

# Declarations

**Acknowledgements:** The authors thank the Taif University Researchers Supporting Program (Project number: TURSP-2020/151), Taif University, Saudi Arabia, for supporting this work.

## Author Contribution

Conceptualization and supervision: A.E.A., R.B.K., and A.A.; Animal treatments, molecular and biochemical methodologies were performed by B.S., K.J.A., K.S.A. O.A., K.E.H., and M.A.E.; Histological methodology and investigation conducted by F.A., H.A.A., H.A., and A.S.F.; data analysis, software, data curation, and visualization were performed by H.K.A., M.S.L., K.F.A., and A.A.; writing-reviewing and editing manuscript performed by B.S., A.E.A., and R.B.K. All authors participated in the design and interpretation of the study and approved the final manuscript.

**Data Availability:** All relevant data are within the paper.

## Declarations

**Ethics Approval:** The study protocol was reviewed and approved by the institutional animal care and use committee, Faculty of Science, Helwan University (HU/2020/Z/AEN0120-01), following the European Community Directive (86/609/EEC).

**Consent to Participate:** Not applicable.

**Consent for Publication:** Not applicable.

**Competing Interests:** The authors declare no competing interests.

# References

1. Carbone M, Donia DT, Sabbatella G, Antiochia R (2016) Silver nanoparticles in polymeric matrices for fresh food packaging. *Journal of King Saud University-Science* 28 (4):273-279
2. Khan I, Saeed K, Khan I (2019) Nanoparticles: Properties, applications and toxicities. *Arabian Journal of Chemistry* 12 (7):908-931
3. Benn TM, Westerhoff P (2008) Nanoparticle silver released into water from commercially available sock fabrics. *Environmental science & technology* 42 (11):4133-4139
4. Gaillet S, Rouanet J-M (2015) Silver nanoparticles: their potential toxic effects after oral exposure and underlying mechanisms—a review. *Food and chemical toxicology* 77:58-63
5. Akter M, Sikder MT, Rahman MM, Ullah AA, Hossain KFB, Banik S, Hosokawa T, Saito T, Kurasaki M (2018) A systematic review on silver nanoparticles-induced cytotoxicity: Physicochemical properties and perspectives. *Journal of advanced research* 9:1-16


6. Sung JH, Ji JH, Song KS, Lee JH, Choi KH, Lee SH, Yu IJ (2011) Acute inhalation toxicity of silver nanoparticles. *Toxicology and Industrial Health* 27 (2):149-154
7. Makama S, Kloet SK, Piella J, van den Berg H, de Ruijter NC, Puntjes VF, Rietjens IM, van den Brink NW (2018) Effects of systematic variation in size and surface coating of silver nanoparticles on their in vitro toxicity to macrophage RAW 264.7 cells. *Toxicological sciences* 162 (1):79-88
8. Sriram MI, Kalishwaralal K, Barathmanikanth S, Gurunathani S (2012) Size-based cytotoxicity of silver nanoparticles in bovine retinal endothelial cells. *Nanoscience Methods* 1 (1):56-77
9. Gurunathan S, Park JH, Han JW, Kim J-H (2015) Comparative assessment of the apoptotic potential of silver nanoparticles synthesized by *Bacillus tequilensis* and *Calocybe indica* in MDA-MB-231 human breast cancer cells: targeting p53 for anticancer therapy. *International journal of nanomedicine* 10:4203
10. Gökce EC, Kahveci R, Gökce A, Cemil B, Aksoy N, Sargon MF, Kısa Ü, Erdoğan B, Güvenç Y, Alagöz F (2016) Neuroprotective effects of thymoquinone against spinal cord ischemia-reperfusion injury by attenuation of inflammation, oxidative stress, and apoptosis. *Journal of Neurosurgery: Spine* 24 (6):949-959
11. Al Aboud D, Baty RS, Alsharif KF, Hassan KE, Zhery AS, Habotta OA, Elmahallawy EK, Amin HK, Abdel Moneim AE, Kassab RB (2021) Protective efficacy of thymoquinone or ebselen separately against arsenic-induced hepatotoxicity in rat. *Environmental Science and Pollution Research* 28 (5):6195-6206
12. Hamdy NM, Taha RA (2009) Effects of *Nigella sativa* oil and thymoquinone on oxidative stress and neuropathy in streptozotocin-induced diabetic rats. *Pharmacology* 84 (3):127-134
13. El Mezayen R, El Gazzar M, Nicolls MR, Marecki JC, Dreskin SC, Nomiyama H (2006) Effect of thymoquinone on cyclooxygenase expression and prostaglandin production in a mouse model of allergic airway inflammation. *Immunology letters* 106 (1):72-81
14. Shehata AM, Salem F, El-Saied EM, El-Rahman A, Sahar S, Mahmoud MY, Noshay PA (2022) Evaluation of the Ameliorative Effect of Zinc Nanoparticles against Silver Nanoparticle-Induced Toxicity in Liver and Kidney of Rats. *Biological trace element research* 200 (3):1201-1211
15. Ellman GL (1959) Tissue sulfhydryl groups. *Arch Biochem Biophys* 82 (1):70-77
16. Ohkawa H, Ohishi N, Yagi K (1979) Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 95 (2):351-358
17. Aebi H (1984) Catalase in vitro. *Methods in enzymology* 105:121-126
18. Paglia DE, Valentine WN (1967) Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med* 70 (1):158-169
19. Factor VM, Kiss A, Woitach JT, Wirth PJ, Thorgeirsson SS (1998) Disruption of redox homeostasis in the transforming growth factor-alpha/c-myc transgenic mouse model of accelerated hepatocarcinogenesis. *The Journal of biological chemistry* 273 (25):15846-15853
20. Xu L, Li X, Takemura T, Hanagata N, Wu G, Chou LL (2012) Genotoxicity and molecular response of silver nanoparticle (NP)-based hydrogel. *Journal of nanobiotechnology* 10 (1):1-11

21. Tsiola A, Pitta P, Callol AJ, Kagiorgi M, Kalantzi I, Mylona K, Santi I, Toncelli C, Pergantis S, Tsapakis M (2017) The impact of silver nanoparticles on marine plankton dynamics: Dependence on coating, size and concentration. *Science of The Total Environment* 601:1838-1848
22. Kim HR, Park YJ, Da Young Shin SMO, Chung KH (2013) Appropriate in vitro methods for genotoxicity testing of silver nanoparticles. *Environmental health and toxicology* 28
23. Jiménez-Lamana J, Laborda F, Bolea E, Abad-Álvaro I, Castillo JR, Bianga J, He M, Bierla K, Mounicou S, Ouerdane L (2014) An insight into silver nanoparticles bioavailability in rats. *Metallomics* 6 (12):2242-2249
24. Wen H, Dan M, Yang Y, Lyu J, Shao A, Cheng X, Chen L, Xu L (2017) Acute toxicity and genotoxicity of silver nanoparticle in rats. *PLOS ONE* 12 (9):e0185554
25. Su C-K, Liu H-T, Hsia S-C, Sun Y-C (2014) Quantitatively profiling the dissolution and redistribution of silver nanoparticles in living rats using a knotted reactor-based differentiation scheme. *Analytical chemistry* 86 (16):8267-8274
26. Tiwari DK, Jin T, Behari J (2011) Dose-dependent in-vivo toxicity assessment of silver nanoparticle in Wistar rats. *Toxicology mechanisms and methods* 21 (1):13-24
27. Al-Brakati A, Albarakati AJA, Daabo H, Baty RS, Salem FEH, Habotta OA, Elmahallawy EK, Abdel-Mohsen DM, Taha H, Akabawy A (2020) Neuromodulatory effects of green coffee bean extract against brain damage in male albino rats with experimentally induced diabetes. *Metabolic brain disease* 35 (7):1175-1187
28. Patlolla AK, Hackett D, Tchounwou PB (2015) Silver nanoparticle-induced oxidative stress-dependent toxicity in Sprague-Dawley rats. *Molecular and cellular biochemistry* 399 (1):257-268
29. Zhang X-F, Shen W, Gurunathan S (2016) Silver nanoparticle-mediated cellular responses in various cell lines: an in vitro model. *International journal of molecular sciences* 17 (10):1603
30. AshaRani P, Low Kah Mun G, Hande MP, Valiyaveetil S (2009) Cytotoxicity and genotoxicity of silver nanoparticles in human cells. *ACS nano* 3 (2):279-290
31. Li L, Cui J, Liu Z, Zhou X, Li Z, Yu Y, Jia Y, Zuo D, Wu Y (2018) Silver nanoparticles induce SH-SY5Y cell apoptosis via endoplasmic reticulum-and mitochondrial pathways that lengthen endoplasmic reticulum-mitochondria contact sites and alter inositol-3-phosphate receptor function. *Toxicology Letters* 285:156-167
32. Fang W, Chi Z, Li W, Zhang X, Zhang Q (2019) Comparative study on the toxic mechanisms of medical nanosilver and silver ions on the antioxidant system of erythrocytes: from the aspects of antioxidant enzyme activities and molecular interaction mechanisms. *Journal of nanobiotechnology* 17 (1):1-13
33. Liu W, Worms I, Slaveykova VI (2020) Interaction of silver nanoparticles with antioxidant enzymes. *Environmental Science: Nano* 7 (5):1507-1517
34. Zhou Y-T, He W, Lo YM, Hu X, Wu X, Yin J-J (2013) Effect of silver nanomaterials on the activity of thiol-containing antioxidants. *Journal of agricultural and food chemistry* 61 (32):7855-7862

35. Albrahim T (2020) Silver nanoparticles-induced nephrotoxicity in rats: the protective role of red beetroot (*Beta vulgaris*) juice. *Environmental Science and Pollution Research* 27 (31):38871-38880
36. Noorbakhsh M-F, Hayati F, Samarghandian S, Shaterzadeh-Yazdi H, Farkhondeh T (2018) An overview of hepatoprotective effects of thymoquinone. *Recent patents on food, nutrition & agriculture* 9 (1):14-22
37. Abd-Elbaset M, Arafa E-SA, El Sherbiny GA, Abdel-Bakky MS, Elgendy ANA (2017) Thymoquinone mitigate ischemia-reperfusion-induced liver injury in rats: a pivotal role of nitric oxide signaling pathway. *Naunyn-Schmiedeberg's archives of pharmacology* 390 (1):69-76
38. Talib WH, AbuKhader MM (2013) Combinatorial effects of thymoquinone on the anticancer activity and hepatotoxicity of the prodrug CB 1954. *Scientia pharmaceutica* 81 (2):519-530
39. Al-Malki AL, Sayed AAR (2014) Thymoquinone attenuates cisplatin-induced hepatotoxicity via nuclear factor kappa- $\beta$ . *BMC complementary and alternative medicine* 14 (1):1-8
40. Laskar AA, Khan MA, Rahmani AH, Fatima S, Younus H (2016) Thymoquinone, an active constituent of *Nigella sativa* seeds, binds with bilirubin and protects mice from hyperbilirubinemia and cyclophosphamide-induced hepatotoxicity. *Biochimie* 127:205-213
41. Nili-Ahmadabadi A, Tavakoli F, Hasanzadeh G, Rahimi H, Sabzevari O (2011) Protective effect of pretreatment with thymoquinone against Aflatoxin B1 induced liver toxicity in mice. *Daru: journal of Faculty of Pharmacy, Tehran University of Medical Sciences* 19 (4):282
42. Mansour MA (2000) Protective effects of thymoquinone and desferrioxamine against hepatotoxicity of carbon tetrachloride in mice. *Life sciences* 66 (26):2583-2591
43. Kim J, Cha Y-N, Surh Y-J (2010) A protective role of nuclear factor-erythroid 2-related factor-2 (Nrf2) in inflammatory disorders. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* 690 (1-2):12-23
44. Wang Z, Xia T, Liu S (2015) Mechanisms of nanosilver-induced toxicological effects: more attention should be paid to its sublethal effects. *Nanoscale* 7 (17):7470-7481
45. Zhang H, Sun S-C (2015) NF- $\kappa$ B in inflammation and renal diseases. *Cell & Bioscience* 5 (1):1-12
46. Choi JE, Kim S, Ahn JH, Youn P, Kang JS, Park K, Yi J, Ryu D-Y (2010) Induction of oxidative stress and apoptosis by silver nanoparticles in the liver of adult zebrafish. *Aquatic Toxicology* 100 (2):151-159
47. Park E-J, Bae E, Yi J, Kim Y, Choi K, Lee SH, Yoon J, Lee BC, Park K (2010) Repeated-dose toxicity and inflammatory responses in mice by oral administration of silver nanoparticles. *Environmental toxicology and pharmacology* 30 (2):162-168
48. Korani M, Rezayat SM, Bidgoli SA (2013) Sub-chronic dermal toxicity of silver nanoparticles in guinea pig: special emphasis to heart, bone and kidney toxicities. *Iranian journal of pharmaceutical research: IJPR* 12 (3):511
49. Gaiser BK, Hirn S, Kermanizadeh A, Kanase N, Fytianos K, Wenk A, Haberl N, Brunelli A, Kreyling WG, Stone V (2013) Effects of silver nanoparticles on the liver and hepatocytes in vitro. *Toxicological sciences* 131 (2):537-547

50. Darakhshan S, Pour AB, Colagar AH, Sisakhtnezhad S (2015) Thymoquinone and its therapeutic potentials. *Pharmacological Research* 95:138-158
51. Piao MJ, Kang KA, Lee IK, Kim HS, Kim S, Choi JY, Choi J, Hyun JW (2011) Silver nanoparticles induce oxidative cell damage in human liver cells through inhibition of reduced glutathione and induction of mitochondria-involved apoptosis. *Toxicology Letters* 201 (1):92-100
52. El-Ghany A, Ragwa M, Sharaf NM, Kassem LA, Mahran LG, Heikal OA (2009) Thymoquinone triggers anti-apoptotic signaling targeting death ligand and apoptotic regulators in a model of hepatic ischemia reperfusion injury. *Drug discoveries & therapeutics* 3 (6)
53. Sethi G, Ahn KS, Aggarwal BB (2008) Targeting nuclear factor- $\kappa$ B activation pathway by thymoquinone: role in suppression of antiapoptotic gene products and enhancement of apoptosis. *Molecular cancer research* 6 (6):1059-1070


## Figures



**Figure 1**


The effect of TQ on AgNPs-induced oxidative stress in the liver and kidney tissues. Data are presented as mean  $\pm$  SD (n = 7). # and \$ means significance difference,  $p < 0.05$ , from control and AgNPs-treated group, respectively.






**Figure 2**

The effect of TQ on the suppressed activity of antioxidant enzymes induced by AgNPs in the liver and kidney tissues. Data are presented as mean  $\pm$  SD (n = 7). # and \$ means significance difference,  $p < 0.05$ , from control and AgNPs-treated group, respectively.



**Figure 3**

The effect of TQ on the mRNA expression of Nrf2 in the liver and kidney tissues following the exposure to AgNPs. Data are presented as mean  $\pm$  SD (n = 7). # and \$ means significance difference,  $p < 0.05$ , from control and AgNPs-treated group, respectively.



**Figure 4**

The protective effect of TQ against AgNPs-induced inflammation response (IL-1b, TNF-  $\alpha$ , and TGF-b) in the liver and kidney tissues. Data are presented as mean  $\pm$  SD (n = 7). # and \$ means significance difference, p < 0.05, from control and AgNPs-treated group, respectively.




Figure 5

the effect of TQ co-administration on NF-κB level in the liver and kidney tissues. Data are presented as mean  $\pm$  SD (n = 7). # and \$ means significance difference,  $p < 0.05$ , from control and AgNPs-treated group, respectively.





Figure 6

The protective effect of TQ on the apoptotic markers (Bax and Bcl-2) in response to AgNPs exposure in the liver and kidney tissues. Data are presented as mean  $\pm$  SD (n = 7). # and \$ means significance difference,  $p < 0.05$ , from control and AgNPs-treated group, respectively.

**Figure 7**

TQ's protective impacts on the histological alterations in liver and kidney tissues following exposure to AgNPs ( $\times 400$ ).



**Figure 8**

Schematic diagram shows the protective effect of TQ against AgNPs induced hepatorenal toxicity. A. illustrates the possible mechanism of TQ antioxidant effect. B. illustrates the possible mechanism of TQ's anti-inflammatory effect. C. illustrates the possible anti-apoptotic effect of TQ anti-apoptotic effect. Blue arrows represent the effect of AgNPs, and red arrows represent the effect of TQ. Up directed arrows (↑) means significant increase, and down directed arrows (↓) means significant decrease.