

**Role of *Mycobacterium avium* sub species *paratuberculosis* in persistent diarrhoea in Egyptian buffaloes**  
**Shohanda M.K.\*; Samia A. Ahmed\*\*; Soliman S.M.\*\*; El-Gedawy, A.A.\* and Y.F. Elnaker\*\*\***

\*Animal Health Research Institute, Dokki-Bacteriology Dept. Tuberculosis Unit

\*\*Faculty of veterinary Medicine, Cairo University, Dept. of Internal medicine and Infectious disease.

\*\*\*Faculty of Veterinary Medicine, New valley, Assiut University, Animal medicine department (Infectious diseases)

Received in 18/5/2016

Accepted in 12/6/2016

### Abstract

Paratuberculosis or John's disease (JD) is a chronic and incurable granulomatous enteric disease affecting cattle, buffaloes, sheep, goats and the other ruminants caused by *M. avium* subspecies *paratuberculosis* (MAP). In this study, A total of 300 buffaloes (240 clinically diseased animals and 60 apparently healthy animals) were examined for the presence of clinical signs of John's disease including incurable chronic diarrhoea, intermittent firstly then intense and continuous which is not responding to treatment, emaciation and progressive weakness. Fecal and serum samples (each of 300) were collected from the examined buffaloes housed in 3 Egyptian Governorates (Sharkia, Kalyoubia and Damietta). Fecal samples were collected then examined according to the pooling procedure and decontaminated by Hexa decylpyridinium chloride solution (HPC 0.9%) prior to culturing on Herrold's Egg Yolk Medium (HEYM). MAP was isolated from 34 of the 60 pooled fecal samples tested (57%). Fecal smears were examined using Ziehl – Neelsen stain (ZN) for the presence of acid fast bacilli revealing 29 fecal smears (48%) of 60 fecal smears were positive. ELISA was conducted on serum samples to detect antibodies against MAP, 212 (71%) of serum samples were positive for antibodies against MAP. Molecular confirmation by PCR IS900 assay was carried out using specific primers directly on fecal sample, Out of the 60 pooled fecal samples, 45 pools (75%) were positive. This study aimed to throw a light on paratuberculosis in Egyptian buffaloes as there is lack of data about this disease in Egypt

**Key words:** Paratuberculosis - John's disease- buffaloes-Persistent diarrhea.

### 1. Introduction

Paratuberculosis or John's disease (JD) is a chronic and incurable granulomatous enteric disease affecting cattle, buffalo, sheep, goats and the other ruminants caused by *M. avium* subspecies *paratuberculosis* (MAP). The infection is typically characterized by long incubation period followed by chronic progressive diarrhoea, decrease in milk yield, submandibular edema, anemia, loss of weight leading to

cachexia and finally to death (Cocito *et al.*, 1994, Harris and Barletta, 2001). Animals are infected by ingestion of food and water contaminated by feces of infected animals. The intrauterine transmission of Map may occur in buffaloes and could represent an important source of infection (Belo-Reis *et al.*, 2016). The incidence of subclinical cases shedding organisms intermittently may be as high as 15%. Calves are susceptible but don't show

signs until adulthood. JD is ranked as a notifiable disease in many countries (**Kennedy, 2011**). The prolonged incubation period of the disease (2-10 years) and the fact that many animals may harbor the organism without ever showing clinical symptoms have made the diagnosis of latent cases difficult. Meanwhile, the lack of any specific chemotherapeutic agent, together with the vaccination drawbacks has made the disease particularly difficult to control. (**Mckenna *et al.*, 2006 and Barry *et al.*, 2011**).

Studies are carried out mostly on cattle, so little is known about paratuberculosis in buffalo. Hence a more comprehensive study of MAP in buffaloes is of a great value to facilitate the design of prevention and control programs.

Although the diagnosis of JD remains difficult; there is a wide range of tests for diagnosis of Johne's disease. Bacteriologic fecal culture remains the most definitive test for identification of both subclinically and clinically infected animals. This fecal culture is time consuming, requiring a long incubation period of 8 to more than 16 weeks for bacterial recovery. This causes problems for producers who need to make quick decisions for the purchase or replacement animals or to cull the infected animals from their herds (**Shaymaa *et al.*, 2015**). The sensitivity of the Ziehl Neelsen test has been estimated at 49% (**Zimmer *et al.*, 1999**) in clinically affected animals. The specificity of the ZN-test was estimated at 83% (**Ris *et al.*, 1988**) in fecal culture-negative animals from paratuberculosis free herds.

Concerning control programs, ELISA is widely used screening assay for the detection of antibodies against MAP in blood or milk because of their advantages like easy performance, high capacity and low costs. PCR technique is a more rapid test which could be an alternative to the time consuming traditional cultural method (**Douarre *et al.*, 2010**). PCR test has led to increase specificity and sensitivity of detection of low shedders. (**Collins *et al.*, 1993; Whittington, 2001; Mason *et al.*, 2001 and Behr and Collins, 2010**).

*Mycobacterium avium* sub species *paratuber-*

*culosis* may have public health importance as clinical similarities have been observed between Johne's disease in animals and Crohn's disease (CD) in humans and both viable MAP and MAP genetic material have been found in some patients diagnosed with Crohn's disease despite almost 100 years of investigation, conflicting data still occur and the responsibility of MAP as the etiological agent of CD was proven in some reports and denied by others. *M. avium* subspecies *paratuberculosis* has been isolated from Crohn's patient tissue worldwide. In spite of that, many laboratories are still reporting their inability to culture it at all, as some human isolates may take up to 6 years to grow even under standard culture and decontamination conditions (**Collins *et al.*, 2000 and Parrish *et al.*, 2008**).

This study aimed to throw light on paratuberculosis in Egyptian buffaloes as there is lack of data about this disease in Egypt and it was achieved through isolation of MAP on HEYM's medium from pooled fecal samples, identification of the MAP organism by Ziehl Neelsen stain of fecal smears, detection of antibodies to MAP in serum samples by indirect ELISA and IS900 PCR technique.

## 2. Materials and Methods

### 2.1. Animals

Animals were suffering from incurable chronic diarrhoea, intermittent firstly then intense and continuous which is not responding to treatment, emaciation and progressive weakness and apparently healthy animals went through general clinical examination parameters including rectal temperature, heart rate, respiratory rate and body condition score.

### 2.2. Samples

A total of 600 random samples (300 fecal samples and 300 serum samples) were collected from 3 Governorate (Sharkia, Kalyoubia and Damietta). Fecal samples were collected and either directly processed or kept frozen in -20 C till processing

Blood samples were drawn in serum vacutainer without anticoagulant and kept in ice

box waiting to be sent to the laboratory. Separated serum was kept frozen in (-20 °C) till examination.

### **2.3. Pooling, Decontamination and cultivation procedures**

Fecal samples were examined on the basis of a strategic pooling procedure by weighing 2 g of feces from each of 5 animals into sterile mortar. A small amount of sterile water (approximately 5ml) was added to make it possible to thoroughly mix the 5 samples with a sterile pestle.

After mixing, a 3g aliquot (2 g feces and 1 ml water) was weighed out and transferred to a second sterile mortar (Kalis *et al.*, 2000). The pooled fecal sample was decontaminated with 0.9% Hexadecylpyridinium Chloride solution (HPC) for 24 h. 3 g of feces were added to a 50mL sterile tube containing 30ml of a 0.9% HPC. This suspension was manually mixed by shaking and vortexing and let in vertical position for 30 min at room temperature to allow the precipitation and sedimentation of big particles. Approximately 20ml of the upper portion of the supernatant was transferred to another 50ml sterile tube. Tubes were placed in vertical position in dark place for 24 h at room temperature. Then, centrifugation of the tubes for 10 min by 3000 rpm. (Fernández-Silva *et al.*, 2011). Supernatant was discarded and Herrold's egg yolk medium (HEYM) slants which are supplemented with mycobactin J (Prepared Culture Media) were inoculated with 300 µl of the decontaminated pellet. The slants were incubated at 37 °C in horizontal position for 1 week with the caps loosened to allow absorption and evaporation of residual moisture on the surface of the medium. Then, caps were tightened and the tubes were returned to vertical position and incubated for 8-16 weeks and checked at 1-2 weeks intervals. The tubes were being checked for growth of gray white rough colonies. Smears were taken from the suspected samples for microscopical examination (Fernández-Silva *et al.*, 2011).

### **2.4. Identification of MAP by Ziehl – Neelsen staining of direct fecal smear**

Fecal smears were prepared and air dried for 10 minutes then heat fixed at 60-70°C for 2hrs. Each slide was flooded with carbol fuchsin and heated gently until a small amount of steam rises. Boiling was avoided. The slides were left for 5-10 minutes then rinsed with tap water. Acid alcohol was added for at least 1-2 minutes then slides were rinsed thoroughly with tap water. The slides were flooded with Methylene blue as a counterstain and were left for 1-2 minutes then rinsed with tap water and air dried. The slides were examined under oil immersion lens. (Australian and New Zealand standard diagnostic procedures, 2010).

### **2.5. Indirect ELISA for detection of MAP antibodies in serum samples.**

The presence of MAP antibodies was detected in serum samples using the PARACHEK2 (Mycobacterium paratuberculosis test kit), (Prionics AG Wagistrasse 27a-8952 Schlieren-Zurich, Switzerland) Results were interpreted according to manufacturer's instructions.

### **2.6. Molecular MAP confirmation by using PCR IS900**

DNA extraction was carried out using QIAamp DNA stool Mini Kit (Qiagen, Germany) according to manufacturer's instructions. The PCR was carried out using primers to detect the presence of MAP-specific IS900 DNA (Table-1).

**Table (1).** Nucleotide sequences of PCR primers

Target gene	Oligonucleotide sequence (5'-3')	Expected amplicon size (bp)
IS900	F:CCG CTA ATT GAG AGA TGC GAT TGG	229 (Vary <i>et al.</i> ,1990)
	R:AAT CAA CTC CAG CAG CGC GGC CTC G	

The amplification product was analyzed using gel electrophoresis on 1.5% agarose gel (Sambrook *et al*, 1989) and was subsequently visualized by UV illumination after ethidium bromide staining and was photographed by a gel documentation system.

**3. Results**


**3.1. Isolation of Mycobacterium avium sub-species paratuberculosis**

The culture results for all pooled fecal samples tested during this study are shown in table (1). MAP was isolated from 34 of the 60 pooled samples tested (57%). After 16 weeks of incubation on Herrold's media containing egg yolk slope, Typical colonies of MAP were observed; very small, convex (hemispherical), soft, non-mucoid and initially colourless and translucent. For some slants, it took about nine months to observe typical colonies

**Table (1).** Results of isolation of fecal culture of MAP on HEYM

Governorate	No. of pooled fecal samples		Total No. of pooled fecal samples	No. and percentage(*) of Positive culture		Total No. and percentage of positive culture
	Diseased animals	Apparently healthy		Diseased	Apparently healthy	
Sharkia	16	4	20	12 (60%)	1(5%)	13(65%)
Kalyoubia	16	4	20	9 (45%)	1 (5%)	10 (50%)
Damietta	16	4	20	11 (55%)	0 (0%)	11(55%)
<b>Total</b>	<b>48</b>	<b>12</b>	<b>60</b>	<b>32</b>	<b>2</b>	<b>34 (57%)</b>

(\*) percentage was calculated according to the total number of the pooled fecal samples.



**Figure (1).** Results of isolation of foecal culture of MAP on HEYM

### 3.2. Microscopic examination of Ziehl–Neelsen stained fecal smears

Microscopic appearance of MAP showed red acid alcohol-fast ZN staining bacilli. Staining

of direct fecal smears revealed 29 (48%) of 60 pooled samples were positive as shown in table (2)

**Table (2).** Results of ZN fecal smear staining.

Governorate	No. of pooled samples		No. and Percentage(*) of positive ZN stained fecal smear		Total No. of Positive ZN stained fecal smear and Percentage (%)
	Diseased animals	Apparently healthy animals	Diseased animals	Apparently healthy animals	
Sharkia	16	4	9 (45%)	1 (5%)	10 (50%)
Kalyoubia	16	4	8 (40%)	0 (0%)	8 (40%)
Damietta	16	4	10 (50%)	1 (5%)	11 (55%)
<b>Total</b>	<b>48</b>	<b>12</b>	<b>29</b>		<b>29(48%)</b>

(\*) percentage was calculated according to the total number of the pooled fecal samples.

### 3.3. ELISA assay results

ELISA was carried out individually to all serum samples of culture-positive animals and culture-negative animals. The ELISA results are shown in table (3). In total, out of 300 se-

rum samples of clinically diseased animals and apparently healthy animals tested using indirect ELISA for detection of antibodies against

**Table (3).** Results of indirect ELISA test

Governorate	No. of tested serum samples		Total No. of serum samples	No. and percentage (*) of Positive ELISA		Total No. and percentage of positive ELISA
	Diseased	Apparently healthy		Diseased animals	Apparently healthy	
Sharkia	80	20	100	70 (70%)	0 (0%)	70 (70%)
Kalyoubia	80	20	100	60 (60%)	1 (1%)	61 (61%)
Damietta	80	20	100	80 (80%)	1 (0%)	81 (81%)
<b>Total</b>	<b>240</b>	<b>60</b>	<b>300</b>	<b>210</b>	<b>2</b>	<b>212 (71%)</b>


(\*) percentage was calculated according to the total number of the pooled fecal samples.

**3.4. Amplification of MAP IS900 gene by using PCR**

Out of 60 pooled fecal samples, 45 pools (75%) were positive.

All positive culture fecal pools were PCR positive, although not every PCR positive sam-

ple was positive culture pool. Results of PCR is summarized in table (4). Representative positive samples are shown in photo (1)



**Photo (1):** Showing results of IS900 PCR amplicons of some fecal samples

**M:** Marker 100pb

**L1:** Control positive

**L2, L3 and L4:** Positive samples

**L5:**Control Negative

**L6, L7 and L8:** Negative samples

**Table (4).** Results of IS900 PCR

Governorate	No. of examined samples		Total No. of Examined samples	Positive PCR (N)		Total No. of positive PCR
	Diseased animals	Apparently healthy animals		Diseased animals	Apparently healthy animals	
<b>Sharkia</b>	16	4	20	14 (70%)	1 (5%)	15 (80%)
<b>Kalyoubia</b>	16	4	20	12(60%)	2 (10%)	14 (70%)
<b>Damietta</b>	16	4	20	15 (75%)	1 (5%)	16 (80%)
<b>Total</b>	48	12	60	41	4	<b>45 (75%)</b>

(\*) percentage was calculated according to the total number of the serum samples in each governorate.

#### 4. Discussion

Johne's disease (JD) is caused by the bacterium *Mycobacterium avium* subspecies *paratuberculosis*. Infections normally affect ruminants but have also been seen in a variety of non-ruminant species, including rabbits, foxes, birds, horses, dogs, and non-human primates have been infected also. JD primarily affects the intestine of all wild and domestic ruminants including cattle, goats, sheep, camels, llamas, giraffes, bison, buffaloes, deer and antelopes (Whittington *et al.* 2011). The disease has been also recorded in a variety of non-ruminant species including rabbits, foxes, birds, horses, donkeys, dogs, cats, pigs, and non-human primates (Beard *et al.* 2001; Glanemann *et al.* 2008 and Stief *et al.* 2012).

Clinical signs of JD including wasting, chronic intermittent diarrhoea, emaciation, decrease in net body weight and milk yield. Animals are infected by ingestion of food and water contaminated by feces of infected individuals. Shedding of infectious agent usually occurred from clinical cases in spite of the incidence of subclinical cases shedding organisms intermittently may be as high as 15%.

Studies are carried out mostly on cattle, So little is known about Paratuberculosis in buffaloes, Hence, a more comprehensive knowledge of *Mycobacterium avium* subsp. *paratuberculosis* in buffaloes is of great value to facilitate the design of prevention and control programs. Understanding the dynamics of the disease is the key element of control as there's lack of practical therapeutic approaches and lack of a vaccine that prevents transmission and the complexity and difficulty of the on farm control strategies hinders the prevention of infection (Behr and Collins, 2010). More recently evidence has accumulated for an association of MAP with Crohn's disease in humans, adding to the pressure on animal health authorities to take precautions to control paratuberculosis.

JD has been reported in almost every continent. Studies from various parts of the world have shown the widespread distribution of the disease. It is mainly a problem reported in Aus-

tralia, Canada, Argentina, USA, Mexico, Brazil, New Zealand, Denmark, Belgium, Norway, Switzerland, Netherlands, France, Spain, Germany, England, Scotland, Ireland, Italy, Greece, Thailand, India, Japan, Saudi Arabia, Iran, Egypt, Morocco and South Africa. However, Sweden and some states in Australia are the only regions that claimed to be free from the disease (Bauerfeind *et al.* 1996; Moreira *et al.* 1999; Pavlik *et al.* 1999; Machackova *et al.* 2004; Salem *et al.* 2005; Palmer *et al.* 2005 and carvalho *et al.* 2009).

Fecal culture is the most practical mean of confirming infection in herd screening programs. Although fecal culture is technically difficult and time-consuming (Van schaik *et al.*, 2003).

The pooling of fecal samples from more than one animal for culture is a logical way to reduce the cost of detection of MAP at herd level and this observation is in accordance with that mentioned by (Whittington *et al.*, 2001; Van Schaik *et al.*, 2003 and Kalis *et al.*, 2004). Current diagnostic tests consider the culture as the gold standard method (Paolicchi *et al.*, 2003).

One of the goals of this study was to evaluate diagnostic methods for MAP diagnosis in Egyptian buffaloes. The tests used to diagnose MAP in buffaloes were carried out in parallel to each other. Pooled fecal samples were cultured using Herrold's egg yolk medium. (HEYM) supplemented with mycobactin J as the egg based media (LJM & HEYM) are the first choice for culture of clinical and environmental samples (Moravoka *et al.*, 2012). The result of the culture was 34 positive slants of the 60 pooled fecal samples tested (57%) which were collected from three Egyptian governorates (Sharkia, Kalyoubia and Damietta). After 16 weeks of incubation on herrold's media containing egg yolk slope, Positive slants show typical colonies of MAP; very small, convex (hemispherical), soft, non mucoid & initially colorless & translucent. Colonies become bigger, more raised, opaque, off-white cream to buff or beige coloured as incubation

continues. Negative slants should not be disposed until 10 months pass as some slants, took about 9 months to observe typical colonies. and our result was almost near the findings of **Asmaa (2014)**, as she reported the result of culture was 34 of the 50 pooled samples tested (68%). The low sensitivity of fecal culture could be due to intermittent shedding of causative agent in the feces (**Stabel, 1997**) or due to substantial decrease in the bacterial load during specimen decontamination protocol (**Reddacliff et al., 2003**) and as reported by **Visser (1999)**, fecal culture may not necessarily give positive results in animals shedding low numbers of MAP in their feces.

Stained fecal smears using Z.N stain were examined microscopically and revealed that 29 (48%) of the 60 pooled samples were positive as it showed the presence of acid fast bacilli at microscopic examination .

Despite the noticeable low figures of sensitivity and specificity of ELISA (Enzyme linked immunosorbent assay ), It is considered as the method of choice for the detection of J.D - positive herds. This is due to the suitability of samples collection, rapid laboratory turnaround time. Low cost, and possibility to test a large number of samples in a short time (**Muskens et al., 2003; Böttcher and Gangl 2004; Collins, 2011**). In this study, Serum samples of diseased individuals and apparently healthy animals were examined by indirect ELISA for detection of antibodies against MAP where 212 (71%) out of 300 serum samples were positive and 88 serum samples were negative at a percentage of (29%) . All individual animals which were positive ELISA for MAP were in the positive pooled fecal cultures. (**Weber et al., 2009**) reported that the result of positive ELISA. serum samples were 57%. The use of ELISA in longstanding infected herds reveals reasonable sensitivity as the antibodies become more abundant in later stages of the disease (**McKenna et al. 2006**). The specificity of the test is markedly increased by the absorption with sonicates from other mycobacteria like *Mycobacterium phlei* in the so-called absorbed ELISA where specificity reaches 99-100% and

sensitivity reaches 53-55% (**Böttcher and Gangl 2004 and Shin et al., 2008**).

**El Gedawy et al., (2015)** recommended the use of both fecal culture method and indirect ELISA as laboratory methods in diagnosis and control programmes of paratuberculosis.

In this study, Molecular MAP confirmation was done by Conventional PCR technique using specific IS900 primers. Recently, the PCR technique as a powerful tool in the microbiological diagnosis of different pathogens led to its extensive application in the diagnosis of JD .The IS900 is commonly used as an abundant reference marker for the molecular detection of *M. avium* subspecies *paratuberculosis* (**Herthnek et al. 2006**). Many publications considered the PCR assay as rapid and powerful tool to amplify DNA of MAP and as a confirmatory test (**Whittington, 2001 and Mason et al., 2001**). In this study, From Each governorate, 20 pooled fecal samples representing apparently healthy cases and pooled fecal samples representing the diseased cases were examined. Out of the 60 pooled fecal samples, 45 pools (75%) were positive; as the DNA amplification gave a fragment size of 229 on 1.5% agarose electrophoresis) (photo (1)). All positive cultured fecal pools were PCR positive although not all PCR positive samples were culture positive. Reduced sensitivity of PCR results can be due to inefficient extraction of mycobacterial DNA from fecal samples, particularly in low count microbial loaded samples and /or as a consequence of the presence of PCR inhibitors. (**Thornton and Passen 2004**). Also, it is known that fecal samples are very difficult to process and IS900 sequence was detected also in mycobacterial strains other than *M. avium* subspecies *paratuberculosis*. (**Englund et al., 2002**). So the specificity of IS900 diagnostic PCR has been questioned since some reports indicated the presence of IS900-like sequences in mycobacterial species other than MAP.

It can be concluded that

- Some points should be considered as the intermittent shedding of the microorganism and sometimes, the low numbers of MAP in feces

and the loss of the microorganism during the exhausting decontamination protocol .

- Identification through ZN staining for fecal smears revealed that 29(48%) out of the 60 pooled samples were positive.

- By ELISA, out of the 300 serum samples, 212 (71%) of tested serum samples were positive. It's unfair to say that it is a true sensitivity; As ELISA in this study was performed on all individual animals serum samples unlike fecal culture was performed according to pooling procedure and the only way to determine that the animal is infected with MAP or not is by reculturing it individually.

Further investigations and more studies are needed to understand the dynamics and the epidemiology of the disease, helping in the development of the on farm control strategies to prevent the infection and limit this hidden threat as it cause important economic losses in ruminants losses due to examination, treatment costs, loss of milk production and animals culling.

Veterinary authorities should work on offering data about the prevalence of Paratuberculosis on the national scale for the different animal species.

## 5. Reference

**Asmaa, E.G. (2014).** Isolation and Molecular characterization of *Mycobacterium paratuberculosis*. Ph. D. thesis, Department of Microbiology, Faculty of Veterinary Medicine, Alexandria University.

**Australian and New Zealand Standard Diagnostic Procedures, April (2002).** (<http://www.scahls.org.au/Procedures/Documents/ANZSDP/jd1.pdf>)

**Barry, C.; Corbett, D.; Bakker, D.; Andersen, P.; McNair, J. and Strain, S. (2011).** The Effect of *Mycobacterium avium* Complex Infections on Routine *Mycobacterium bovis* Diagnostic Tests. Veterinary Medicine International, 2011, 145092-145100.

**Bauerfeind, R.; Benazzi, S.; Weiss, R.; Schli-**

**esser, T.; Willems, H. and Baljer, G. (1996).** Molecular characterization of *Mycobacterium paratuberculosis* isolates from sheep, goats, and cattle by hybridization with a DNA probe to insertion element IS900. Journal of Clinical Microbiology, 34, 1617–1621.

**Beard, P.M.; Daniels, M.J. and Henderson, D. (2001).** Paratuberculosis infection of non-ruminant wildlife in Scotland. J. Clin. Microbiol., 39: 1517 – 1521.

**Behr, M.A. and Collins, D.M. (2010).** Paratuberculosis: Organism, Disease, Control. CAB International 2010, Wallingford, Oxfordshir, UK

**Belo-Reis,, A.S.; Salvarani, F.M.; Brito, M.F.; Fonseca, A.A.; Silva, N.S.; Silveira, J.A.S.; Reis, J.K.P.; Silva, J.B.; Oliveira, C.M.C. and Barbosa, J.D. (2016).** Intrauterine Infection by *Mycobacterium avium* subsp. *paratuberculosis* in buffalo (*Bubalus bubalis*). J Vet Sci. Med. Diagn 2016, 5: 10

**Böttcher, J. and Gangl, A. (2004).** *Mycobacterium avium* ssp. *Paratuberculosis* combined serological testing and classification of individual animals and herds. Journal of veterinary medicine. B, Infectious diseases and veterinary public health, 51, 443–448

**Carvalho, I.A.; Silva, A.; Jr., Campos, V.E. and Moreira, M.A. (2009).** Short communication: detection of *Mycobacterium avium* subspecies *paratuberculosis* by polymerase chain reaction in bovine milk in Brazil. Journal of Dairy Science, 92, 5408–5410

**Cocito, C.; Gilot, P.; Coene, M.; De Kesel, M.; Poupard, P. and Vannuffel, P. (1994).** Paratuberculosis. Clin. Microbiol. Rev., 7(3): 328-345.

**Collins, M.T. (2011).** Diagnosis of paratuberculosis. Veterinary Clinics of North America: Food Animal Practice, 27, 581–91

- Collins, M.T.; Lisby, G.; Moser, C.; Chicks, D.; Christensen, S.; Reichelderfer, M.; Hoiby, N.; Harms, B.A.; Thomsen, O.O.; Skibsted, U. and Binder, V. (2000).** Results of Multiple Diagnostic Tests for *Mycobacterium avium* subsp. *paratuberculosis* in Patients with Inflammatory Bowel Disease and in Controls. *Journal of Clinical Microbiology*, 38, 4373–4381.
- Collins, M.T. and Sockett, D.C. (1993).** Accuracy and economic of the USDA-licensed enzyme linked immunosorbent assay for bovine paratuberculosis. *J. Am Vet Med Assoc.*, 203: 1456 – 1463.
- Douarre, P.E.; Cashman, W.; Buckley, J.; Coffey, A. and O Mahony, J.M. (2010).** Isolation and detection of *Mycobacterium avium* subsp. *paratuberculosis* (*Map*) from cattle in Ireland using traditional culture and molecular based methods. *Gut Pathogens*. 2:11.
- El-Gedawy, A.A.; Y.F. El-naker and M.A. Elbeskawy (2015).** Diagnosis of Johne's disease in cattle in some Egyptian governorates with special reference to culture and ELISA methods. *Egy. J. of Appl. Sci.*, 30(4) 2015
- Englund, S. Bolske, G, Johansson, K.E. (2002).** An IS900-like sequence found in a *Mycobacterium* sp. other than *Mycobacterium avium* subsp. *paratuberculosis*. *FEMS Microbiol Lett*. 2002 Apr 9; 209(2): 267-71.
- Fernández-Silva, J.A.; Abdulmawjood, A. and Bülte, M. (2011).** Diagnosis and Molecular Characterization of *Mycobacterium avium* subsp. *paratuberculosis* from Dairy Cows in Colombia. *Journal of Veterinary Medicine International*, Vol. 2011, Article ID 352561, 12 pages
- Glanemann, B.; Schonenbrucher, H.; Bridger, N.; Abdulmawjood, A.; Neiger, R. and Bulte, M. (2008).** TI—Detection of *Mycobacterium avium* subspecies *paratuberculosis*-specific DNA by PCR in intestinal biopsies of dogs. *Journal of Veterinary Internal Medicine*, 4, 1090–1094.
- Harris, N.B. and Barletta, R.G. (2001).** “*Mycobacterium avium* subsp. *paratuberculosis* in veterinary medicine,” *Clinical Microbiology Reviews*, 14(3): 489–512.
- Herthnek, D. and G. Boelske (2006).** New PCR systems to confirm real time PCR detection of *Mycobacterium avium* subsp. *paratuberculosis* *J.BMC Microbiology*.6: 87.
- Kalis, C.H.J.; Collins, M.T.; Barkema, H.W. and Hesselink, J.W. (2004).** Certification of herds as free of *Mycobacterium paratuberculosis* infection: actual pooled faecal results versus certification model predictions. *Preventive Veterinary Medicine* 65, 189–204.
- Kalis, C.H.J.; Hesselink, J.W.; Barkema, H.W. and Collins, M.T. (2000).** Culture of strategically pooled bovine fecal samples as a method to screen herds for paratuberculosis. *J Vet Diagn Invest*, 12: 547– 551.
- Kennedy, D.J. and Benedictus, G. (2001).** Control of *Mycobacterium avium* subsp. *paratuberculosis* infection in agricultural species. *Rev. Sci. Tech.*, 20: 151–179.
- Machackova, M.; Svastova, P.; Lamka, J.; Parmova, I.; Liska, V.; Smolik, J.; Fischer, O.A. and Pavlik, I. (2004).** Paratuberculosis in farmed and free-living wild ruminants in the Czech Republic (1999–2001). *Veterinary Microbiology*, 101, 225–234
- Mason O.; Marsh I.B. and Whittington R.J. (2001).** Comparison of immunomagnetic bead separation-polymerase chain reaction and fecal culture for the detection of *Mycobacterium avium* sub species *paratuberculosis* in feces. *Aust Vet. J.*, 79: 497-500.
- McKenna, S.L.; Keefe, G.P.; Tiwari, A.; Van Leeuwen, J. and Barkema, H.W.**

- (2006). Johne's disease in Canada part II: disease impacts, risk factors, and control programs for dairy producers. *Canadian Veterinary Journal*, 47, 1089–1099.
- Moravkova, M.; Babak, V.; Kralova, A.; Pavlik, I. and Slana, I. (2012).** Culture- and Quantitative IS900 Real-Time PCR-Based Analysis of the Persistence of *Mycobacterium avium subsp. paratuberculosis* in a Controlled Dairy Cow Farm Environment. *Appl. Environ. Microbiol.*, 78: 6608–6614.
- Moreira, A.R.; Paolicchi, F.; Morsella, C.; Zumarraga, M.; Cataldi, A.; Fabiana, B.; Alicia, A.; Piet, O.; Van Soolingen, D. and Isabel, R.M. (1999).** Distribution of IS900 restriction fragment length polymorphism types among animal *Mycobacterium avium subsp. paratuberculosis* isolates from Argentina and Europe. *Veterinary Microbiology*, 70, 251–259.
- Muskens, J.; Mars, M.H.; Elbers, A.R.; Van Maanen, K. and Bakker, D. (2003).** The results of using faecal culture as confirmation test of paratuberculosis-seropositive dairy cattle. *Journal of veterinary medicine. B, Infectious diseases and veterinary public health*, 50, 231–234
- Palmer, M.V.; Stoffregen, W.C.; Carpenter, J.G. and Stabel, J.R. (2005).** Isolation of *Mycobacterium avium subsp paratuberculosis* (Map) from Feral Cats on a Dairy Farm with Map-infected Cattle. *Journal of Wildlife Diseases*, 41, 629–635.
- Paolicchi, F.A.; Zumarraga, M.J.; Gioffre, A.; Zamorano, P.; Morsella, C.; Verna, A.; Cataldi, A.; Alito, A. and Romano, M. (2003).** Application of different methods for the diagnosis of paratuberculosis in dairy cattle herd in Argentina. *J. Vet. Med. B. Infect. Dis. Vet. Public health*. 50(1): 20 – 26.
- Parrish, N.M.; Radcliff, R.P.; Brey, B.J.; Anderson, J.L.; Clark, D.L.; Jr., Koziarczyk, J.J.; Ko, C.G.; Goldberg, N.D.; Brinker, D.A.; Carlson, R.A.; Dick, J.D. and Ellingson, J.L. (2008).** Absence of *Mycobacterium avium subsp. Paratuberculosis* in Crohn's patients. *Inflammatory bowel disease*, 15, 556-558.
- Pavlik, I.; Horvathova, A.; Dvorska, L.; Bartl, J.; Svastova, P.; du Maine, R. and Rychlik, I. (1999).** Standardisation of restriction fragment length polymorphism analysis for *Mycobacterium avium* subspecies *paratuberculosis*. *Journal of Microbiological Methods*, 38, 155–167.
- Reddacliff, L.A.; Vadali, A. and Whittington, R.J. (2003).** The effect of decontamination protocols on the numbers of sheep strain *Mycobacterium avium subsp. paratuberculosis* isolated from tissues and faeces. *Vet. Microbiol.*, 95 (4): 271–282.
- Ris, D.R.; Hamel, K.L. and Ayling, J.M. (1988).** The detection of *Mycobacterium paratuberculosis* in bovine faeces by isolation and the comparison of isolation with the examination of stained smears by light microscopy. *N. Z. Vet. J.* 36, 112-114.
- Salem, M.; Zeid, A.A.; Hassan, D.; El-Sayed A. and Zschoeck, M. (2005).** Studies on Johne s disease in Egyptian Cattle. *Journal of Vet. Med. Series B* 52: 134 – 137.
- Sambrook, J.; Fritsch, E.F. and Maniatis, E. (1989).** *Molecular cloning. A laboratory manual*. Vol 1., Cold spring Harbor Laboratory press, New York.
- Shaymaa Abd El-Malek, Kh. F. Mohammed, R.Soliman, M. El Hariri (2015).** Comparison of the sensitivity of different primers used for identification of *Mycobacterium avium subsp. paratuberculosis* isolates recovered from clinically suspected cases. *Int. J. of Sci. Tech. and management* Vol. 4, issue 09.

- Shin, S.J.; Cho, D. and Collins, M.T. (2008).** Diagnosis of Bovine Paratuberculosis by a Novel Enzyme-Linked Immunosorbent Assay Based on Early Secreted Antigens of *Mycobacterium avium* subsp. *paratuberculosis*. *Clinical and Vaccine Immunology*, 15, 1277–1281
- Stabel J.R. (1997).** An improved method for cultivation of *Mycobacterium paratuberculosis* from bovine fecal samples and comparison to three other methods. *J Vet Diagn Invest* 9:375–380.
- Stief, B.; Mobius, P.; Turk, H.; Horugel, U.; Arnold, C. and Pohle, D. (2012).** [Paratuberculosis in a miniature donkey (*Equus asinus f. asinus*)]. *Berliner und Munchener Tierarztliche Wochenschrift*, 125, 38–44.
- Thornton, C.G. and Passen, S. (2004).** Inhibition of PCR amplification by phytic acid, and treatment of bovine fecal specimens with phytase to reduce inhibition. *Journal of Microbiological Methods*, 59, 43–52.
- Van Schaik, G.; Stehman, M.S.; Schukken, H.Y.; Rossiter, R.C. and Shin, J.S. (2003).** Pooled fecal culture sampling for *Mycobacterium avium* Subsp. *paratuberculosis* at different herd sizes and prevalence. *J. VET Diagn Invest*. 15(3): 233-241.
- Vary, P.H.; Andersen, P.R. and Green, E. (1990).** Use of highly specific DNA probes and the polymerase chain reaction to detect *Mycobacterium paratuberculosis* in Johne's disease. *J. Clin. Microbiol.* 28, 933–937.
- Visser, I. (1999).** Reproducibility of a fecal culture method for *Mycobacterium paratuberculosis*. *Proceedings of the 6<sup>th</sup> International colloquium on paratuberculosis, Australia*, pp. 41.
- Weber, M.F.; J. Verhoeff; G. Van Schaik, and C. Van Maanen (2009).** Evaluation of Ziehl-Neelsen stained faecal smear and ELISA as tools for surveillance of clinical paratuberculosis in cattle in the Netherlands. *Prev. Vet. Med.*, 92: 256-266.
- Whittington, R.J.; Marsh, I.B.; Saunders, V.; Grant, I.R.; Juste, R.; Sevilla, I.A.; Manning, E.J.; and Whitlock, R.H. (2011).** Culture phenotypes of genomically and geographically diverse *Mycobacterium avium* subsp. *paratuberculosis* isolates from different hosts. *Journal of Clinical Microbiology*, 49, 1822–1830.
- Whittington, R.J. and Sergeant, E.S.G. (2001).** Progress towards understanding the spread, detection and control of *Mycobacterium avium* subsp. *Paratuberculosis* in animal populations. *Aust Vet. J.*, 4: 267 – 278.
- Yue, R.; Liu, C.; Barrow, P.; Liu, F.; Cui, Y.; Yang, L. and Zhou, X. (2016).** The isolation and molecular characterization of *Mycobacterium avium* subsp. *paratuberculosis* in Shandong province, China. *Gut Pathogens*, 8, 9. <http://doi.org/10.1186/s13099-016-0092-6>
- Zimmer, K.; Drager, K.G.; Klawonn, W. and Hess, R.G. (1999).** Contribution to the diagnosis of Johne's disease in cattle. Comparative studies on the validity of Zeihl-Neelsen staining, faecal culture and a commercially available DNA probe test in detecting *Mycobacterium paratuberculosis* in feces from cattle. *Zentralbl Vet. Med. B.*, 46(2): 137 – 140.